

A database of Holocene sediment cores for England

Short Communication

A.J. Suggitt^{1,2,*}, R.T. Jones³, C.J. Caseldine³, B. Huntley⁴, J.R. Stewart⁵, S.J. Brooks⁶, E. Brown⁷, D. Fletcher^{1,5}, P.K. Gillingham⁵, J. Larwood⁷, N.A. Macgregor⁷, B. Silva⁷, Z. Thomas³, R.J. Wilson² & I.M.D. Maclean^{1,2,*}.

¹ Environment and Sustainability Institute, University of Exeter.

² School of Biosciences, University of Exeter.

³ Department of Geography, University of Exeter.

⁴ School of Biological and Biomedical Sciences, University of Durham.

⁵ Conservation Ecology and Environmental Sciences Group, University of Bournemouth.

⁶ Natural History Museum, London.

⁷ Natural England, Nobel House, 17 Smith Square, London.

* Joint corresponding authors email: a.suggitt@exeter.ac.uk , i.m.d.maclean@exeter.ac.uk

tel: +44 (0) 1326 255968.

Abstract

Extracting sediment cores for palaeoecological and archaeological investigations has occurred extensively across England since the early 20th Century. Surprisingly, there has been comparatively little collation of these valuable publications and potential sources of data; for example, a search on the European Pollen Database (1st Aug 2014 edition) found just 118 core sites for the whole of Great Britain. Here, using a combination of systematic meta-searching and knowledge of the unpublished ('grey') literature, we have assembled a database of some 763 sediment cores documented across 273 scientific studies. The majority of these (>90%) were sediment cores upon which pollen analyses had been performed, but other types of evidence, such as plant macrofossil (preserved plant remains large enough to be visible without a microscope) and faunal records were also identified. We are making this database publically available, in the hope that it will assist further investigations into vegetation history, palaeoecology, and environmental change.

Keywords

Pollen diagram, radiocarbon, recolonisation, carbon dating, climatic change.

Introduction

Palaeoecological studies have been undertaken in England, and more widely in Great Britain and Ireland, since the early 20th century. Seminal early syntheses were provided by Jessen (1949) of late Quaternary deposits and the floral-history of Ireland, Godwin (1956) of the history of British flora (a subsequent more widely used edition was published in 1975) and Pennington (1969) of the history of British vegetation. Many early studies were focused specifically on aspects of vegetation history, rather than the use of pollen and macrofossils to infer climatic and/or anthropogenic impacts. For instance, there was a sporadic series of papers under the overall title of 'Studies in the post-glacial history of British vegetation' in a range of journals including *Philosophical Transactions of the Royal Society* and *New Phytologist*, the first appearing in 1938 (Godwin & Clifford 1938). By the 1970s there was a growing understanding of vegetation history that included studies of plant colonisation in the early Holocene and thereafter, including work from areas with specialised flora such as Upper Teesdale (Turner et al. 1973). These works, although pioneering, suffer from one major drawback in that they preceded the widespread use of radiocarbon dating. They were instead dated by correlation with Godwinian pollen zones, constructed on a very small sample of radiocarbon dates. Nevertheless, the availability of these studies, of which the work of Turner et al. (1973) is a prime example, offers an opportunity to evaluate potential sites for re-analysis with more modern overall approaches. As the spatial coverage of sediment cores increased, so they became used for a number of broad scale studies seeking to elucidate patterns of change within the flora, especially for key species and not just for England but for the whole of the British Isles (Birks 1989) and for Europe as a whole (Huntley & Birks 1983). Thus, the analytical strength of combining palaeoecological records was demonstrated in the literature from an early stage.

From the mid-1970s onwards, studies tended to be more hypothesis driven, looking at specific problems. These included human impacts on the landscape from the Mesolithic through to recent historical changes, with a special interest in the earliest agriculture, or climatic change, particularly during the Lateglacial. Greater numbers of radiocarbon dates were often obtained for individual sites, resulting in better-dated records than previously, and sequences were often analysed with higher temporal resolution, although with a tendency to focus on specific periods rather than to examine the entire Holocene. As a result, we have a good knowledge of vegetation history across a range of regions from which suitable sites have been analysed, although the data are in some cases rather isolated in time and space, especially where there are severe restrictions on the availability of lake or peat sediments. This is particularly true of England when compared to Wales, Scotland or Ireland. In these latter countries, the greater proportion of uplands and the glacial history result in greater numbers of sites where sediments suitable for analysis have accumulated.

Despite this rich history of palaeoecological research, there has been no concerted effort at the national scale to document and collate these records. The European Pollen Database (EPD, Fyfe et al. 2009), and more recently the European Modern Pollen Database (Davis et al. 2013), both represent invaluable resources for researchers working at the European level, while a catalogue of pollen diagrams has recently been compiled for Ireland (Mitchell et al. 2013). However, a search of the EPD for English records (1st Aug 2014 edition) found just 118 core sites available for analysis, for the whole of Britain. Here, we present the findings of an extensive meta-search and examination of the grey literature to determine as many sediment coring sites in England as possible (Supplementary Table 1).

Search Protocol

To ensure we captured as many published studies involving sediment coring as possible, we conducted a systematic meta-search of the scientific literature using the search engines 'Web of Science' and Google Scholar. We used a number of relevant search terms (Supplementary Table 2) to identify suitable studies, while also pursuing literature cited within these studies. Because the searches in Google Scholar generated a large number of returns (e.g. more than 75,000 documents for 'Holocene' + 'pollen'), we sorted these returns by 'relevance', restricting our searches to the 100 most relevant. We also sourced a number of additional articles known to the author team and/or outside the peer-reviewed literature. Studies were included in the database if they documented the findings from at least one sediment core taken during the Holocene (although in practice almost no cores from our search returns had temporal coverage entirely limited to the Late Glacial, see also Figure 1c). The resulting database of studies (Supplementary Table 2) is provided in the supplementary information.

Geographical coverage of the database

A total of 273 palaeoecological studies were identified. These studies represented 763 locations from which palaeoecological data were obtained. The majority of these (>90%) were sediment cores upon which pollen analyses had been performed, but other types of evidence, such as plant macrofossil (preserved plant remains large enough to be visible without a microscope) and faunal records were also identified. The number of individual samples examined from each core ranges widely (from 3 to almost 200). Very few sites had more than a small number of plant macrofossil or faunal records.

The locations of coring sites, together with the number of dated samples associated with these sites, are shown in Figure 1. There are large variations in the number of sites in different regions of the country, largely in line with a combination of the availability of deposits, adopted research questions and researcher interests. There are particularly abundant data from North East England (especially the North Pennines), Cumbria, the North York Moors and the Humber Estuary area. There is also a moderate amount of data from the Welsh Marches, Dartmoor, Exmoor and areas in the Breckland and Fens. Data from elsewhere are scarce or entirely non-existent, principally due to a lack of suitable deposits rather than untapped resources.

Temporal coverage of the database

Although we searched for studies from both the Pleistocene and the Holocene epochs, most studies considered only the Holocene, with only thirty studies providing data from the terminal stage of the Pleistocene (~15.0 – 11.7 ka cal BP; Figure 1c). The longest pollen record was estimated as beginning 17,400 ka cal BP (Bennett 1988), although the validity of the older dates from this site (Saham Mere, Norfolk) has been questioned. The rise in temporal coverage of dated sediment cores coincides with the beginning of the Holocene, amelioration in the climate, and the beginning of peat formation. There is a notable Mid-Holocene peak in the coverage of dated cores at 4-3 kYBP (Figure 1c). Note that dating of cores was not adjusted for year of publication, e.g. a date of 0 kYBP as reported in a 1964 article was not adjusted by 50 years relative to our baseline of 2014.

Concluding remarks and relevance for biological conservation

Our database brings together over 80 years of palaeoecological research in England. This research extends to 763 coring locations and documented across 273 studies. By bringing these studies together, we hope to facilitate further research by providing an internally consistent, simple means of establishing the research history of a particular fieldsite, a region of interest, or indeed England as a whole. Of note is that the database is a reflection of: a) the spatial availability of suitable deposits, which is variable across England, and b) the priorities of researchers, which have undoubtedly changed over time (see Introduction).

Conservation biology is increasingly conducted at the landscape scale to better inform policy goals (Lawton et al. 2010), and the development of statistical and spatial-analytic techniques continues apace. Both these research directions are data hungry, and we hope that by making this database publicly available, further deductions into vegetation history, palaeoecology and other priority areas of research will be facilitated. For

example, we imagine these studies will provide a useful steer for initiatives relating to the management or restoration of degraded habitats (Seddon et al. 2014), and the question of what state a particular habitat should be restored to. Of note here is the rich history of human disturbance in England, from simple hunting and/or harvesting to fire management and direct vegetation manipulation (examples include Peglar et al. 1989; Bennett et al. 1990). Therefore, it follows that conservation goals designed to return ecosystems to 'natural' states- as perceived in the modern era- must be balanced with pragmatism over anthropogenic drivers of habitat status (and change).

Like many countries with a history of deforestation and intensivised agriculture, English habitats are now fragmented and in variable condition. It has been demonstrated repeatedly that, in order to respond successfully to the multitude of environmental changes underway, species require these habitats to be in good condition, and managed appropriately (e.g. Lawson et al. 2014). These management actions should be based on the best available evidence: both from contemporary studies of the (currently) resident species, but also from palaeoecological studies of the site or region, where available. Such studies will often contain valuable information on the persistence of species in the longer-term.

Acknowledgements

This project was funded by Natural England (Ref #: 24837). Conversations with Jacqui P. Huntley improved our coverage of studies in the North Pennines substantially. We are also grateful to the academic colleagues who gave up their time to participate in the Natural England Evidence Review which preceded this work: Jane Bunting, Frank Chambers, Dan Charman, Mary Edwards, Peter Langdon, Danielle Schreve, Helen Shaw and Nikki Whitehouse.

References

- Bennett KD (1988) Holocene pollen stratigraphy of central East Anglia, England, and comparison of pollen zones across the British Isles. *New Phytol* 109:237-253
- Bennett KD, Simonson WD, Peglar SM (1990) Fire and man in post-glacial woodlands of Eastern England. *J Archaeol Sci* 17:635-642
- Birks HJB (1989) Holocene isochrone maps and patterns of tree-spreading in the British Isles. *J Biogeogr* 16:503-540

Davis, BS *et al.* (81 authors, 2013) The European Modern Pollen Database (EMPD) project. *Veget Hist Archaeobot* 22:521-530

Fyfe RM, de Beaulieu JL, Binney H, Bradshaw RHW, Brewer S, Le Flao A, Finsinger W, Gaillard MJ, Giesecke T, Gil-Romera G, Grimm EC, Huntley B, Kunes P, Kuhl N, Leydet M, Lotter AF, Tarasov PE, Tonkov S (2009) The European Pollen Database: past efforts and current activities. *Veget Hist Archaeobot* 18:417-424

Godwin H, Clifford MH (1938) Studies of the post-glacial history of British vegetation I Origin and stratigraphy of Fenland deposits near Woodwalton, Hunts II Origin and stratigraphy of deposits in Southern Fenland. *Philos T Roy Soc B* 229:323-406

Godwin H (1956) The history of the British flora. Cambridge University Press, Cambridge

Huntley B, Birks HJB (1983) An atlas of past and present pollen maps of Europe: 0-13,000 years ago. Cambridge University Press, Cambridge

Huntley B (1991) Historical lessons for the future. In: Spellerberg IF, Goldsmith FB, Morris MG (eds) The scientific management of temperate communities for conservation. Blackwell Scientific Publications, Oxford.

Huntley B (1996) Quaternary palaeoecology and ecology. *Quaternary Sci Rev* 15:591-606

Huntley J (2011) Alston Moor Miner-Farmer landscape: palaeoenvironmental assessment. English Heritage, Portsmouth

Jessen K (1949) Studies in late Quaternary deposits and flora-history of Ireland. *P Roy Irish Acad B* 52:85-290.

Lawson CR, Bennie JJ, Thomas CD, Hodgson JA, Wilson RJ (2014) Active management of protected areas enhances metapopulation expansion under climate change. *Cons Lett* 7:111-118.

Lawton J *et al.* (14 authors, 2010) Making Space for Nature: A review of England's Wildlife Sites and Ecological Network. Defra, London

Peglar SM, Fritz SC, Birks HJB (1989) Vegetation and Land-Use History at Diss, Norfolk, U.K. *J Ecol* 77:203-222


Pennington W (1969) History of vegetation in Britain. English Universities Press, London

Seddon AWR *et al.* (69 authors, 2014) Looking forward through the past: identification of 50 priority research questions in palaeoecology. *J Ecol* 102:256-267


Turner J, Hewetson VP, Hibbert FA, Lowry KH, Chambers C (1973) The history of the vegetation and flora of Widdybank Fell and the Cow Green reservoir basin, Upper Teesdale. *Philos T Roy Soc B* 265:327-408

Fig. 1 Location of palaeoecological records derived from literature obtained by systematic searching of Web of Knowledge and Google Scholar, or from the personal knowledge of the authors. The location of all dated and undated sites is shown in (a). The number of dated samples at each site is shown in (b), with the size and colour of symbol proportional to the number of dated samples. Panel (c) shows the number of dated cores available for each millennium over the last 15,000 years.


(a)



(b)



(c)



Supplementary Table 1 The English sediment core database

A database, provided as a Microsoft Excel spreadsheet accompanying this manuscript, provides details of all studies from which data were extracted. The geographic location, duration of the record, the total number of samples and the number of dates is listed for each record.

Supplementary Table 2 Search terms used to compile the English sediment core database

- | | |
|----|---------------------------------|
| 1 | 'Holocene' + 'pollen' |
| 2 | 'Quaternary' + 'pollen' |
| 3 | 'Pleistocene' + 'pollen' |
| 4 | 'Holocene' + 'palynolog*' |
| 5 | 'Quaternary' + 'palynolog*' |
| 6 | 'Pleistocene' + 'palynolog*' |
| 7 | 'Holocene' + 'fossil' |
| 8 | 'Quaternary' + 'fossil' |
| 9 | 'Pleistocene' + 'fossil' |
| 10 | 'paleoecologic*' |
| 11 | 'pal*o* + refug*', |
| 12 | 'pal*oecolog* + climat* change' |
| 13 | 'pollen' + 'Engl*' |
| 14 | 'fossil' + 'Engl*' |
| 15 | 'palynolog* + 'Engl*' |
| 16 | 'glaci* ref*' |
-