

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass and global biogeochemical cycles

C. Le Quéré¹, E. T. Buitenhuis¹, R. Moriarty¹, S. Alvain², O. Aumont³, L. Bopp⁴, S. Chollet⁵, C. Enright¹, D. J. Franklin⁶, R. J. Geider⁷, S. P. Harrison⁸, A. Hirst^{9,10}, S. Larsen¹¹, L. Legendre^{12,13}, T. Platt¹⁴, I. C. Prentice^{8,15}, R. B. Rivkin¹⁶, S. Sathyendranath¹⁴, N. Stephens¹⁴, M. Vogt¹⁷, S. Sailley¹⁴, and S. M. Vallina¹⁸

¹Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia, Norwich Research Park, NR4 7TJ, Norwich, UK

²Laboratoire d'Océanologie et de Géosciences – UMR LOG 8187, Université Lille Nord de France, BP 8062930 Wimereux, France

³Laboratoire d'Océanographie et de Climatologie: Expérimentation et Approches Numériques, IRD/IPSL, Plouzané, France

⁴Lab. des Sciences du Climat et de l'Environnement, Orme des Merisiers, Bat. 709, 91191 Gif-sur-Yvette, France

⁵School of Environmental Sciences, University of East Anglia, Norwich Research Park, NR4 7TJ, Norwich, UK

⁶Faculty of Science and Technology, Bournemouth University, Talbot Campus, Poole, BH12 5BB, UK

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

⁷School of Biological Sciences, University of Essex, Colchester CO4 3SQ, UK

⁸Department of Biological Sciences, Macquarie University, North Ryde, NSW 2109, Australia
and School of Archaeology, Geography and Environmental Sciences (SAGES), University of
Reading, Whiteknights, Reading, RG6 6AB, UK

⁹School of Biological and Chemical Sciences, Queen Mary University of London, London, E1
4NS, UK

¹⁰Centre for Ocean Life, National Institute for Aquatic Resources, Technical University of
Denmark, Kavalergården 6, 2920 Charlottenlund, Denmark

¹¹Norwegian Institute of Marine Research, Nye Flødevigveien 20, His, 4817, Norway

¹²Sorbonne Universités, UPMC Univ Paris 06, UMR 7093, LOV, Observatoire océanologique,
06230, Villefranche-sur-Mer, France

¹³CNRS, UMR 7093, Laboratoire d'Océanographie de Villefranche, 06230
Villefranche-sur-Mer, France

¹⁴Plymouth Marine Laboratory, Prospect Place, Plymouth, PL1 3DH, UK

¹⁵AXA Chair of Biosphere and Climate Impacts, Grand Challenges in Ecosystems and the
Environment and Grantham Institute – Climate Change and the Environment, Department of
Life Sciences, Silwood Park Campus, Buckhurst Road, Ascot SL5 7PY, UK

¹⁶Department of Ocean Sciences, Memorial University of Newfoundland, St. John's, NL A1C
3T3 Canada

¹⁷Institute for Biogeochemistry and Pollutant Dynamics, Universitätsstrasse 16, 8092 Zürich,
Switzerland

¹⁸Institute of marine Sciences (CSIC), Department Marine Biology and Oceanography, 08003
Barcelona, Spain

Received: 9 June 2015 – Accepted: 11 June 2015 – Published: 30 July 2015

Correspondence to: C. Le Quéré (c.lequere@uea.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Abstract

Global ocean biogeochemistry models currently employed in climate change projections use highly simplified representations of pelagic food webs. These food webs do not necessarily include critical pathways by which ecosystems interact with ocean biogeochemistry and climate. Here we present a global biogeochemical model which incorporates ecosystem dynamics based on the representation of ten plankton functional types (PFTs); six types of phytoplankton, three types of zooplankton, and heterotrophic bacteria. We improved the representation of zooplankton dynamics in our model through (a) the explicit inclusion of large, slow-growing zooplankton, and (b) the introduction of trophic cascades among the three zooplankton types. We use the model to quantitatively assess the relative roles of iron vs. grazing in determining phytoplankton biomass in the Southern Ocean High Nutrient Low Chlorophyll (HNLC) region during summer. When model simulations do not represent crustacean macrozooplankton grazing, they systematically overestimate Southern Ocean chlorophyll biomass during the summer, even when there was no iron deposition from dust. When model simulations included the developments of the zooplankton component, the simulation of phytoplankton biomass improved and the high chlorophyll summer bias in the Southern Ocean HNLC region largely disappeared. Our model results suggest that the observed low phytoplankton biomass in the Southern Ocean during summer is primarily explained by the dynamics of the Southern Ocean zooplankton community rather than iron limitation. This result has implications for the representation of global biogeochemical cycles in models as zooplankton faecal pellets sink rapidly and partly control the carbon export to the intermediate and deep ocean.

1 Introduction

Phytoplankton, zooplankton and heterotrophic bacteria (including both *Bacteria* and *Archaea*, herein called “bacteria”) in the oceans control important ecosystem processes

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



and services (Ducklow, 2008), including primary, secondary and export production. Primary production, i.e. the production of organic matter by photoautotrophs using inorganic nutrients, can be either particulate and serve as food for small heterotrophs from protists to fish larvae, or dissolved and used by bacteria. Secondary production, the fraction produced by zooplankton grazing on phytoplankton, smaller zooplankton, or organic detritus, serves as food for larger organisms in the ocean, including fish and mammals. Export production, the fraction of primary production that sinks below the mixed layer, exerts an influence on marine biogeochemistry and climate as most sinking organic matter is remineralized to inorganic matter at depths where it becomes isolated from the atmosphere for decades to centuries. Export production responds primarily to the activity of larger plankton, particularly the production and sinking of faecal pellets by meso- and macrozooplankton and larger organisms as well as the aggregation of diatoms, for example, during intense blooms. Export production lowers the surface concentration of inorganic carbon and maintains atmospheric CO₂ about 200 ppm lower than it would be in the absence of biological activity (Maier-Reimer et al., 1996). In contrast, bacteria and small zooplankton remineralize and recycle organic matter in the upper ocean, thus reducing the quantity of organic matter that is exported. These ecosystem processes are controlled by the state of the environment (e.g. temperature, light, available nutrients, vertical mixing), and are modulated by ecosystem structure 15 the plankton community.

Dynamic Green Ocean Models have been developed and used in global biogeochemical studies to understand and quantify the interactions between marine ecosystems and the environment. In these models, phytoplankton and zooplankton are grouped by taxa into plankton functional types (PFTs) according to their specific role in 20 marine biogeochemical cycles (Hood et al., 2006; Le Quéré et al., 2005). Although generally only a small number of PFTs are treated explicitly, their inclusion has been shown to improve the realism of model simulations. For example, the explicit inclusion of diatoms in marine ecosystem models is required to reproduce the observed response to natural or purposeful iron fertilisation in the ocean (Aumont and Bopp, 2006), and

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



observed changes in export production during glacial cycles (Bopp et al., 2003). The representation of diazotrophs (i.e. N₂-fixing organisms) is necessary to simulate the feedbacks between iron and the nitrogen inventories of the ocean (Moore et al., 2006; Moore and Doney, 2007), of coccolithophores to simulate large blooms of phytoplankton (i.e. chlorophyll) biomass (Gregg and Casey, 2007) and phytoplankton succession (Gregg et al., 2003), and of *Phaeocystis* to reproduce the ecosystem structure in the Southern Ocean (Wang and Moore, 2011).

Fewer studies have examined the role of different zooplankton PFTs in global ocean biogeochemistry, even though there are data sets on zooplankton physiology (e.g. Hirst and Bunker, 2003; Straile, 1997). Zooplankton can influence the fate of exported materials through various processes, including grazing, repackaging of organic matter in faecal pellets, and vertical migrations in the mesopelagic realm (e.g. Stemmann et al., 2000). Furthermore, there are important interactions among grazing, nutrient cycles, and environmental conditions as has been shown from studies based on regional models and observations in the equatorial Pacific (Landry et al., 1997; Price et al., 1994), North Pacific (Frost, 1991), the Atlantic (Daewel et al., 2014; Steinberg et al., 2012) and the Southern Ocean (Banse, 1995; Bishop and Wood, 2009). The importance of grazing has also been highlighted during iron enrichment experiments (Henjes et al., 2007; Latasa et al., 2014), in part explaining why some experiments led to increased carbon export and others did not (Martin et al., 2013). Thus, a more explicit representation of different zooplankton PFTs in global models could provide important clues for the functioning of marine biogeochemistry. More mechanistic parameterisations of zooplankton dynamics constrained by observations have been shown to improve simulations of phytoplankton biomass (Buitenhuis et al., 2006, 2010), the choice of grazing formulation to influence phytoplankton diversity (Prowe et al., 2012; Vallina et al., 2014b) and the resulting food web dynamics (Sailley et al., 2013; Vallina et al., 2014a), and to have implications for energy flow to higher trophic levels (Stock et al., 2014).

Here, we present a new global ocean biogeochemistry model with ten PFTs. The parameterisation of vital rates associated with these PFTs is based on an extensive

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

synthesis of published information on growth rates and other relevant parameters. We use the model to examine a long-standing paradox in biological oceanography: the low phytoplankton biomass in the Southern Ocean despite the high concentrations of macronutrients. This phenomenon has been attributed to lack of iron (Fe) because of the distance to continental dust sources (Geider and La Roche, 1994; Martin, 1990). Increases in phytoplankton biomass have been produced in more than a dozen open ocean iron fertilisation experiments (Boyd and al., 2007; Smetacek et al., 2012). The influx of Fe has been proposed as a driver for the drawdown of atmospheric CO₂ during glaciations (Kohfeld et al., 2005; Watson et al., 2000), and intentional Fe-fertilisation has been considered as a means to both geo-engineer climate (Rickels et al., 2012) and to sell carbon credits (Tollefson, 2012). However, ocean biogeochemistry models that explicitly include the effect of Fe-limitation on phytoplankton growth fail to reproduce the low Chl biomass observed during summer in the Southern Ocean (Aumont and Bopp, 2006; Dutkiewicz et al., 2005; Le Quéré et al., 2005; Moore et al., 2004). This raises the question of the relative control exerted by Fe-limitation on biomass vs. that exerted by the grazing pressure of zooplankton (Banse, 1996; Price et al., 1994) and more generally on the suitability of the current generation of models to explore ecosystem – climate interactions. Our study addresses this question directly.

2 Methods

2.1 Model description and development

The PlankTOM10 Dynamic Green Ocean Model is a global ocean biogeochemistry model that includes plankton ecosystem processes based on the representation of ten PFTs and their interactions with the environment. PlankTOM10 incorporates six autotrophic and four heterotrophic PFTs: picophytoplankton (pico-eukaryotes and non N₂-fixing cyanobacteria such as *Synechococcus* and *Prochlorococcus*), N₂-fixers (*Trichodesmium* and N₂-fixing unicellular cyanobacteria), coccolithophores, mixed-

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



phytoplankton (e.g. autotrophic dinoflagellates and chrysophytes), diatoms, colonial *Phaeocystis*, bacteria (here used to subsume both heterotrophic *Bacteria* and *Archaea*), protozooplankton (e.g. heterotrophic flagellates and ciliates), mesozooplankton (predominantly copepods), and crustacean macrozooplankton (euphausiids, amphipods, and others, called “macrozooplankton” for simplicity; Fig. 1). Gelatinous macrozooplankton are not included in the model. Diversity within PFTs is not considered, and the physiological parameters for each PFT are the same everywhere in the ocean, although some vary as a function of environmental conditions (i.e. nutrients, food, temperature).

The current version of the PlankTOM10 model was developed from the model of Buitenhuis et al. (2013a), using the strategy for regrouping PFTs described by Le Quéré et al. (2005). It does not include new parameterisations compared with previous versions of the PlankTOM model, but it includes an additional trophic level in the zooplankton PFTs (i.e. macrozooplankton). Parameterisations are based on more data related to the vital rates of individual PFTs, where new information was available. Previous studies have shown that model results are highly sensitive to PFT growth rates (Buitenhuis et al., 2006, 2010), and considerable effort was made to constrain these rates using observations from LaRoche and Breitbarth (2005), Bissinger et al. (2008), Buitenhuis et al. (2008, 2010), Sarthou et al. (2005), Schoemann et al. (2005), Rivkin and Legendre (2001), Hirst and Bunker (2003), and Hirst et al. (2003).

The complete set of model equations and parameter values are provided in the Supplement. Here, we describe the elements that are most important for the analysis of the Southern Ocean and the strategy used to determine parameter values for PFT growth and loss processes.

PlankTOM10 simulates the growth of ten PFTs in response to environmental conditions. The model includes three detrital pools: large and small particulate organic matter, and semi-labile dissolved organic matter. The sinking speed of large particles is based on the mineral (ballast) content of particles following Buitenhuis et al. (2001), while the sinking speed of small particles is constant at 3 md^{-1} . The model includes

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

full cycles of carbon (C), oxygen (O_2), and phosphorus (P), which are assimilated and released by biological processes at a constant ratio of 122 : 172 : 1 (Anderson and Sarmiento, 1994). It also includes a full cycle of silica (Si) as in Maier-Reimer (1993), and simplified cycles for Fe and nitrogen (N). CO_2 and O_2 are exchanged with the atmosphere using the gas exchange formulation of Wanninkhof (1992). The Fe cycle is represented as in Aumont and Bopp (2006). Iron is deposited with dust particles using the monthly fields of Jickells et al. (2005), the Fe content of dust is assumed to be 3.5 % everywhere. We use an Fe solubility of 1 % (Jickells et al., 2005). Iron is also delivered to the ocean via river fluxes following the outflow scheme of da Cunha et al. (2007) with 5 95 % sedimentation in estuaries. Dissolved inorganic nitrogen (DIN) is the sum of nitrate and ammonium. The N : P ratio of organic processes is set to the Redfield ratio of 10 16 : 1. N_2 -fixers can use N_2 and thus have access to unlimited N from the atmosphere.

The growth rate parameters for the ten PFTs in PlankTOM10 are based on a compilation of growth rates as a function of temperature (Sect. 2.2). Phytoplankton PFT 15 growth rates are also limited by light and inorganic nutrients (P, N, Si, and Fe). Light limitation is constrained by the slope of the photosynthesis-irradiance curve (α) and the maximum Chl/C ratio (θ_{max}). We could not distinguish PFT-specific values for α (Geider et al., 1997) and used a mean value of $1.0 \text{ mol C m}^2 (\text{g Chl mol photons})^{-1}$ for all PFTs. Observed θ_{max} for diatoms are systematically higher than those of other PFTs 20 (Geider et al., 1997). There are too few direct observations to parameterize θ_{max} for other PFTs, so we fitted the observations (Geider et al., 1997) for θ_{max} to the maximum growth rate (μ^{max}) presented in that paper. The fit showed θ_{max} increasing with growth rate ($n = 19$, $p = 0.02$). We thus used a θ_{max} higher than average for *Phaeocystis* and diatoms, and a lower than average θ_{max} for N_2 -fixers.

We used a two-step approach to define the nutrient limitation parameters, which 25 are not well constrained by observations. Firstly, we assigned PFT-specific limitation parameters to each phytoplankton PFT based on literature-derived values, using the value for a similar-sized PFT when PFT-specific information was not available. We then examined the covariations of surface Chl concentrations with each limiting nutrient and

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

zooplankton PFT and adjusted the magnitude of the limiting parameters of both the phytoplankton and zooplankton PFTs but keeping the ratios of these parameter values between phytoplankton PFTs and between zooplankton PFTs approximately the same as the original ratios.

- Initial values for the half saturation concentrations of P (k_P) and N (k_N) for phytoplankton growth rates were based on observations. For N₂-fixers, coccolithophores and diatoms, the half-saturation values for growth were computed using the half-saturation values of uptake reported in Riegman et al. (1998), LaRoche et al. (2005), and Sarthou et al. (2005) multiplied by the minimum/maximum N:C ratio (0.33) to account for the acclimation of nutrient saturated vs. nutrient limited growth (Morel, 1987). For picophytoplankton, reported values for the half-saturation extend over three orders of magnitude. We assigned low half-saturation values as these organisms grow even under very low nutrient conditions (Timmermans et al., 2005). For mixed phytoplankton, we assigned a value intermediate between picophytoplankton and diatoms. For *Phaeocystis*, we used half saturation values that characterise colonies (Schoemann et al., 2005). The selected set of parameter values were tuned to reproduce the observed covariation (or lack of covariation) between Chl and N, P, and Fe distributions (Fig. 3, Table 2).

- Iron uptake was computed using a cell quota model (Buitenhuis and Geider, 2010; Geider et al., 1997), where the Fe uptake by phytoplankton PFTs is explicitly regulated by the light conditions. The three parameters needed are the minimum, the maximum and the optimal Fe quotas. The minimum and maximum quotas were set at the same value of 2.5 and 20 µmol Fe (mol C⁻¹) for all PFTs. The optimal quota was set to the minimum quota plus $2 \cdot \mu_{20}^{\text{max}}$ based on (Sunda and Huntsman, 1995) for all PFTs. In addition, phytoplankton PFT also respond to the concentration of Fe in water which is parameterised with a half saturation. The half saturation of Fe uptake (k_{Fe}) is lower for picophytoplankton (Timmermans et al., 2005) than other phytoplankton, and higher for N₂-fixers (LaRoche and Breitbarth, 2005) and diatoms (Sarthou et al., 2005). Intermediate values for k_{Fe} have been reported for the other phytoplankton PFTs (Le Vu, 2005;

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Schoemann et al., 2005). The selected set of parameter values after adjustments in k_{Fe} produces no systematic covariation between Chl and Fe, as observed (Fig. 3, Table 2).

The half-saturation parameters of zooplankton grazing rate were initially set to a constant value of $20 \mu\text{mol CL}^{-1}$ for zooplankton PFTs and $60 \mu\text{mol CL}^{-1}$ for bacteria, based on the relationship between metabolic rates and body volume of Hansen et al. (1997), and subsequently adjusted to reproduce the Chl-zooplankton covariation (Fig. 3, Table 2).

Zooplankton food preferences were assigned based on predator-prey size ratio (Table 3), as there were insufficient data to determine these parameters directly. This approach assumes that protozooplankton generally have a high preference for bacteria and a low preference for diatoms, that mesozooplankton have a higher preference for protozooplankton and a low preference for N₂-fixers and bacteria, and macrozooplankton have a lower preference for N₂-fixers, picophytoplankton and bacteria than other groups. We assume that all zooplankton graze on both organic particles (Table 3) but prefer to graze on other PFTs. The gross growth efficiency (the part of grazing that is incorporated into biomass) was defined based on the mean across available observations: 0.21 for bacteria (data from Rivkin and Legendre, 2001), and 0.29, 0.25, and 0.30 for protozooplankton, mesozooplankton and macrozooplankton, respectively (data from Straile, 1997). Respiration and mortality parameters were based on observations from Buitenhuis et al. (2010) for protozooplankton, Buitenhuis et al. (2006) for mesozooplankton, and Moriarty (2013) for macrozooplankton. The temperature-dependence of respiration and mortality was fitted to all data as for the growth rate (Sect. 2.2), except for the mortality of macrozooplankton and mesozooplankton. There are nine observations on macrozooplankton mortality and we tuned this term based on the resulting biomass. The fitted relationship for the mortality of mesozooplankton was reduced by a factor of ~ 2 to account for the explicit mortality from macrozooplankton represented in the model.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2.2 Growth rates as a function of temperature

The most important trait that distinguishes the various PFTs is the rate at which they grow under different conditions (Buitenhuis et al., 2006, 2010). We compiled maximum growth rates as a function of temperature (Table 1). We fit an exponential growth relationship to the observations by optimising the relation $\mu^T = \mu_0 \cdot Q_{10}^{T/10}$ where T and μ^T are the observed temperature and associated growth rate, μ_0 is the growth at 0 °C, and Q_{10} is the derived temperature-dependence of growth (Table 1).

5 The parameter values for μ_0 and Q_{10} were estimated by minimising the error, quantified as the least squares cost function $\sum (\mu^T - \mu_{\text{obs}}^T)^2 / \mu_{\text{obs}}^T$. Normalising to observations helps ensure a good fit
10 of μ^T in cold waters where growth rates are low. We used exponential growth, rather than a temperature-optimal growth, to avoid biases caused by the lack of observations for some PFTs at low or high temperatures. The p value of a linear regression between observations and the exponential fit (Table 1) provides a measure of how well the relationship is constrained by the observations. The fit assigns equal weight for all the
15 data, rather than following the 99 % quantile (e.g. Eppley, 1972; Bissinger et al., 2008) to provide a better representation of the mean community for each PFT.

Growth rate parameters estimated with this method are well constrained (p values < 0.05) for seven of the ten PFTs, including all of the heterotrophic PFTs (Table 1). There are insufficient data to provide significant constraints on the growth rates of
20 N_2 -fixers ($p = 0.76$), and some uncertainty in the growth data for coccolithophores ($p = 0.06$) and *Phaeocystis* ($p = 0.23$; Table 1). However, the growth of N_2 -fixers is less than that of other phytoplankton PFTs (Fig. 2), and the fitted relationship produces μ^T less than that of other PFTs despite these uncertainties. An exponential function may not be appropriate for growth rates of coccolithophores and *Phaeocystis* (Schoemann et al., 2005). The growth rate of coccolithophores was overestimated at low temperatures due to high growth rates at 20 °C and the absence of observations for temperatures below 5 °C. We reduced the fitted growth rate of coccolithophores linearly to 0
25

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

below 10 °C to match the observed reduced coccolithophore biomass in cold regions (O'Brien et al., 2013).

2.3 Covariation between Chl and nutrients or zooplankton

We used relationships between observed concentrations of Chl and both inorganic nutrients (e.g. NO_3 , PO_4 and Fe), and zooplankton biomasses (protozooplankton, mesozooplankton and macrozooplankton; Fig. 3) to provide additional constraints on model parameters. Specifically, we used observations for in situ NO_3 and PO_4 concentrations from the World Ocean Atlas 2009; in situ Fe concentration data from Tagliabue et al. (2012); protozooplankton biomass data from Buitenhuis et al. (2010); mesozooplankton biomass data from Buitenhuis et al. (2006); macrozooplankton biomass data from Atkinson et al. (2004) and Moriarty et al. (2013). All the data were binned into $1 \times 1^\circ$ grid boxes. Most observations are for the surface ocean. Mesozooplankton and macrozooplankton data are from depth-integrated tows of typically 200 m depth and may underestimate surface concentrations (by a factor 1.5–2 based on our model simulations). All data are monthly except for mesozooplankton, which are seasonal. Chl concentration is from SeaWiFS satellite averaged over 1998–2009 and interpolated to the same grid. The model output was averaged over the same time period, and sampled for the same month and on the same grid box as the observations. The data intervals were chosen to include approximately the same number of grid boxes, except for macrozooplankton where the lowest interval was set to 0–0.05 $\mu\text{mol CL}^{-1}$ because of the large number of grid boxes with very low macrozooplankton concentration. Ten concentration intervals were used for the nutrients (Fig. 3).

Chlorophyll concentrations covary with NO_3 concentrations at $< 3 \mu\text{mol L}^{-1}$, and with PO_4 in the range $0.3\text{--}0.5 \mu\text{mol L}^{-1}$ (Fig. 3; Spearman ranked correlations for data in the 25–75 % interquartile range gives $r = 0.72$ for NO_3 and $r = 0.73$ for PO_4). These relationships are consistent with our understanding of the growth limitation of phytoplankton in the subtropics, where NO_3 and PO_4 concentrations are low. There is no observed covariation between Chl and Fe concentration ($r = -0.16$). The strongest covaria-

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

tions are found between Chl and protozooplankton at concentrations $< 0.6 \mu\text{mol CL}^{-1}$ ($r = 0.83$) and mesozooplankton at concentrations $< 0.3 \mu\text{mol CL}^{-1}$ ($r = 0.77$). There is no covariation between Chl concentration and macrozooplankton biomass ($r = -0.19$; Fig. 3). We use these relationships to tune the growth limitations parameters in the model, so that the functional relationships between Chl and nutrients or zooplankton are close to the observed relationships overall.

2.4 Simulations

PlankTOM10 is coupled to the Ocean General Circulation Model (OGCM) NEMO version 3.1 (NEMOv3.1). We used the global configuration (Madec and Imbard, 1996), which has a resolution of 2° of longitude and a mean resolution of 1.5° of latitude, with enhanced resolution up to 0.3° in the tropics and at high latitudes. The model resolves 30 vertical levels, with 10 m depth resolution in the upper 100 m. NEMOv3.1 calculates vertical diffusion explicitly and represents eddy mixing using the parameterisation of Gent and McWilliams (1990). NEMOv3.1 is coupled to a dynamic-thermodynamic sea-ice model (Timmermann et al., 2005).

PlankTOM10 is initialised from observations for dissolved inorganic carbon (DIC) and alkalinity from Key et al. (2004), O_2 and nutrients from Garcia et al. (2006a) and Garcia et al. (2006b), and temperature and salinity from the World Ocean Atlas 2005 (Antonov et al., 2006; Locarnini et al., 2006). Fe is initialised at a constant concentration of $0.6 \text{ nmol Fe L}^{-1}$ north of 30°S and $0.2 \text{ nmol Fe L}^{-1}$ in the Southern Ocean, consistent with observations (Parekh et al., 2005; Tagliabue et al., 2012). The PFTs equilibrated within 3 years and were not influenced by initialisation. The model is forced by daily winds and precipitation from the ECMWF interim reanalysis (Simmons et al., 2006) from 1989 to 2009. Results for standard simulations are averaged over 1998–2009; sensitivity tests are run for a shorter ten-year period and averaged over 1995–1999.

To understand the interaction pathways among ecosystems, biogeochemistry and climate, we developed a simplified version of the model that included only six PFTs

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



(PlankTOM6) (Fig. 1). PlankTOM6 is identical to PlankTOM10 except that the growth rates of N₂-fixers, mixed-phytoplankton, *Phaeocystis*, and macrozooplankton are zero, and the mortality of the mesozooplankton is increased to account for the lack of macrozooplankton predation until the point when primary production is at its maximum. Given the otherwise similar model structure, parameters, initialisation and simulation protocol, comparison of results from PlankTOM6 and PlankTOM10 allows us to isolate the specific roles of zooplankton dynamics in the model.

3 Results

3.1 Temperature and size – dependence of PFT growth rates

- The data show systematic patterns in growth rates that differ among PFTs. The growth rates of all PFTs increase with increasing temperature, but not to the same extent (Fig. 2). The growth rate of phytoplankton PFTs increases with PFT size, from 0.15 d⁻¹ for N₂-fixers to 1.87 d⁻¹ for *Phaeocystis*, and the growth rate of heterotrophic PFTs decreases with size, from 1.22 d⁻¹ for bacteria to 0.19 d⁻¹ for macrozooplankton (Table 1).
- The sign of the relationship between growth rate and size between phytoplankton PFTs is the opposite of the sign of this relationship within specific PFTs, including diatoms (Sarthou et al., 2005), picophytoplankton (Chen and Liu, 2010) and coccolithophores (Buitenhuis et al., 2008). From these relationships, we conclude that the observed phytoplankton growth rates may be more influenced by eco-evolutionary determinants (e.g. reproduction strategies) than by environmental physical constraints (e.g. diffusion rates across cell walls).

3.2 Ecosystem properties in the PlankTOM10 model

PlankTOM10 reproduces the main characteristics of observed surface Chl, with high concentrations in the high latitudes and low concentrations in the subtropics, higher

Chl concentration in the Northern compared to the Southern Hemisphere, and in the South Atlantic compared to the South Pacific Ocean (Fig. 4). The global biogeochemical fluxes simulated by PlankTOM10 are generally below or at the low end of the range of observed values (in Table 4, “model” and “data”, respectively), with global primary production of 42.4 PgCyr^{-1} , export production of 7.6 PgCyr^{-1} , export of CaCO_3 and SiO_2 of 0.4 PgCyr^{-1} and 2.9 PgSiyr^{-1} , respectively, and N_2 fixation of 165 TgNyr^{-1} .

PlankTOM10 produces distinctive geographical distributions of carbon biomasses among PFTs (Fig. 5). About a third of the phytoplankton biomass occurs as picophytoplankton, followed in descending abundance by diatoms and *Phaeocystis*, mixed-phytoplankton, coccolithophores and N_2 -fixers (Table 4). This distribution is broadly consistent with observations (Buitenhuis et al., 2013b) but the simulated phytoplankton biomass is generally on the low side of the observational range, which is consistent with the results from the global rates. The simulated biomass of coccolithophores is overestimated (i.e. 0.077 PgC compared with $0.001\text{--}0.032 \text{ PgC}$) although CaCO_3 export is underestimated, suggesting either that the model calcification or aggregation rates are too low or that zooplankton calcifiers contribute significantly to CaCO_3 export.

The model underestimates bacterial biomass by a factor of 10 compared with observations. This probably reflects the fact that the model only represents high activity bacteria and a substantial fraction of observed biomass is from low activity and ghost cells. The model underestimates protozooplankton by a factor of 1.5–5 (in absolute value) or 2–3 (as a fraction of total biomass value) compared to observations (Table 4). This discrepancy could be caused by the underestimation of bacterial biomass, as bacteria are an important source of food for protozooplankton. The simplified representation of the range of protozooplankton grazers in a single PFT (instead of having heterotrophic nanoflagellates and microzooplankton) could also play a role. Simulated mesozooplankton biomass is only slightly less than the observed range, while simulated macrozooplankton biomass is within the observed range, although the uncertainty here is large ($0.010\text{--}0.64 \text{ PgC}$). Overall the balance is slightly skewed towards relatively more biomass than observed in the larger zooplankton (53 % compared to

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
-
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

3–47 %) compared to the smaller zooplankton groups (13 % compared to 27–31 %; Table 4).

The geographic distribution of each simulated PFT is also distinctive (Figs. 6 and 7). Satellite data products indicate that small phytoplankton (picophytoplankton and N₂-fixers) are generally dominant in the tropics, haptophytes (coccolithophores and *Phaeocystis*) in mid to high latitudes, and diatoms in high latitudes (Alvain et al., 2005; Brewin et al., 2010). The simulated phytoplankton distribution generally matches the distribution inferred from satellite normalised radiance (Fig. 6), except in the temperate zones where observations suggest a balance between picophytoplankton and haptophytes and the model shows a dominance of haptophytes. PlankTOM10 also reproduces the locations of blooms of colonial *Phaeocystis* and coccolithophores (Fig. 7). The simulated geographic distributions of zooplankton PFTs are particularly distinctive, with protozooplankton abundant in the tropics and subtropics, mesozooplankton at high latitudes of both hemisphere, and macrozooplankton with high biomass in the North Pacific and South Atlantic and along the coasts (Fig. 5). There is generally high contrast between high and low biomass regions.

3.3 Comparison of PlankTOM6 and PlankTOM10

PlankTOM10 and PlankTOM6 generally produce similar results in surface Chl concentration, nutrient distribution, primary and export production (Fig. 8), except that PlankTOM6 fails to reproduce the observed low Chl concentration in summer in the Southern Ocean (Fig. 4; Sect. 3.4). The overall difference between the two models, quantified statistically using a Taylor distribution (Taylor, 2001), are less than 0.1 in either correlation or normalised standard deviation (Fig. 8). PlankTOM10 does slightly better for the distribution of Chl, primary and export production, but slightly worse for the distribution of silica and nitrate, with similar performance for phosphate (Fig. 8). These differences are small in part because of the short duration of the simulations presented here (20 years), which allow equilibration of the ocean surface only.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

3.4 Role of zooplankton dynamics for HNLC regions

The observed phytoplankton biomass, including the low Chl concentrations in HNLC regions, reflects the balance between phytoplankton growth and loss. Phytoplankton growth rates vary with temperature, light, and nutrient supply, whereas losses result mainly from grazing by zooplankton, respiration and cell death, viral lysis, sinking to depth, and dilution by vertical mixing. Any process that reduces the net rate of increase of phytoplankton biomass (i.e. differences between growth and loss) may lead to low residual Chl concentration, even in nutrient-replete environments such as in HNLC regions. For example, Platt et al. (2003a) showed that deep mixing by wind dilutes the Chl in the surface layer and reduces the average irradiance experienced by the phytoplankton and results in low growth rate and demand for nitrate; the conditions generally observed in HNLC regions. Here we further examine the consequences of high zooplankton-mediated grazing losses.

We use the North/South ratio in surface Chl concentration as a metric to quantify model performance, focusing on the Pacific Ocean where the contrast between the Northern Hemisphere and the Southern Ocean is most pronounced. This metric is simple and easy to quantify with data (geographic locations: boxes in Fig. 4). Satellite observations indicate a North/South Chl ratio of 2.16 ± 0.35 (1998–2009 mean $\pm 2\text{SD}$ of annual values). To ensure that the ratio is not affected by potential biases in the SeaWiFS Southern Ocean data (Johnson et al., 2013), we also used in situ data from the World Ocean Atlas which indicates a similar North/South Chl ratio of 2.0. This ratio is 1.72 ± 0.051 in the PlankTOM10, and 1.21 ± 0.074 in the PlankTOM6 simulations (Fig. 9). Controlling factors on this ratio are examined here through a set of sensitivity tests.

3.4.1 Role of trophic level and top zooplankton

We tested the specific effect of macrozooplankton on Chl by running four additional model experiments (Fig. 9): in the Z1 simulation, we added macrozooplankton to Plank-

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



TOM6, in Z2 we parameterised the top grazer in PlankTOM6 using the same growth and loss rate parameters as macrozooplankton, in Z3 we removed macrozooplankton from PlankTOM10, and in Z4 we parameterised the top grazer in PlankTOM10 using the same growth and loss rate parameters as mesozooplankton. These sensitivity studies were identical to the PlankTOM10 (or PlankTOM6) simulation in all other respects. Experiments Z1 and Z2 both include macrozooplankton, but in different food-web positions. These experiments maintain a high North/South Chl ratio of 1.64 and 1.46, respectively (Fig. 9). Experiments Z3 and Z4 did not include macrozooplankton but had grazing structures as in the standard PlankTOM6 and PlankTOM10 models, the North/South Chl ratio was 1.26 and 1.11 respectively. These four experiments show that the presence in the model of slow-growing zooplankton, such as macrozooplankton, plays a pivotal role in determining the relative average concentrations of Chl in the Northern vs. Southern Hemisphere (difference between PlankTOM6 and both Z1 and Z2). More realistic patterns are achieved by including a third zooplankton food-web compartment (higher ratio in Z1 than in Z2) and three additional phytoplankton compartments (higher ratio in PlankTOM10 than in Z1).

3.4.2 Role of macrozooplankton growth rate

We examined the impact of macrozooplankton grazing in sensitivity tests in which the grazing rate of macrozooplankton was varied within the range of observations (Fig. 10). These simulations show that macrozooplankton grazing rate has a strong influence on the Chl North/South ratio. The PlankTOM10 simulation that uses the mean growth rate from observations (Sect. 2.2) produces results that are closest to the observed North/South Chl ratio. When the grazing rate is decreased (by up to 2σ), the macrozooplankton biomass decreases by over 50 % and the North/South Chl ratio decreases from 1.72 to 1.05. When the grazing rate is increased, the macrozooplankton biomass also decreases because of pressure on the food sources (Fig. 10) and the Chl North/South ratio also decreases. These simulations suggest that the observed Chl North/South distributions are a consequence of trophic balances among PFTs.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

3.4.3 Role of atmospheric iron deposition

We tested the relative role of atmospheric iron deposition compared with grazing for the North/South Chl distribution by applying five different dust deposition scenarios, all (except one) with realistic but different regional distributions, to the PlankTOM10 and PlankTOM6 models: D0 is an extreme case with no atmospheric dust deposition (where phytoplankton use iron sources from deep waters), D1 dust deposition including the effect of dust particle size on iron solubility (Mahowald et al., 2009), and D2-D4 iron deposition using the three distinct dust fields (Ginoux et al., 2001; Mahowald and Luo, 2003; Tegen et al., 2004) averaged by Jickells et al. (Jickells et al., 2005). The simulated North/South Chl ratios vary from 1.62 and 1.85 in these experiments (Fig. 9). These differences are smaller than the differences between the PlankTOM10-like (1.46–1.85) and the PlankTOM6-like simulations (1.08–1.26) for all experiments. In PlankTOM6, even the simulation with no iron deposition from dust (D0) produces Southern Ocean Chl concentrations that are too high during summer. This result is consistent with the observation that although Fe is lower in the Southern Ocean than elsewhere, concentrations average around $0.3 \text{ nmol Fe L}^{-1}$ (range of $0.15\text{--}0.6 \text{ nmol Fe L}^{-1}$) in the summer (January and February, $n = 79$) in the Subantarctic region (Tagliabue et al., 2012), which is near the half-saturation for growth of most phytoplankton as well as those used in the model (Le Quéré et al., 2005; Sarthou et al., 2005). Thus Fe concentrations may be limiting for phytoplankton growth, but nevertheless the observed very low Chl concentration during summer months seem to reflect loss processes due to grazing mortality rather than reduced growth rates from low Fe supply.

As a means of validating the model results, we also tested the response of PlankTOM10 to Fe-fertilisation to verify that the model reproduced the observed Chl blooms under Fe enrichment conditions (Boyd and al., 2007). This was done by saturating the surface layer of the ocean with Fe for one month (February). In this experiment, surface Chl south of 40° S increased by $2.1 \pm 2.2 \text{ mg Chl a m}^{-3}$ (mean $\pm 1\text{SD}$) with a maximum concentration of $14.2 \text{ mg Chl m}^{-3}$. This is similar to the responses observed at sea dur-

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

ing Fe-fertilisation experiments (Boyd and al., 2007). The response of the model to Fe enrichment provides further support of our hypothesis that grazing is responsible for the low Chl concentration in the Southern Ocean during summer.

3.4.4 Role of combined effects

- 5 Model simulations could be influenced by the model structure and parameters, the physical transport, meteorological data, or the choice of dust deposition fields. We assessed the combined effects of model choices by comparing our results with outputs from seven other models: a version of the PISCES model (Aumont and Bopp, 2006), the CCSM-BECs model (Doney et al., 2009), and the NEMURO model
10 (Kishi et al., 2007), IPSL-CM5A-LR (Dufresne et al., 2013), GRDL-ESM2M (Jones et al., 2011), HadGEM2-ES (Giorgetta et al., 2015), and CanESM2 (Arora et al., 2011). All of these other models focus on the representation of phytoplankton groups and parameterise grazing pathways in a simpler fashion than PlankTOM10. They produce a North/South Chl ratio in the range from 0.60 to 1.36, lower than the value (1.72)
15 obtained using PlankTOM10. Previous studies have suggested that the overestimation of Chl may result from a generalised model bias towards too shallow mixing depth in the Southern Ocean in summer, but Séférian et al. (2013) have shown that while better representation of sub-grid scale processes and mixed layer depth improves the simulation of Chl overall it does not lead to a more realistic North/South Chl ratio (Fig. 9).
20 Thus, the comparison between PlankTOM10 and other ocean biogeochemistry models supports our contention that it is important to simulate grazing pathways explicitly.

4 Discussion

The development of PlankTOM10 has capitalised on the existence of the very extensive range of observations to develop realistic parameterisations of key processes, particularly PFT growth rates. Although the simulated global biogeochemical fluxes are
25

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



generally below or at the low end of the range of observed values, the model reproduces both the relative importance of different PFTs and the geographic patterns in their abundance. Thus, while not perfect, the model is sufficient to explore the role of ecosystem dynamics in determining ocean biogeochemistry.

Our analyses suggest that Southern Ocean phytoplankton biomass during summer is primarily controlled by zooplankton grazing and the structure of the pelagic food web, rather than the low supply rate of iron. Trophic cascading appears to account for the differences between the results from PlankTOM10 and PlankTOM6. For example, protozooplankton graze on phytoplankton (and bacteria), which reduces their biomass. However, mesozooplankton graze on phytoplankton and protozooplankton, and macrozooplankton graze on phytoplankton and both protozooplankton and mesozooplankton. Thus the grazing pressure of larger zooplankton on smaller zooplankton can indirectly reduce the overall grazing pressure on phytoplankton. In PlankTOM10, macrozooplankton concentration is higher in winter in the Northern Hemisphere Pacific sector where the surface layer is more stratified and food is abundant, compared with the Southern Ocean Pacific sector where the surface layer is more mixed and food is scarce. Thus when the spring bloom starts in the North, the biomass and grazing pressure exerted by macrozooplankton are high enough to reduce the biomass of smaller zooplankton consequently reducing the grazing pressure on Chl and leading to an increase in Chl biomass. In the South, however, macrozooplankton biomass is too low to cause significant losses of smaller zooplankton. Hence, the high proto- and mesozooplankton biomasses prevent a phytoplankton bloom from developing in that region.

The higher concentration of macrozooplankton biomass in the North compared to the South is consistent with the observations (Moriarty et al., 2013). A similar contrast is found between the Atlantic and Pacific sectors of the Southern Ocean, where the high macrozooplankton biomass observed in the Atlantic (Atkinson et al., 2004) would reduce the abundance of smaller zooplankton resulting in higher Chl concentrations in the Atlantic sector, as simulated in PlankTOM10 (Fig. 4). Such trophic cascades have

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



been observed in diverse ecosystems on land and in the ocean (Casini et al., 2009). Furthermore, many observational-based studies have highlighted the important role of zooplankton grazing for controlling phytoplankton biomass (Atkinson et al., 2001; Banse, 1996; Dubischar and Bathmann, 1997; Granéli et al., 1993). Although some processes are missing from the model (e.g. vertical migration of zooplankton, which mostly contributes to downward export), the model suggests that the primary cascading effect of grazing is sufficient to account for a large part of the North/South Chl differences.

There are a number of limitations to the current version of PlankTOM10, including simplified overwintering strategies for zooplankton, the use of a simple Fe model, and the lack of representation of the more refractory organic matter which provides an additional food source for bacteria and may therefore contribute to the model overestimation of macrozooplankton biomass compared to protozooplankton and bacteria. In addition, the model does not include some ecosystem pathways, such as viral lysis (Evans et al., 2009). The realism of the simulations may also be affected by the relatively coarse resolution of the physical ocean model. However these biases affect both PlankTOM6 and PlankTOM10, and thus the experiments still provide information on the processes that differ between the two models. Our work suggests that improved representation of the zooplankton components in model could help further constrain the processes that regulate Chl distribution in models.

Our results on the important role of grazing do not contradict the results on the importance of Fe-fertilisation as highlighted in Fe enrichment experiments (Boyd and al., 2007), because additional Fe will trigger further growth provided that Fe is initially below an optimal concentration (Blain et al., 2007). However, our results suggest that low Fe concentrations by themselves are insufficient to account for the very low Chl levels observed in the Southern Ocean HNLC region in summer, and that differences in zooplankton trophic and community structure, and concomitant grazing dynamics have an important role in controlling phytoplankton blooms and maintaining very low Chl levels in that region. Although previous studies emphasised the role of phytoplankton com-

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

5 Conclusions

The development of global marine ecosystem models is hampered in particular because of our poor understanding of several critical processes and interactions, and the lack of global-scale observation of rates and biomass for parameterisation and validation (Le Quéré and Pesant, 2008). For example, the wide range in observed growth rates for the same temperature is an indication of the challenges met by modellers, particularly in representing the within-PFT diversity, which is unaccounted for in our model. Much more work is needed to understand the specific pathways by which matter circulates within ecosystems, taking into account the regional distributions of zooplankton groups and interactions with the environment including seasonal mixed layer dynamics.

The role of macrozooplankton highlighted here has implications for carbon export to depth because faecal pellets of some macrozooplankton have very fast sinking rates (Fortier et al., 1994; Turner, 2002). Hence, a more explicit representation of the pelagic food web in global models is needed to capture the full range of interactions between marine ecosystems, marine biogeochemistry and climate. The synthesis and analysis of observations and model results by the MAREDAT and MAREMIP projects provide valuable insights into the processes that control marine ecosystems, including the con-

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

tributions that different PFTs make to ocean biomass (Buitenhuis et al., 2013a; Hashioka et al., 2012; Salliey et al., 2013).

Our simulations examining the effects of grazing on phytoplankton biomass raise questions about the biological and biogeochemical bases for the current projections of the feedbacks between climate (and other environmental changes) and marine ecosystems. It also raises potential complications for the proposed use of purposeful Fe-fertilisation to enhance the deep ocean storage of CO₂ (Caias et al., 2013). Assessments of the impact of such geoengineering will be unreliable, at least until the full ecosystem response including the grazing pathways (Landry et al., 1997) and the relationship to deep water carbon export (Smetacek et al., 2012) can be reproduced with models, which could be used to make quantitative predictions.

The Supplement related to this article is available online at
doi:10.5194/bgd-12-11935-2015-supplement.

Acknowledgements. The structure of PlankTOM10 was developed through a series of seven international workshops funded in part by the Max-Planck Institute for Biogeochemistry, and hosted by the Villefranche Oceanography Laboratory, France. We thank C. Klaas and D. Wolf-Gladrow for their input on model development and interpretation, S. Pesant for support with data compilations, G. Madec and the NEMO team for assistance with the physical model, A. Tagliabue for providing the iron database, and N. Mahowald for providing dust deposition fields. C. Le Quéré and E. T. Buitenhuis were funded by the UK-NERC projects NE/C516079/1 and NE/K001302/1, and European Commission project 282672. R. Moriarty was funded by the EU FAASIS project MEST/CT/2004/514159. M. Vogt was funded by the Marie Curie Research and Training Network GREENCYCLES project MC-RTN-512464 and EUR-OCEANS project 282672. Model simulations were run on the High Performance Computing Cluster of the University of East Anglia.

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

References

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Bishop, J. K. B. and Wood, T. J.: Year-round observations of carbon biomass and flux variability in the Southern Ocean, *Global Biogeochem. Cy.*, 23, GB2019, doi:10.1029/2005GB002591, 2009.
- Bissinger, J. E., Montagnes, D. J. S., Sharples, J., and Atkinson, D.: Predicting marine phytoplankton maximum growth rates from temperature: improving on the Eppley curve using quantile regression, *Limnol. Oceanogr.*, 53, 487–493, 2008.
- Blain, S., Quéguiner, B., Armand, L., Belviso, S., Bomblé, B., Bopp, L., Bowie, A., Brunet, C., Brussaard, C., Carlotti, F., Christaki, U., Corbière, A., Durand, I., Ebersbach, F., Fuda, J.-L., Garcia, N., Gerringa, L., Griffiths, B., Guiguer, C., Guillerm, C., Jacquet, S., Jeandel, C., Laan, P., Lefèvre, D., Lo Monaco, C., Malits, A., Mosseri, J., Obernosterer, I., Park, Y.-H., Picheral, M., Pondaven, P., Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scouranec, L., Souhaut, M., Thuiller, D., Timmermans, K., Trull, T., Uitz, J., van Beek, P., Veldhuis, M., Vincent, D., Viollier, E., Vong, L., and Wagener, T.: Effect of natural iron fertilization on carbon sequestration in the Southern Ocean, *Nature*, 446, 1070–1075, 2007.
- Bopp, L., Kohfeld, K. E., Le Quere, C., and Aumont, O.: Dust impact on marine biota and atmospheric CO₂ in glacial periods, *Paleoceanography*, 18, 1046, doi:10.1029/2002PA000810, 2003.
- Boyd, P. W., Jickells, T., Law, C. S., Blain, S., Boyle, E. A., Buesseler, K. O., Coale, K. H., Cullen, J. J., de Baar, H. J. W., Follows, M., Harvey, M., Lancelot, C., Levasseur, M., Owens, N. P. J., Pollard, R., Rivkin, R. B., Sarmiento, J., Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S., and Watson, A. J.: Mesoscale iron enrichment experiments 1993–2005: synthesis and future directions, *Science*, 315, 612–617, 2007.
- Brewin, R. J. W., Lavender, S. J., and Hardman-Mountford, N. J.: Mapping size-specific phytoplankton primary production on a global scale, *J. Maps.*, 6, 448–462, 2010.
- Buitenhuis, E., Le Quéré, C., Aumont, O., Beaugrand, G., Bunker, A., Hirst, A., Ikeda, T., O'Brien, T., Piontkovski, S., and Straile, D.: Biogeochemical fluxes through mesozooplankton, *Global Biogeochem. Cy.*, 20, GB2003, doi:10.1029/2005GB002511, 2006.
- Buitenhuis, E. T. and Geider, R. J.: A model of phytoplankton acclimation to iron-light colimitation, *Limnol. Oceanogr.*, 55, 714–724, 2010.
- Buitenhuis, E. T., van der Wal, P., and de Baar, H.: Blooms of *Emiliania huxleyi* are sinks of atmospheric carbon dioxide; a field and mesocosm study derived simulation, *Global Biogeochem. Cy.*, 15, 577–588, 2001.

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Buitenhuis, E. T., Pangerc, T., Franklin, D. J., Le Quere, C., and Malin, G.: Growth rates of six coccolithophorid strains as a function of temperature, Limnol. Oceanogr., 53, 1181–1185, doi:10.4319/lo.2008.53.3.1181, 2008.

Buitenhuis, E. T., Rivkin, R. B., Sailley, S., and Le Quere, C.: Biogeochemical fluxes through microzooplankton, Global Biogeochem. Cy., 24, GB4015, doi:10.1029/2009GB003601, 2010.

Buitenhuis, E. T., Li, W. K. W., Lomas, M. W., Karl, D. M., Landry, M. R., and Jacquet, S.: Picoheterotroph (*Bacteria* and *Archaea*) biomass distribution in the global ocean, Earth Syst. Sci. Data, 4, 101–106, doi:10.5194/essd-4-101-2012, 2012.

Buitenhuis, E. T., Li, W. K. W., Vaulot, D., Lomas, M. W., Landry, M. R., Partensky, F., Karl, D. M., Ulloa, O., Campbell, L., Jacquet, S., Lantoine, F., Chavez, F., Macias, D., Gosselin, M., and McManus, G. B.: Picophytoplankton biomass distribution in the global ocean, Earth Syst. Sci. Data, 4, 37–46, doi:10.5194/essd-4-37-2012, 2012.

Buitenhuis, E. T., Hashioka, T., and Le Quéré, C.: Combined constraints on global ocean primary production using observations and models, Global Biogeochem. Cy., 27, 847–858, 2013a.

Buitenhuis, E. T., Vogt, M., Moriarty, R., Bednaršek, N., Doney, S. C., Leblanc, K., Le Quéré, C., Luo, Y.-W., O'Brien, C., O'Brien, T., Peloquin, J., Schiebel, R., and Swan, C.: MAREDAT: towards a world atlas of MARine Ecosystem DATa, Earth Syst. Sci. Data, 5, 227–239, doi:10.5194/essd-5-227-2013, 2013b.

Casini, M., Hjelm, J., Molinero, J.-C., Lövgren, J., Cardinale, M., Bartolino, V., Belgrano, A., and Kornilovs, G.: Trophic cascades promote threshold-like shifts in pelagic marine ecosystems, P. Natl. Acad. Sci. USA, 106, 197–202, doi:10.1073/pnas.0806649105, 2009.

Chen, B. Z. and Liu, H. B.: Relationships between phytoplankton growth and cell size in surface oceans: interactive effects of temperature, nutrients, and grazing, Limnol. Oceanogr., 55, 965–972, 2010.

Cho, J. C. and Giovannoni, S. J.: Cultivation and growth characteristics of a diverse group of oligotrophic marine Gammaproteobacteria, Appl. Environ. Microbiol., 70, 432–440, 2004.

Caias, P., Sabine, C., Govindasamy, B., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R., Piao, S., and Thornton, P.: Chapter 6: Carbon and other biogeochemical cycles, in: Climate Change 2013 The Physical Science Basis, edited by: Stocker, T., Qin, D., and Plattner, G.-K., Cambridge University Press, Cambridge, 465–570, 2013.

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

da Cunha, L. C., Buitenhuis, E. T., Le Quere, C., Giraud, X., and Ludwig, W.: Potential impact of changes in river nutrient supply on global ocean biogeochemistry, *Global Biogeochem. Cy.*, 21, GB4007, doi:10.1029/2006GB002718, 2007.

Daewel, U., Saetre Hjollo, S., Huret, M., Rubao, J., Maar, M., Niiranen, S., Travers-Trolet, M., Peck, M. A., and van de Wolfshaar, K. E.: Predation control of zooplankton dynamics: a review of observations and models, *ICES J. Mar. Sci.*, 71, 254–271, 2014.

Doney, S. C., Lima, I., Feely, R. A., Glover, D. M., Lindsay, K., Mahowald, N., Moore, J. K., and Wanninkhof, R.: Mechanisms governing interannual variability in upper-ocean inorganic carbon system and air-sea CO₂ fluxes: physical climate and atmospheric dust, *Deep-Sea Res. Pt. II*, 56, 640–655, 2009.

Dubischar, C. D. and Bathmann, U. V.: Grazing impact of copepods and salps on phytoplankton in the Atlantic sector of the Southern ocean, *Deep-Sea Res. Pt. II*, 44, 415–433, doi:10.1016/S0967-0645(96)00064-1, 1997.

Ducklow, H. W.: Microbial services: challenges for microbial ecologists in a changing world, *Aquat. Microb. Ecol.*, 53, 13–19, 2008.

Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benshila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., de Noblet, N., Duvel, J.-P., Ethé, C., Fairhead, L., Fichefet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J.-Y., Guez, L., Guilyardi, E., Hauglustaine, D., Hourdin, F., Idelkadi, A., Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G., Labeyrie, S., Lahellec, A., Lefebvre, M.-P., Lefevre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N., and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, *Clim. Dynam.*, 40, 2123–2165, 2013.

Dutkiewicz, S., Follows, M. J., and Parekh, P.: Interactions of the iron and phosphorus cycles: a three-dimensional model study, *Global Biogeochem. Cy.*, 19, GB1021, doi:10.1029/2004GB002342, 2005.

Eppley, R. W.: Temperature and phytoplankton growth in the sea, *Fisheries Bulletin*, 70, 1063–1085, 1972.

Evans, C., Pearce, I., and Brussaard, C. P. D.: Viral-mediated lysis of microbes and carbon release in the sub-Antarctic and polar frontal zones of the Australian Southern Ocean, *Environ. Microbiol.*, 11, 2924–2934, doi:10.1111/j.1462-2920.2009.02050.x, 2009.

- Fortier, L., Lefevre, J., and Legendre, L.: Export of biogenic carbon to fish and to the deep-ocean – the role of large planktonic microphages, *J. Plankton Res.*, 16, 809–839, doi:10.1093/plankt/16.7.809, 1994.
- Frost, B. W.: The role of grazing in nutrient-rich areas of the open ocean, *Limnol. Oceanogr.*, 39, 1616–1630, 1991.
- Garcia, H. E., Locarnini, R. A., Boyer, T. P., and Antonov, J. I.: World Ocean Atlas 2005, Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Oxygen Saturation, in: NOAA Atlas NESDIS, edited by: Levitus, S., US Government Printing Office, Washington D. C., 342, 2006a.
- Garcia, H. E., Locarnini, R. A., Boyer, T. P., and Antonov, J. I.: World Ocean Atlas 2005, Volume 4: Nutrients (Phosphate, Nitrate, Silicate), in: NOAA Atlas NESDIS, edited by: Levitus, S., US Government Printing Office, Washington DC, 396, 2006b.
- Geider, R. J. and La Roche, J.: The role of iron in phytoplankton photosynthesis, and the potential for iron-limitation of primary productivity in the sea, *Photosynth. Res.*, 39, 275–301, 1994.
- Geider, R. J., MacIntyre, H. L., and Kana, T. M.: Dynamic model of phytoplankton growth and acclimation: responses of the balanced growth rate and the chlorophyll a:carbon ratio to light, nutrient-limitation and temperature, *Mar. Ecol.-Prog. Ser.*, 148, 187–200, 1997.
- Gent, P. R. and McWilliams, J. C.: Isopycnal mixing in the ocean circulation models, *J. Phys. Oceanogr.*, 20, 150–155, 1990.
- Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S. J.: Sources and distributions of dust aerosols simulated with the GOCART model, *J. Geophys. Res.-Atmos.*, 106, 20255–20273, 2001.
- Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T., Kinne, S., Kornblueh, L., Matei, D., Mauritsen, T., and Mikolajewicz, U.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM CMIP5 simulations for the coupled model intercomparison project phase 5, *Journal of Advances in Modeling Earth Systems*, 5, 572–597, 2015.
- Granéli, E., Granéli, W., Rabbani, M., Daugbjerg, N., Fransz, G., Roudy, J., and Alder, V.: The influence of copepod and krill grazing on the species composition of phytoplankton communities from the Scotia Weddell sea, *Polar. Biol.*, 13, 201–213, doi:10.1007/BF00238930, 1993.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

- Gregg, W. W. and Casey, N. W.: Modeling coccolithophores in the global oceans, Deep-Sea Res. Pt. II., 54, 447–477, 2007.
- Gregg, W. W., Ginoux, P., Schopf, P. S., and Casey, N. W.: Phytoplankton and iron: validation of a global three-dimensional ocean biogeochemical model, Deep-Sea Res. Pt. II., 50, 3143–3169, 2003.
- Gruber, N.: The marine nitrogen cycle: Overview of distributions and processes, in: Nitrogen in the Marine Environment, edited by: Capone, D. G., Elsevier, Amsterdam, 1–50, 2008.
- Hansen, P. J., Bjornsen, P. K., and Hansen, B. W.: Zooplankton grazing and growth: scaling within the 2–2000 µm body size range, Limnol. Oceanogr., 42, 687–704, 1997.
- Hashioka, T., Vogt, M., Yamanaka, Y., Le Quéré, C., Buitenhuis, E. T., Aita, M. N., Alvain, S., Bopp, L., Hirata, T., Lima, I., Sailley, S., and Doney, S. C.: Phytoplankton competition during the spring bloom in four plankton functional type models, Biogeosciences, 10, 6833–6850, doi:10.5194/bg-10-6833-2013, 2013.
- Henjes, J., Assmy, P., Klaas, C., Verity, P., and Smetacek, V.: Response of microzooplankton (protists and small copepods) to an iron-induced phytoplankton bloom in the Southern Ocean (EisenEx), Deep-Sea Res. Pt. I., 54, 363–384, doi:10.1016/j.dsr.2006.12.004, 2007.
- Hirst, A. G. and Bunker, A. J.: Growth of marine planktonic copepods: global rates and patterns in relation to chlorophyll *a*, temperature, and body weight, Limnol. Oceanogr., 48, 1988–2010, 2003.
- Hirst, A. G., Roff, J. C., and Lampitt, R. S.: A synthesis of growth rates in marine epipelagic invertebrate zooplankton, Adv. Mar. Biol., 41, 1–142, 2003.
- Hood, R. R., Laws, E. A., Armstrong, R. A., Bates, N. R., Brown, C. W., Carlson, C. A., Chai, F., Doney, S. C., Falkowski, P. G., Feely, R. A., Friedrichs, M. A. M., Landry, M. R., Moore, J. K., Nelson, D. M., Richardson, T. L., Salihoglu, B., Schartau, M., Toole, D. A., and Wiggert, J. D.: Pelagic functional group modeling: progress, challenges and prospects, Deep-Sea Res. Pt. II., 53, 459–512, 2006.
- Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mabhowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global iron connections between desert dust, ocean biogeochemistry, and climate, Science, 308, 67–71, 2005.
- Johnson, R., Strutton, P. G., Wright, S. W., McMinn, A., and Meiners, K. M.: Three improved satellite chlorophyll algorithms for the Southern Ocean, J. Geophys. Res.-Oceans, 118, 3694–3703, doi:10.1002/jgrc.20270, 2013.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Johnson, Z. I., Zinser, E. R., Coe, A., McNulty, N. P., Woodward, E. M. S., and Chisholm, S. W.: Niche partitioning among *Prochlorococcus* ecotypes along ocean-scale environmental gradients, *Science*, 311, 1737–1740, 2006.

Jones, C. D., Hughes, J. K., Bellouin, N., Hardiman, S. C., Jones, G. S., Knight, J., Liddicoat, S., O'Connor, F. M., Andres, R. J., Bell, C., Boo, K.-O., Bozzo, A., Butchart, N., Cadule, P., Corbin, K. D., Doutriaux-Boucher, M., Friedlingstein, P., Gornall, J., Gray, L., Halloran, P. R., Hurt, G., Ingram, W. J., Lamarque, J.-F., Law, R. M., Meinshausen, M., Osprey, S., Palin, E. J., Parsons Chini, L., Raddatz, T., Sanderson, M. G., Sellar, A. A., Schurer, A., Valdes, P., Wood, N., Woodward, S., Yoshioka, M., and Zerroukat, M.: The HadGEM2-ES implementation of CMIP5 centennial simulations, *Geosci. Model Dev.*, 4, 543–570, doi:10.5194/gmd-4-543-2011, 2011.

Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy, C., and Peng, T.-H.: A global ocean carbon climatology: results from Global Data Analysis Project (GLODAP), *Global Biogeochem. Cy.*, 15, GB4031, doi:10.1029/2004GB002247, 2004.

Kishi, M. J., Kashiwai, M., Ware, D. M., Megrey, B. A., Eslinger, D. L., Werner, F. E., Noguchi-Aita, M., Azumaya, T., Fujii, M., Hashimoto, S., Huang, D., Iizumi, H., Ishida, Y., Kang, S., Kantakov, G. A., Kim, H.-C., Komatsu, K., Navrotksy, V. V., Lan Smith, S., Tadokoro, K., Tsuda, A., Yamamura, O., Yamanaka, Y., Yokouchi, K., Yoshie, N., Zhang, J., Zuenko, Y. I., and Zvalinsky, V. I.: NEMURO – introduction to a lower trophic level model for the North Pacific marine ecosystem, *Ecol. Model.*, 202, 12–25, 2007.

Kohfeld, K. E., Le Quere, C., Harrison, S. P., and Anderson, R. F.: Role of marine biology in glacial-interglacial CO₂ cycles, *Science*, 308, 74–78, 2005.

Landry, M. R., Barber, R. T., Bidigare, R. R., Chai, F., Coale, K. H., Dam, H. G., Lewis, M. R., Lindley, S. T., McCarthy, J. J., Roman, M. R., Stoecker, D. K., Verity, P. G., and White, J. R.: Iron and grazing constraints on primary production in the central equatorial Pacific: an EqPac synthesis, *Limnol. Oceanogr.*, 42, 405–418, 1997.

LaRoche, J. and Breitbarth, E.: Importance of the diazotrophs as a source of new nitrogen in the ocean, *J. Sea. Res.*, 53, 67–91, 2005.

Latasa, M., Henjes, J., Scharek, R., Assmy, P., Rottgers, R., and Smetacek, V.: Progressive decoupling between phytoplankton growth and microzooplankton grazing during an iron-induced phytoplankton bloom in the Southern Ocean (EIFEX), *Mar. Ecol.-Prog. Ser.*, 513, 39–50, doi:10.3354/meps10937, 2014.

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

Le Quéré, C. and Pesant, S.: Plankton functional types in a new generation of biogeochemical models: integration of plankton abundance data for the evaluation of marine biogeochemical models, EOS, 90, 30–31, doi:10.1029/2009EO040004, 2008.

Le Quéré, C., Harrison, S. P., Prentice, I. C., Buitenhuis, E. T., Aumont, O., Bopp, L., Claustre, H., Da Cunha, L. C., Geider, R., Giraud, X., Klaas, C., Kohfeld, K. E., Legendre, L., Manizza, M., Platt, T., Rivkin, R. B., Sathyendranath, S., Uitz, J., Watson, A. J., and Wolf-Gladrow, D.: Ecosystem dynamics based on plankton functional types for global ocean biogeochemistry models, Glob. Change Biol., 11, 2016–2040, 2005.

Le Vu, B.: La biocalcification dans l'océan actuel à travers l'organisme modèle *Emiliania huxleyi*, PhD, Université Pierre et Marie Curie, Paris, 2005.

Leblanc, K., Arístegui, J., Armand, L., Assmy, P., Beker, B., Bode, A., Breton, E., Cornet, V., Gibson, J., Gosselin, M.-P., Kopczynska, E., Marshall, H., Peloquin, J., Piontkovski, S., Poulton, A. J., Quéguiner, B., Schiebel, R., Shipe, R., Stefels, J., van Leeuwe, M. A., Varela, M., Widdicombe, C., and Yallop, M.: A global diatom database – abundance, biovolume and biomass in the world ocean, Earth Syst. Sci. Data, 4, 149–165, doi:10.5194/essd-4-149-2012, 2012.

Lee, K.: Global net community production estimated from the annual cycle of surface water total dissolved inorganic carbon, Limnol. Oceanogr., 46, 1287–1297, 2001.

Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., and Garcia, H. E.: World Ocean Atlas 2005, Volume 1: Temperature, US Government Printing Office, Washington, 182, 2006.

Luo, Y.-W., Doney, S. C., Anderson, L. A., Benavides, M., Berman-Frank, I., Bode, A., Bonnet, S., Boström, K. H., Böttjer, D., Capone, D. G., Carpenter, E. J., Chen, Y. L., Church, M. J., Dore, J. E., Falcón, L. I., Fernández, A., Foster, R. A., Furuya, K., Gómez, F., Gundersen, K., Hynes, A. M., Karl, D. M., Kitajima, S., Langlois, R. J., LaRoche, J., Letelier, R. M., Marañón, E., McGillicuddy Jr., D. J., Moisander, P. H., Moore, C. M., Mouríño-Carballido, B., Mulholland, M. R., Needoba, J. A., Orcutt, K. M., Poulton, A. J., Rahav, E., Raimbault, P., Rees, A. P., Riemann, L., Shiozaki, T., Subramaniam, A., Tyrrell, T., Turk-Kubo, K. A., Varela, M., Villareal, T. A., Webb, E. A., White, A. E., Wu, J., and Zehr, J. P.: Database of diazotrophs in global ocean: abundance, biomass and nitrogen fixation rates, Earth Syst. Sci. Data, 4, 47–73, doi:10.5194/essd-4-47-2012, 2012.

Madec, G. and Imbard, M.: A global ocean mesh to overcome the North Pole singularity, Clim. Dynam., 12, 381–388, 1996.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Mahowald, N. M. and Luo, C.: A less dusty future?, *Geophys. Res. Lett.*, 30, 1903, doi:10.1029/2003GL017880, 2003.

Mahowald, N. M., Engelstaedter, S., Luo, C., Sealy, A., Artaxo, P., Benitez-Nelson, C., Bonnet, S., Chen, Y., Chuang, P. Y., Cohen, D. D., Dulac, F., Herut, B., Johansen, A. M., Kubilay, N., Losno, R., Maenhaut, W., Paytan, A., Prospero, J. A., Shank, L. M., and Siefert, R. L.: Atmospheric iron deposition: global distribution, variability, and human perturbations, *Annu. Rev. Mar. Sci.*, 1, 245–278, 2009.

Maier-Reimer, E.: Geochemical cycles in an ocean general-circulation model – preindustrial tracer distributions, *Global Biogeochem. Cy.*, 7, 645–677, 1993.

Maier-Reimer, E., Mikolajewicz, U., and Winguth, A.: Future ocean uptake of CO₂: interaction between ocean circulation and biology, *Clim. Dynam.*, 12, 711–721, 1996.

Martin, J. H.: Glacial-interglacial CO₂ change: the iron hypothesis, *Paleoceanography*, 5, 1–13, 1990.

Martin, P., van der Loeff, M. R., Cassar, N., Vandromme, P., d'Ovidio, F., Stemmann, L., Renagarajan, R., Soares, M., Gonzalez, H. E., Ebersbach, F., Lampitt, R. S., Sanders, R., Barnett, B. A., Smetacek, V., and Naqvi, S. W. A.: Iron fertilization enhanced net community production but not downward particle flux during the Southern Ocean iron fertilization experiment LOHAFEX, *Global Biogeochem. Cy.*, 27, 871–881, doi:10.1002/gbc.20077, 2013.

Moore, J. K. and Doney, S. C.: Iron availability limits the ocean nitrogen inventory stabilizing feedbacks between marine denitrification and nitrogen fixation, *Global Biogeochem. Cy.*, 21, GB2001, doi:10.1029/2006GB002762, 2007.

Moore, J. K., Doney, S. C., and Lindsay, K.: Upper ocean ecosystem dynamics and iron cycling in a global three-dimensional model, *Global Biogeochem. Cy.*, 18, GB4028, doi:10.1029/2004GB002220, 2004.

Moore, J. K., Doney, S. C., Lindsay, K., Mahowald, N., and Michaels, A. F.: Nitrogen fixation amplifies the ocean biogeochemical response to decadal timescale variations in mineral dust deposition, *Tellus B*, 58, 560–572, 2006.

Moore, L. R., Goericke, R., and Chisholm, S. W.: Comparative physiology of *synechococcus* and *prochlorococcus* – influence of light and temperature on growth, pigments, fluorescence and absorptive properties, *Mar. Ecol.-Prog. Ser.*, 116, 259–275, 1995.

Morel, F. M. M.: Kinetics of nutrient uptake and growth in phytoplakton, *J. Phycol.*, 23, 137–150, 1987.

BGD

12, 11935–11985, 2015

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

- Moriarty, R. and O'Brien, T. D.: Distribution of mesozooplankton biomass in the global ocean, *Earth Syst. Sci. Data*, 5, 45–55, doi:10.5194/essd-5-45-2013, 2013.
- Moriarty, R., Buitenhuis, E. T., Le Quéré, C., and Gosselin, M.-P.: Distribution of known macrozooplankton abundance and biomass in the global ocean, *Earth Syst. Sci. Data*, 5, 241–257, doi:10.5194/essd-5-241-2013, 2013.
- O'Brien, C. J., Peloquin, J. A., Vogt, M., Heinle, M., Gruber, N., Ajani, P., Andrleit, H., Arístegui, J., Beaufort, L., Estrada, M., Karentz, D., Kopczyńska, E., Lee, R., Poulton, A. J., Pritchard, T., and Widdicombe, C.: Global marine plankton functional type biomass distributions: coccolithophores, *Earth Syst. Sci. Data*, 5, 259–276, doi:10.5194/essd-5-259-2013, 2013.
- Parekh, P., Follows, M. J., and Boyle, E. A.: Decoupling of iron and phosphate in the global ocean, *Global Biogeochem. Cy.*, 19, GB2020, doi:10.1029/2004GB002280, 2005.
- Platt, T., Broomhead, D. S., Sathyendranath, S., Edwards, A. M., and Murphy, E. J.: Phytoplankton biomass and residual nitrate in the pelagic ecosystem, in: *Proceedings of the Royal Society London A*, Royal Society Publishing London, 459, 1063–1073, 2003a.
- Platt, T., Sathyendranath, S., Edwards, A. M., Broomhead, D. S., and Ulloa, O.: Nitrate supply and demand in the mixed layer of the ocean, *Mar. Ecol.-Prog. Ser.*, 254, 3–9, 2003b.
- Price, N. M., Ahner, B. A., and Morel, F. M. M.: The Equatorial Pacific Ocean: grazer-controlled phytoplankton populations in an iron-limited ecosystem, *Limnol. Oceanogr.*, 39, 520–534, 1994.
- Prowe, A. E. F., Pahlow, M., Dutkiewicz, S., Follows, M. J., and Oschlies, A.: Top-down control of marine phytoplankton diversity in a global ecosystem model, *Prog. Oceanogr.*, 101, 1–13, 2012.
- Rickels, W., Rehdanz, K., and Oschlies, A.: Economic prospects of ocean iron fertilization in an international carbon market, *Resour. Energy. Econ.*, 34, 129–150, 2012.
- Riegman, R., Flaming, I. A., and Noordeloos, A. A. M.: Size-fractionated uptake of ammonium, nitrate and urea and phytoplankton growth in the North Sea during spring 1994, *Mar. Ecol.-Prog. Ser.*, 173, 85–94, 1998.
- Rivkin, R. B. and Legendre, L.: Biogenic carbon cycling in the upper ocean: effects of microbial respiration, *Science*, 291, 2398–2400, 2001.
- Sailley, S. F., Vogt, M., Doney, S. C., Aita, M. N., Bopp, L., Buitenhuis, E. T., Hashioka, T., Lima, I., Le Quéré, C., and Yamanaka, Y.: Comparing food web structures and

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- dynamics across a suite of global marine ecosystem models, *Ecol. Model.*, 261, 43–57, doi:10.1016/j.ecolmodel.2013.04.006, 2013.
- Sarmiento, J. L., Dunne, J., Gnanadesikan, A., Key, R. M., Matsumoto, K., and Slater, R.: A new estimate of the CaCO_3 to organic carbon export ratio, *Global Biogeochem. Cy.*, 16, 1107, doi:10.1029/2002GB001919, 2002.
- Sarthou, G., Timmermans, K. R., Blain, S., and Treguer, P.: Growth physiology and fate of diatoms in the ocean: a review, *J. Sea. Res.*, 53, 25–42, 2005.
- Schlitzer, R.: Export production in the Equatorial and North Pacific derived from dissolved oxygen, nutrient and carbon data, *J. Oceanogr.*, 60, 53–62, 2004.
- Schoemann, V., Becquevert, S., Stefels, J., Rousseau, W., and Lancelot, C.: *Phaeocystis* blooms in the global ocean and their controlling mechanisms: a review, *J. Sea. Res.*, 53, 43–66, 2005.
- Simmons, A., Uppala, S., Dee, D., and Kobayashi, S.: ERAInterim: new ECMWF reanalysis products from 1989 onwards, *ECMWF Newsletter*, 110, 25–35, 2006.
- Smetacek, V., Klaas, C., Strass, V. H., Assmy, P., Montresor, M., Cisewski, B., Savoye, N., Webb, A., d'Ovidio, F., Arrieta, J. M., Bathmann, U., Bellerby, R., Berg, G. M., Croot, P., Gonzalez, S., Henjes, J., Herndl, G. J., Hoffmann, L. J., Leach, H., Losch, M., Mills, M. M., Neill, C., Peeken, I., Röttgers, R., Sachs, O., Sauter, E., Schmidt, M. M., Schwarz, J., Terbrüggen, A., and Wolf-Gladrow, D.: Deep carbon export from a Southern Ocean iron-fertilized diatom bloom, *Nature*, 487, 313–319, 2012.
- Steinberg, D. K., Lomas, M. W., and Cope, J. S.: Long-term increase in mesozooplankton biomass in the Sargasso Sea: linkage to climate and implications for food web dynamics and biogeochemical cycling, *Global Biogeochem. Cy.*, 26, GB1004, doi:10.1029/2010GB004026, 2012.
- Stemmann, L., Picheral, M., and Gorsky, G.: Diel variation in the vertical distribution of particulate matter ($> 0.15 \text{ mm}$) in the NW Mediterranean Sea investigated with the underwater video profiler, *Deep.-Sea Res.*, 47, 505–531, 2000.
- Stock, C. A., Dunne, J. P., and John, J. G.: Drivers of trophic amplification of ocean productivity trends in a changing climate, *Biogeosciences*, 11, 7125–7135, doi:10.5194/bg-11-7125-2014, 2014.
- Straile, D.: Gross growth efficiencies of protozoan and metazoan zooplankton and their dependence of food concentration, predator-prey weight ratio, and taxonomic group, *Limnol. Oceanogr.*, 42, 1375–1385, 1997.

- Sunda, W. G. and Huntsman, S. A.: Iron uptake and growth limitation in oceanic and coastal phytoplankton, *Mar. Chem.*, 50, 189–206, doi:10.1016/0304-4203(95)00035-p, 1995.
- Tagliabue, A., Mtshali, T., Aumont, O., Bowie, A. R., Klunder, M. B., Roychoudhury, A. N., and Swart, S.: A global compilation of dissolved iron measurements: focus on distributions and processes in the Southern Ocean, *Biogeosciences*, 9, 2333–2349, doi:10.5194/bg-9-2333-2012, 2012.
- Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *J. Geophys. Res.*, 106, 7183–7192, 2001.
- Tegen, I., Werner, M., Harrison, S. P., and Kohfeld, K. E.: Relative importance of climate and land use in determining present and future global soil dust emission, *Geophys. Res. Lett.*, 31, L05105, doi:10.1029/2003GL019216, 2004.
- Timmermann, R., Goosse, H., Madec, G., Fichefet, T., Ethe, C., and Duliere, V.: On the representation of high latitude processes in the ORCA-LIM global coupled sea ice–ocean model, *Ocean Model.*, 8, 175–201, 2005.
- Timmermans, K. R., van der Wagt, B., Veldhuis, M. J. W., Maatman, A., and de Baar, H. J. W.: Physiological responses of three species of marine pico-phytoplankton to ammonium, phosphate, iron and light limitation, *J. Sea. Res.*, 53, 109–120, 2005.
- Tollefson, J.: Ocean-fertilization off Canada sparks furore, *Nature*, 490, 458–459, 2012.
- Tréguer, P., Nelson, D. M., Van Bennekom, A. J., DeMaster, D. J., Leynaert, A., and Quéguiner, B.: The silica balance in the world ocean: a reestimate, *Science*, 268, 375–379, 1995.
- Turner, J. T.: Zooplankton fecal pellets, marine snow and sinking phytoplankton blooms, *Aquat. Microb. Ecol.*, 27, 57–102, doi:10.3354/ame027057, 2002.
- Vallina, S. M., Follows, M. J., Dutkiewicz, S., Montoya, J. M., Cermeno, P., and Loreau, M.: Global relationship between phytoplankton diversity and productivity in the ocean, *Nature Communications*, 5, 4299, doi:10.1038/ncomms5299, 2014a.
- Vallina, S. M., Ward, B. A., Dutkiewicz, S., and Follows, M. J.: Maximal feeding with active prey-switching: a kill-the-winner functional response and its effect on global diversity and biogeography, *Prog. Oceanogr.*, 120, 93–109, 2014b.
- Vogt, M., O'Brien, C., Peloquin, J., Schoemann, V., Breton, E., Estrada, M., Gibson, J., Karantz, D., Van Leeuwe, M. A., Stefels, J., Widdicombe, C., and Peperzak, L.: Global marine plankton functional type biomass distributions: *Phaeocystis* spp., *Earth Syst. Sci. Data*, 4, 107–120, doi:10.5194/essd-4-107-2012, 2012.

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Wang, S. and Moore, J. K.: Incorporating *Phaeocystis* into a Southern Ocean ecosystem model, *J. Geophys. Res.*, 116, C01019, doi:10.1029/2009JC005817, 2011.
- Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.*, 97, 7373–7382, 1992.
- Watson, A. J., Bakker, D. C. E., Ridgwell, A. J., Boyd, P. W., and Law, C. S.: Effect of iron supply on Southern Ocean CO₂ uptake and implications for glacial atmospheric CO₂, *Nature*, 407, 730–733, 2000.

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.

Table 1. Growth rates of PFTs at 0 and 20 °C (μ_0 and μ_{20}), and rate increase for a 10 °C increase in temperature (Q_{10}). The uncertainty in μ_0 and Q_{10} represents ± 1 standard deviation from an optimal parameter value in the parameter space. Full references for the phytoplankton growth rate data are provided in the Supplement. The zooplankton growth rate data are from published data synthesis cited here.

PFT	μ_0	Q_{10}	μ_{20}	number of obs.	p values	Size range (μm)	Main references
Autotrophs							
N ₂ -fixers	0.05 ± 0.05	1.83 ± 0.71	0.15	34	0.76	0.5–2.0	LaRoche and Breitbarth (2005)*
Picophytoplankton	0.26 ± 0.06	1.81 ± 0.18	0.89	150	< 0.01	0.7–2.0	Agawin et al. (1998); Johnson et al. (2006); Moore et al. (1995)
Coccolithophores							
Mixed-phytoplankton	0.70 ± 0.17	1.14 ± 0.17	0.90	322	0.06	5–10	Buitenhuis et al. (2008); S. Larsen (this paper)
Diatoms	0.35 ± 0.05	1.57 ± 0.12	0.87	95	< 0.01	2–200	Bissinger et al. (2008)*
Phaeocystis	0.44 ± 0.02	1.93 ± 0.07	1.63	439	< 0.01	20–200	Sarthou et al. (2005)*
Heterotrophs							
Bacteria	0.68 ± 0.07	1.66 ± 0.16	1.87	67	0.23	120–360	Schoemann et al. (2005) ^g
Protozooplankton	0.66 ± 0.04	1.45 ± 0.06	1.22	1429	< 0.01	0.3–1.0	Rivkin and Legendre (2001)*; Cho and Giovannoni (2004)
Mesozooplankton	0.46 ± 0.07	1.48 ± 0.13	1.03	1057	0.01	5–200	Buitenhuis et al. (2010)*
Macrozooplankton	0.31 ± 0.02	1.27 ± 0.05	0.49	2745	< 0.01	200–2000	Hirst and Bunker (2003)*
	0.03 ± 0.01	3.01 ± 0.52	0.19	253	< 0.01	> 2000	Hirst et al. (2003)*

*These references include syntheses of data from other papers.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 2. Model parameters constraining the resource limitations of growth rates. See model equations in Supplement for definitions of parameters.

PFT		Light						Nutrients half saturation ^b					
		α^a	θ_{\max} g Chl g C ⁻¹	Fe^{opt} μmol Fe mol C ⁻¹	k_{Fe} nmol L ⁻¹	k_P μmol L ⁻¹	k_N μmol L ⁻¹						
N ₂ -fixers		1	0.025	8.6	40	0.2	13						
Picophytoplankton		1	0.033	8.6	10	0.13	2						
Coccolithophores		1	0.033	8.6	25	0.13	2						
Mixed-phytoplankton		1	0.033	8.6	25	0.1	2						
Diatoms		1	0.058	8.6	40	0.06	2						
Phaeocystis		1	0.042	8.6	25	0.8	3						
Heterotrophs													
	Food half saturation												
Bacteria			K_{Food} μmol CL ⁻¹										
Protozooplankton			10										
Mesozooplankton			10										
Macrozooplankton			9										

^a Units of mol C g Chl⁻¹ m² (molphotons)⁻¹

^b The reported values are half saturation for uptake for Fe, and half saturation for growth for P and N.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 3. Relative preference of zooplankton for food. The preferences are weighted with the biomass to obtain the model parameter value as in Buitenhuis et al. (2010).

Plankton Functional Type	protozooplankton	mesozooplankton	macrozooplankton
Autotrophs			
N ₂ -fixers	2	0.1	0.1
Picophytoplankton	2	0.75	0.5
Coccolithophores	2	0.75	1
Mixed-phytoplankton	2	0.75	1
Diatoms	1	1	1
Phaeocystis	2	0.75	1
Heterotrophs			
Bacteria	4	0.1	0.1
protozooplankton	0	2	1
Mesozooplankton	0	0	1
Macrozooplankton	0	0	0
Particulate matter			
Small organic particles	0.1	0.1	0.1
Large organic particles	0.1	0.1	0.1

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Table 4. Global mean values for rates and biomass from observations (data) and PlankTOM10 (model) averaged over 1998–2009. The reported confidence level are from the author's assessment of confidence with high: most likely within $\pm 25\%$ of reported value; medium: most likely within $\pm 50\%$ of reported value; low: could be more than $\pm 50\%$ of reported value. For the biomass of phytoplankton and zooplankton, the percentage of the total biomass is also indicated in parentheses (excluding mixed-phytoplankton for which no observations are available).

	model	data	confidence	reference for the data
Rates				
Primary production (PgCy^{-1})	42.4	51–65	high	Buitenhuis et al. (2013b)
Export production at 100 m (PgCy^{-1})	7.6	9–10	medium	Schlitzer (2004); Lee (2001)
CaCO_3 export at 100 m (PgCy^{-1})	0.4	0.6–1.1	medium	Lee (2001); Sarmiento et al. (2002)
SiO_2 export at 100 m (Pg Si)	2.9	3.4	high	Tréguer et al. (1995)
N_2 fixation (TgNy^{-1})	165	60–200	high	Gruber (2008)
Phytoplankton biomass 0–200 m (PgC)*				
N_2 -fixers	0.062 (11 %)	0.008–0.12 (2–8 %)	medium	Luo et al. (2012)
Picophytoplankton	0.21 (38 %)	0.28–0.52 (35–68 %)	medium	Buitenhuis et al. (2012b)
Coccolithophores	0.077 (14 %)	0.001–0.032 (0.2–2 %)	medium	O'Brien et al. (2013)
Mixed-phytoplankton	0.079			
Phaeocystis	0.080 (15 %)	0.11–0.69 (27–46 %)	medium	Vogt et al. (2012)
Diatoms	0.12 (22 %)	0.013–0.75 (3–50 %)	medium	Leblanc et al. (2012)
Heterotrophs biomass 0–200 m (PgC)*				
Bacteria	0.031	0.25–0.26	high	Buitenhuis et al. (2012a)
Protozooplankton	0.067 (13 %)	0.10–0.37 (27–31 %)	medium	Buitenhuis et al. (2010)
Mesozooplankton	0.18 (34 %)	0.21–0.34 (25–66 %)	medium	Moriarty and O'Brien (2013)
Macrozooplankton	0.28 (53 %)	0.010–0.64 (3–47 %)	low	Moriarty et al. (2013)

* The biomass ranges have been computed using the method described in Buitenhuis et al. (2013b).

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.





Figure 1. Schematic representation of the PlankTOM10 (top) and PlankTOM6 (bottom) marine ecosystem models.




Figure 2. Maximum growth rates for 10 Plankton Functional Types (PFTs) as a function of temperature for the phytoplankton PFTs (left) and for the heterotrophic PFTs (right). The PFTs are presented from the smallest (top) to the largest (bottom) in size. The fit to the data used in the model is shown in black, using the parameter values from Table 1. See Table 1 for references.




Figure 3. Covariation between Chl concentration and (left) potentially limiting nutrients and (right) biomass of zooplankton groups for the World Ocean. Chlorophyll data from SeaWiFS satellite are the same in each panel, and are averaged over 1998–2009. The NO₃ and PO₄ data are from the World Ocean Atlas 2009, updated from (Garcia, 2006b). Fe data are from (Tagliabue et al., 2012). The protozooplankton biomass data are updated from Buitenhuis et al. (2010), the mesozooplankton biomass data from Buitenhuis et al. (2006), and the macrozooplankton biomass data include all krill data from Atkinson et al. (2004) and other crustacean data from (Moriarty et al., 2013). All data are monthly averages except for the mesozooplankton, which are seasonal. The black lines are medians, and grey shadings the 25–75 % interquartile range for Chl concentration. The median from the PlankTOM10 model is shown in red.

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.




Figure 4. Surface Chl (mg m^{-3}) for (left) Southern Ocean winter (June–August) and (right) Southern Ocean summer (November–January). Data are from (top) SeaWiFS satellite, (middle) PlankTOM10, and (bottom) PlankTOM6. All datasets are averages for 1998–2009. The boxes highlight the regions used in Fig. 9.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.




Figure 5. Annual mean surface carbon biomasses for individual Plankton Functional Types as simulated by the PlankTOM10 model ($\mu\text{mol CL}^{-1}$).

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.




Figure 6. Dominance of picophytoplankton (top), haptophytes (middle) and diatoms (bottom) in the ocean surface (fraction of time). Left panels show the frequency of the dominance of each PFT detected from satellite data by Alvain et al. (2005) for each pixel during 1998–2006. Right panels show model results, as the surface Chl for each PFT divided by the total Chl. For the model results, picophytoplankton include both the picophytoplankton and N₂-fixers groups; haptophytes include coccolithophores, DMSP-producers and mixed-phytoplankton. The data provides information on the spatial patterns, but not on the absolute amplitude of the dominance. To best highlight the spatial patterns in the model, a PFT is assumed to be dominant if it accounts for at least 45 % of the biomass for picophytoplankton and haptophytes, and 30 % of the biomass for diatoms. The dark red represents area with highest dominance of a PFT, while in the lightest red the PFT is absent.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.




Figure 7. Frequency of blooms of *Phaeocystis* (top) and coccolithophores (bottom) in the surface ocean. *Phaeocystis* data are from Alvain et al. (2005); coccolithophore blooms are updated from Brown and Yoder (1994). A bloom is defined in the model when the PFT accounts for at least 30 % of the biomass and when Chl exceeds $0.3 \text{ mg Chl m}^{-3}$. The dark red represents area with highest dominance of a PFT, while in the lightest red the PFT is absent.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.




Figure 8. Taylor diagram comparing the distributions of surface concentration in annual and monthly mean Chl (Chl and Chl_s), NO₃, PO₄, Si, primary production (pp) and export production (exp) for PlankTOM10 (in grey) and PlankTOM6 (in white) with observations. Chl, biomass and nutrient observations are as in Fig. 3. Export production is from Schlitzer (2004) and represents annual mean flux at 100 m. Primary production is from Buitenhuis et al. (2013) and includes monthly mean values for the surface 300 m. The black dot shows the location where the model results should be if it was perfect and there were no errors in the observations. The distance from the black dot quantifies the performance of the model (Taylor, 2001).

Role of zooplankton dynamics for Southern Ocean phytoplankton biomass

C. Le Quéré et al.




Figure 9. North/South ratio of surface Chl concentration in the Pacific Ocean. Observations are from SeaWiFS. Model results in green correspond to model runs with slow-growing zooplankton: PlankTOM10 (includes macrozooplankton), (Z1) PlankTOM6 plus macrozooplankton, (Z2) PlankTOM6 with mesozooplankton parameterised like macrozooplankton, (D0-D4) PlankTOM10 with no dust deposition or with dust fields from Mahowald et al. (2009), Tegen et al. (2004), Ginoux et al. (2001) and Mahowald et al. (2003), respectively. Model results in blue correspond to model runs without slow growing zooplankton: PlankTOM6, (Z3) PlankTOM10 minus macrozooplankton, (Z4) PlankTOM10 with macrozooplankton parameterised like mesozooplankton, and (D0*-D4*) as above with PlankTOM6. Results from (F1–F3) are model simulations available through the MARine Ecosystem Model Intercomparison Project and (C1–C4) the Climate Model Intercomparison Project 5 (Arora et al., 2011; Dufresne et al., 2015; Giorgetta et al., 2015; Jones et al., 2011). Results from Séférian et al. (2012) mainly differ through their representation of sub-grid scale processes, with improvements in the representation of summer mixed layer depth from Model 1 to Model 3. All data are averaged between 30 and 55 degrees latitude in both hemispheres; 140–240° E in the North and 140–290° E in the South as highlighted in Fig. 1.




Figure 10. North/South ratio of surface Chl concentration as in Fig. 9 vs. the surface biomass of macrozooplankton (PgC yr^{-1}). The standard PlankTOM10 results are shown by the filled circle. Results from ten sensitivity tests are shown by the empty circles, where the maximum growth rate of macrozooplankton is varied within $\pm 2\sigma$ within the range of the data (Fig. 2).

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)