


Accepted Manuscript

Detecting ecological thresholds and tipping points in the natural capital assets of a protected coastal ecosystem

Stephen C.L. Watson, Francis G.C. Grandfield, Roger J.H. Herbert, Adrian C. Newton



PII: S0272-7714(17)31061-2

DOI: [10.1016/j.ecss.2018.10.006](https://doi.org/10.1016/j.ecss.2018.10.006)

Reference: YECSS 5996

To appear in: *Estuarine, Coastal and Shelf Science*

Received Date: 31 October 2017

Revised Date: 20 September 2018

Accepted Date: 9 October 2018

Please cite this article as: Watson, S.C.L., Grandfield, F.G.C., Herbert, R.J.H., Newton, A.C., Detecting ecological thresholds and tipping points in the natural capital assets of a protected coastal ecosystem, *Estuarine, Coastal and Shelf Science* (2018), doi: <https://doi.org/10.1016/j.ecss.2018.10.006>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Detecting Ecological Thresholds and Tipping Points in the Natural Capital Assets of a Protected Coastal Ecosystem.

Stephen C.L. Watson^a, Francis G. C. Grandfield^a, Roger J. H. Herbert^a, Adrian C. Newton^a.

^a Bournemouth University. Faculty of Science and Technology, Centre for Ecology, Environment and Sustainability, Talbot Campus, Poole, Dorset BH12 5BB, UK.

Key words: Thresholds, Tipping points, Marine Protected Areas, Multiple stressors, Natural capital, Poole Harbour.

Abstract

Concern about abrupt and potentially irreversible ecosystem thresholds and tipping points is increasing, as they may have significant implications for natural capital and human wellbeing. Although well established in theory, there are few empirical studies that provide evidence for these phenomena in coastal and estuarine ecosystems, despite their high value for provision of ecosystem services. To determine the likelihood of such events, we tested two statistical methods; sequential T-test analysis (STARS) and generalized additive models (GAMs) in a harbour ecosystem. These methods were applied to time series data spanning up to 25 years coupled with analysis of the relationships between drivers and natural capital asset flows. Results of the STARS analysis identified nonlinear thresholds in three of the natural capital assets of the harbour; mudflat area, Manila clam stocks and wader/wildfowl numbers, as well as an increase in several drivers affecting the harbour. The most prominent threshold was recorded in the Manila clam fisheries of the harbour, with stocks in two locations of the the harbour declining by 73-78% between 2006-2008. We suggest that the historic decline in the Manila clam stocks of the harbour were partly attributable to illegal fishing pressure although other factors such as disease and lease bed holders switching to other species were also likely to have contributed. More recently (2015-onwards) wild clam stocks of the harbour have increased thanks to improved management measures by local authorities. Generalized additive models also identified the contribution of macroalgal mats, sediment shoaling and river flows to historic changes in mudflat area, saltmarsh area and wader/wildfowl numbers. We conclude that information on thresholds and tipping points obtained using these approaches can potentially be of value in a management context, by focusing attention on the interactions and positive feedbacks between drivers that may cause abrupt change in coastal ecosystems.

1 Introduction

Concern about abrupt and potentially irreversible ecosystem transitions is growing rapidly, as they may have significant implications for human wellbeing and are forecast to increase with intensifying climatic change and environmental degradation (Scheffer *et al.*, 2001; Rockström *et al.*, 2009). Such transitions may result from an abrupt change in underlying drivers (e.g. land cover change, nutrient inputs), from an interaction between drivers, or from an abrupt change in the state of the ecosystem with a small or smooth change in drivers (Andersen *et al.*, 2009). Another possibility is a threshold driven by a positive feedback loop, which is often referred to as a tipping point (Scheffer *et al.*, 2009; 2012). While identifying such thresholds and tipping points can be challenging to identify in practice, evidence is increasingly indicating that nonlinear threshold responses could be widespread. Incorporating information about such responses into management plans can facilitate improved management outcomes (Huggett, 2005; Foley *et al.*, 2015). Issues of particular importance to environmental policy and practice include development of techniques to identify where and when thresholds are likely to be encountered (Bestelmeyer *et al.*, 2011; Newton, 2016) and identification

of the underlying mechanisms so that appropriate management responses can be identified (e.g. in the relationships between shorebird mortality and shellfish stock resources; Goss-Custard *et al.*, 2004).

While the importance of ecological thresholds, tipping-points and associated phenomena is increasingly being recognised (e.g. deYoung *et al.*, 2008; Hughes *et al.*, 2013; Levin & Möllmann, 2015), few previous studies have examined their occurrence in transitional systems such as estuaries and harbours (although see Hewitt *et al.*, 2010). This is surprising as such systems typically deliver a number of valuable goods and services (Barbier *et al.*, 2011) but at the same time are subject to more human-induced pressures than most other marine systems (McLusky & Elliott, 2004). In particular, harbours (which may be classified as estuaries or lagoons; Humphreys, 2005) often provide examples of conflicts between high ecological value and intensive human use. The current research was designed to help address this knowledge gap. The purpose of this research was to use a combination of time series data and statistical techniques to examine the occurrence of thresholds and tipping points in Poole Harbour, UK, a Special Protection Area (SPA) of high ecological and socio-economic value. Owing to the breadth of definitions surrounding the concept of tipping points, we start by outlining the definitions adopted here and the underlying theory.

2 Defining tipping points in the natural capital components of ecological systems

Tipping points have been defined in a number of different ways. For example, in their consideration of the Earth's climate system, Lenton *et al.* (2008) defined a tipping point as the critical point at which the future state of the system is qualitatively altered by a small perturbation. Similarly Scheffer *et al.* (2012) referred to a tipping point as a situation where a local perturbation can cause a domino effect resulting in a system transition. Tipping points in complex systems have been widely interpreted as equivalent to critical transitions, phase transitions or fold bifurcations (Lenton *et al.*, 2008; Scheffer *et al.*, 2009; Ashwin *et al.*, 2012). Such concepts derive from theories of dynamical systems, including bifurcation and catastrophe theories. Application of these theories has highlighted a number of ways in which tipping points can occur, for example by a change in the external conditions of a system, or a change in the state of the system itself (Ashwin *et al.*, 2012, van Nes *et al.*, 2016).

While application of dynamical systems theory to the climate system is now well established (Lenton *et al.*, 2008), its application to understand the dynamics of terrestrial and marine ecosystems has been the focus of some debate. Policy makers and land managers increasingly want to understand how different forms of environmental change might affect the condition of natural capital (NC), and the flow of multiple ecosystem services (ES) to human society (Mace *et al.*, 2015). As dynamical systems models are typically defined in relation to a single independent variable, simultaneous consideration of multiple and potentially interacting drivers of ecological change represents a significant analytical challenge. As noted by Donahue *et al.* (2016), the multidimensionality of ecological responses requires explicit consideration of multidimensional disturbances or causes of change. The challenges of applying dynamical systems theory to real-world ecosystems are illustrated by the concept of ecological resilience. Much of the recent literature on this concept is based on the assumption that ecosystems have multiple stable equilibria, with tipping points occurring between them (Donahue *et al.*, 2016). Definitions of ecological resilience focus on the capacity of a system to maintain its essential structure and function when confronted with external perturbations (Quinlan *et al.*, 2016). Yet the empirical evidence for the existence of such multiple stable states is very limited (Petraitis, 2013); most ecosystems are far from the equilibria assumed by theory (Donahue *et al.*, 2016), and other assumptions on which the underlying theory is based are often not met in field situations (Newton, 2016). Consequently, ecological resilience has proved very difficult to measure in practice (Quinlan *et al.*, 2016, Biggs *et al.*, 2012, Cantarello *et al.*, 2017).

Together with the semantic confusion surrounding resilience, these problems have resulted in the concept being misapplied in both policy and practice (Newton, 2016).

We therefore follow van Nes *et al.* (2016) in applying the term ‘tipping point’ to any situation where accelerating change caused by a positive feedback drives the system to a new state. We make no assumptions about whether the ecosystem in question is characterised by the existence of multiple stable states (Petraitis, 2013), and we do not make an explicit link between tipping points and dynamical systems theory. As highlighted by van Nes *et al.* (2016), this broader definition of a tipping point is consistent with the work of Gladwell (2000), who did so much to popularize the concept. The existence of an intrinsic positive feedback process that drives accelerating change differentiates concept tipping point from a broader category of abrupt ecosystem change, which we refer to as an ecological threshold. Any situation where there is an abrupt change in ecosystem structure or function can be considered as an ecological threshold (Groffman *et al.*, 2006). Ecological thresholds may also usefully be differentiated from decision or management thresholds, or regulatory limits (Johnson, 2013), which are based on values of system state variables that should prompt specific management actions (Martin *et al.*, 2009). Following van Nes *et al.* (2016), we therefore restrict the term ‘tipping point’ to a subcategory of ecological threshold where the abrupt change is driven by a positive feedback mechanism.

Here we examine the occurrence of thresholds and tipping points in relation to provision of multiple ecosystem services in a coastal ecosystem. To achieve this, we employ a conceptual framework based on the reviews conducted by Mace *et al.* (2015) and the Natural Capital Committee (NCC, 2014). Here, NC is defined as assets, stocks or the elements of nature that directly and indirectly produce value or benefits to people (NCC, 2014), such as ecological communities or habitat types. Following Mace *et al.* (2015), the status of these natural assets can be measured using metrics of the area, and condition of these communities. In the context of environmental degradation and its potential impact on human society, the form of the relationship between the condition of a natural asset and provision of benefits is of particular importance. Environmental degradation may lead to a decline in natural asset status, which will reduce the benefits provided to people. The form of this decline represents a key knowledge gap (Folke *et al.*, 2011; NCC, 2014), but could potentially include threshold responses or tipping points (Figure 1 (I)). In addition, we hypothesize that the relationship between anthropogenic drivers (or pressures) and NC status may also demonstrate a threshold response or a tipping point (Figure 1 (II,III,IV)).

The relationships between anthropogenic drivers (or pressures) and NC status may also vary over time, demonstrating either linear or nonlinear trends (Figure 1 (V-VII)). If an environmental driver intensified over time, then it could produce a threshold response in NC status, or a tipping point if a positive feedback mechanism were influential. Tipping events (IV & VII) are often considered difficult to reverse because of a phenomenon known as hysteresis (Meyer, 2016). This implies that the system cannot recover by retracing the path followed during degradation. Instead, the environmental driver that caused the transition has to be reduced further than the threshold value that caused the initial transition. Ultimately, if environmental degradation leads to an abrupt decline in natural asset status, this will reduce the benefits provided to people, either temporarily or permanently.




Figure 1: (I) Alternative forms of forms of natural capital asset–benefit relationships, as hypothesized by Mace *et al.* (2015). The solid black line illustrates how the value of benefits might change in response to variation in the status or condition of natural assets, which could be caused by environmental degradation. The dashed line shows a threshold response (or tipping point). Panels (II–IV) show the relationship between NC status to changing conditions or environmental drivers which might be: II. Linear response. III. Nonlinear, non-hysteretic response of ecosystem state as a function of a pressure (threshold) or IV. Tipping point (hysteretic), representing a nonlinear change driven by an intrinsic positive feedback mechanism and with respect to changing conditions or environmental drivers. Finally, panels (V–VII) show how a responding system may change through time when they respond to an escalating driver according to the linear or abrupt equilibrial behaviour shown in (II–IV).

3 Methods

3.1 Details of study area: Poole Harbour

Poole Harbour is a large natural harbour of nearly 4,000 ha (Underhill-Day, 2006) located on the coast of Dorset in southern England (Lat. 50° 42' 44" Long. 2° 03' 30" W) in the United Kingdom (Figure 2). Although classified as an estuary (as several rivers flow into it), Poole Harbour has many of the qualities of a large lagoon, owing to the narrow entrance and limited tidal range (Humphreys, 2005). A diverse set of habitats from saltmarsh and reedbed (*Phragmites australis*) to valley mire and lowland heathland provide a host of different ecosystem services such as recreation, coastal protection and increased water quality to a catchment of over 142,100 people (Office for National Statistics, 2010). Ecologically, the intertidal mudflats, sandflats and marshes support large numbers of wintering wildfowl and waders that are of national and international significance. The harbour and its adjacent landscape also hold a number of other national statutory designations that serve to protect the natural environment, including being classified as a Site of Special Scientific Interest (SSSI), a Special Protection Area (SPA) designated under the EU Birds Directive and a Ramsar site. Under the EC Shellfish Waters Directive, Poole Harbour (with the exception of Holes Bay) is also designated as a shellfish water and is the location of fishing and aquaculture activities, which in 2005 were worth in excess of £2 million per year to the local economy (Jensen *et al.*, 2004). However, despite its high economic and conservation value, the occurrence of ecological thresholds and tipping points in the NC assets of Poole Harbour has not been examined previously.




Figure 2 Map of Poole Harbour ©Crown Copyright and database right (2010) Ordnance Survey Licence Number 1000022021. Open water, Saltmarsh & Sediment data from East Dorset Habitat map© Environment Agency, 2010.

3.2 Data collection

Data for four different categories of NC components were gathered for the period 1980-2015 (Table 1). Three NC stocks of interest (mudflat area, saltmarsh area and wader/wildfowl numbers) were chosen owing to their immediate importance for conservation within the SPA, while the potential stocks of the Manila clam in the harbour (*Ruditapes philippinarum*) were also investigated based on their significant commercial importance and the potential benefit flows provided by the landings of clams into Poole Harbour. To test potential pressure-state relationships, data for possible drivers in

the harbour were sourced from the literature, environmental data-bases and monitored instrument records (Table 2). For example we used tidal river flow and water quality data from the River Frome at East Stoke gauging station (ID: 44207) to represent a county level watershed driver. In the absence of long-term fishing effort data (e.g. fishing effort, frequency trawled) fleet capacity (i.e. number of licenced clam boats) was used as a proxy for fishing pressure (Piet *et al.*, 2006). As fishermen in Poole Harbour utilise a unique “pump-scoop” dredge to harvest the Manila clam (95% of catch is typically clam landings; Clarke *et al.*, 2017) fleet capacity is likely an effective pressure indicator that describes the impact induced by fishing activities on the system.

Table 1: Proxies used for assessing natural capital assets (stocks) in Poole Harbour.

Natural capital assets (stock)	Potential ecosystem services	Indicator	Time series	Data source
Intertidal mudflat (area)	Carbon storage, (Regulating) Marine invertebrate habitat (Supporting/Habitat)	Area of mudflat and other littoral sediment (excluding saltmarsh and macroalgal mats) in Poole Harbour as a whole (ha). Areas derived from aerial photography, Compact Airborne Spectrographic Imaging and direct survey.	1980-2015	Environment Agency field data. (Bryan <i>et al.</i> , 2013).
Manila clam stocks (<i>Ruditapes philippinarum</i>)	Food (Provisioning) Nutrient cycling (Regulating).	Annual stock surveys for Manila clam were obtained for three sites in the harbour: Arne Bay, Seagull Island and Round Island. Samples were collected at each site using a trailed pump scoop dredge which was towed along the seabed in circular motions for two minutes. During these two minutes the number of rotations made by the vessel was recorded. The dredge was then lifted aboard the vessel and the contents were emptied into a sample bucket. Three replicate samples were taken at each site and the mean density ($N.m^{-2}$) of each catch recorded.	2003-2015	Southern Inshore Fisheries and Conservation Authority (IFCA) field data. The methodology is described in: SIFCA, (2017).
Saltmarsh (area)	Nutrient cycling and coastal protection (Regulating), Marine invertebrate habitat (Supporting/Habitat)	Trends in saltmarsh area (ha) in Poole Harbour derived from aerial photography, Compact Airborne Spectrographic Imaging.	1980-2013	Raybould (2005); Gardiner (2015).
Wildfowl and waders	Birdwatching (Cultural)	The harbour wide average density of all species of wildfowl and waders known per year (N).	1980-2015	Wetland Bird Survey (WeBS) data.

Table 2: Indicators of environmental drivers selected for analysis in the Poole Harbour system.

Drivers	Indicator	Time series	Data source
Fishing pressure (Manila clam <i>Ruditapes philippinarum</i>)	Number of licenced Manila clam boats in Poole Harbour. Clams are removed from the seabed using a pump scoop dredge which is towed along the seabed by small (under 10 m) fishing vessels.	1994-2015	Information on fishing activity has been obtained with consultation from a range of sources and organisations including Southern IFCA, and Poole Harbour Commissioner reports (Simpson 2004).
Macroalgal mats (area)	Areas of macroalgal mats (ha) on mudflat and other littoral sediment (excluding saltmarsh) with $\geq 75\%$ cover and $> 2 \text{ kgm}^{-2}$ biomass (ha) in Poole Harbour as a whole. Areas derived from aerial photography, Compact Airborne Spectrographic Imaging and direct survey.	1980-2015	Environment Agency field data (Bryan <i>et al.</i> , 2013)
Nutrient loading (Nitrates)	Dissolved nitrate concentration ($\text{mg NO}_3\text{-N l}^{-1}$)	1980-2015	River Frome at East Stoke - Centre for Ecology & Hydrology, & FBA (Freshwater Biological Association); Bowes <i>et al.</i> (2011).
Nutrient loading (Phosphates)	Soluble reactive phosphorus concentration ($\mu\text{g l}^{-1}$)	1980-2015	River Frome at East Stoke - Centre for Ecology & Hydrology, & FBA (Freshwater Biological Association); Bowes <i>et al.</i> (2011)
Riparian water flows.	Mean annual river flow (m^3s^{-1}) within the Frome and Piddle rivers.	1980-2015	National River Flow Archive; The Centre for Ecology & Hydrology (CEH)
Sediment shoaling	Mean channel depth (m) Wareham Channel.	1980-2015	Poole Harbour Commissioners (PHC); Raybould (2005)
Water temperature	Monthly recorded sea surface temperatures were averaged across the Poole Harbour time series data ($^{\circ}\text{C}$)	1980-2015	Cefas Coastal Temperature Network Station 23; Channel Coastal Observatory from 2011.

3.3 Data analysis

Based on criteria outlined by Collie *et al.* (2004), Bestelmeyer *et al.* (2011), Carpenter (2011) and Samhouri *et al.* (2017) we followed a step-wise process for detecting and characterising thresholds and their driver-response interactions. The workflow can be summarised in three parts: (1) explore the potential for nonlinear relationships in the time series data, (2) determine appropriate pressure-state relationships, and (3) identify any pressure-state thresholds and the location (inflection point) and strength of the thresholds. Before any analysis was conducted, we normalised each set of ecological and environmental time series data by subtracting the mean and scaling by the standard deviation. Where necessary, we averaged intra-annual measures to create a single annual time series for each variable, noting that this may increase the possibility of detecting significant thresholds and tipping points (Samhouri *et al.*, 2017).

The first step was to locate and statistically test one or more breakpoints in time series data with the purpose of identifying the potential existence of nonlinear thresholds occurring over time. Significant breakpoints in each time-series data set (Table 1 and 2) were identified by performing a sequential analysis of mean values using the sequential T-test analysis (STARS) method (Rodionov, 2004). The STARS algorithm was set to detect significant ($p \leq 0.01$) shifts in the mean value and the magnitude of fluctuations in the time series data by using a modified two-sided Student's t-test. Three different cut-off lengths ($l = 5$, $l = 10$ and $l = 15$) were used to test the sensitivity of results obtained from STARS analyses. Tipping points are often associated with short periods of variability and so an initial cut-off length of 5 was chosen.

To determine appropriate pressure-state relationships, model selection tests were then carried out using stepwise generalised additive models (GAMs) performed using R 3.4.5 statistical software (R Development Core Team, 2016). Similar techniques have successfully been used to detect threshold responses in ecological data (Large *et al.*, 2013) as they are non-parametric and capable of modelling nonlinear responses. They are robust and more flexible than linear methods when using unequally

spaced data (Large *et al.*, 2013), while offering a robust approach for detecting threshold responses (Toms & Villard, 2015). As change in one element of NC stocks can either directly or indirectly affect the dependence of other NC stocks or their associated benefit flows (Beaumont *et al.*, 2008), we also tested interrelationships between these variables. For example, biomass of invertebrates in mudflat often provides an important food source for waders and wildfowl, thus any change in a mudflats total area may affect such populations.

For statistically significant pressure-state relationships ($p \leq 0.01$), we fitted separate generalised additive models (GAMs) in R to test for nonlinearities. A smoothing function was applied to each explanatory variable. If smoothing functions are not properly fitted in the model, complex overfitting is likely to result. To minimise this risk, we used integrated model cross-validation algorithms to ensure that the models selected were as robust as possible (Rodionov & Overland, 2005). An eigenvalue optimisation process was carried out to prevent overfitting using the “mgcv” package in R (Wood, 2011). Generalised cross validation (GCV) was used to estimate a smoothing parameter for each term. Smoothing terms with penalised regression splines with an added penalty for each term were used so that the number of knots (the x-value at which the two pieces of the model connect) for each term could be reduced to zero. Through this eigenvalue optimization process, smoothing terms with linear functions in response to pressure variables could effectively be removed from the model if it did not improve the fit (Wood, 2004). As the goal of this research was to identify possible nonlinear threshold values that can inform decision criteria, we rejected GAM models that were more adequately explained using a linear model (Wood & Augustin, 2002). Model selection tests using Akaike's Information Criterion (AIC) were performed on GAMs with different knot combinations to find the knot allocation that resulted in the best fit to the data. The relative importance or explained variance (R^2) of each pressure-state variable in the regression model was calculated and checked using the LMG metric with the relaimpo package in R (Groemping, 2007). From this analysis, we calculated 95% confidence intervals *via* bootstrapping of the residuals in order to allow for autocorrelation (Vinod & López-de-Lacalle, 2009). This procedure generated a range of pressure-state values where a GAMs smoothing function changes trajectory and indicates where threshold might occur. Quantitative estimates of a threshold were defined as the point of inflection where the second derivative changes sign (e.g. Samhouri *et al.*, 2010, Large *et al.*, 2013; 2015).

4 Results

4.1: Time-series trends, thresholds and ecosystem responses

Breakpoint (STARS) analysis of the time series data available for Poole Harbour provided empirical evidence of recent environmental degradation in three of the four NC assets: mudflat area, saltmarsh area and Manila clam stocks (Figure 3). A brief description of the results for each NC asset follows, along with the results from the assembled driver data (Table 3), (see also Appendix 1).

Following their introduction to the harbour in the late 1980's Manila clam stocks increased considerably in the Arne and Seagull Island areas between 2003 and 2007, but experienced strong abrupt shifts between 2007 and 2008 with reduced clam densities persisting at below pre 2006 mean values until 2014. Results of the STARS algorithm (Table 3) suggest that the magnitude of the changes detected in 2008 were the greatest of any variable tested (~3.18-3.26 respectively). There were however, strong indications of recovery in recorded stocks at these sites in 2015, with significant breakpoints ranging in magnitude from (~ 2.93-3.02 respectively). Stocks of Manila clam recorded at Round Island have conversely increased steadily across the time period, with little signs of abrupt changes. Towards the intertidal areas of the harbour, the mudflats and saltmarshes both showed significant signs of erosion across their respective time periods. The decline in mudflat area

over the twenty-five year interval was the more pronounced of the two assets, declining by up to two standard deviations away from the mean value in 1980. Over this time interval, saltmarsh area declined for the first decade then remained relatively stable. This was associated with an increase in mudflat area from 1988 until the mid-1990s, values declining thereafter. Populations of waders and wildfowl increased after 1980 reaching a peak in the mid-1990s, thereafter declining such that by 2005-2010 values were close to those encountered in the early 1980s. Since then, numbers have increased somewhat. It should be noted that these trends only give a “snapshot” of the overall status of the resident bird populations and do not reveal trends for individual species.

The highest STARS value for the driver data was obtained for phosphate values in the harbour (1.95) which have declined considerably since the 1980's. The second strongest shift in the drivers (1.46 & 1.86) was marked by an increase in macroalgal mats across the harbour between 1996 and 2010, followed by a marginal decline from 2011-2015 (Appendix 1). Changes in nitrate concentrations and the water temperature both showed increasing trends over the multidecadal period, leading towards a catchment with a high eutrophic status. River flow trends for the catchment also indicate a year on year increase in flow rate. A single low STARS value (0.12) was detected for sediment shoaling in our proxy site of the Wareham channel, with sediment initially increasing the depth of the channel between 1980 and 1995, before crossing a threshold and thereafter decreasing channel depth. A plausible shift in fishing pressure in 2004 and 2008 can also be seen, coinciding with a decline in Manila clam stocks (Figure 3).




Figure 3: STARS threshold detection of the four normalised natural capital assets in Poole Harbour, Manila clam stocks ($N m^2$), mudflat area (excluding saltmarsh and macroalgal mats) (ha), saltmarsh area (ha) and waders/wildfowl (no. individuals). The horizontal line(s) indicates the direction (positive or negative) of the trend representing a significant deviation from zero (i.e. the proxy mean over the time period). Vertical black lines represent statistically significant ($p \leq 0.01$) breakpoints for individual trends from sequential Student's t-tests.

Table 3: Summary of the STARS index values of the environmental drivers and natural assets (stocks).

Drivers/Natural capital stocks	Best estimate of threshold: Time series (STARS)	Magnitude of responses (STARS)
Fishing pressure	2004, 2007	1.78, 1.42
Macroalgal mats (area)	1989, 1996, 2010	0.85, 1.46, 1.86
Nitrates	1996, 2005, 2008	0.34, 0.32, 0.98
Phosphates	2011	1.95
River flow	N/A	N/A
Sediment shoaling	1996	0.12
Water temperature	1985, 1989	0.27, 0.56,
Manila clam stocks (Arne)	2006, 2008, 2014	2.26, 3.18, 2.93
Manila clam stocks (Seagull)	2006, 2008, 2014	2.25, 3.26, 3.02
Manila clam stocks (Round)	-	-
Mudflat excluding saltmarsh and macroalgal mats (area)	1990, 2000, 2007	0.26, 0.65, 0.62
Saltmarsh (area)	1986, 1992	0.54, 0.67
Waders and wildfowl	1991, 2006, 2012	1.59, 1.83, 1.76

4.2 The relative contribution of multiple pressures to natural capital stocks

Based on multi-model inference with GAMs we quantified the relative importance of environmental variables to influence each of the four selected NC stocks. Of the thirty-six possible GAM models, ten were significant (Table 4) with the smoothing function included ($p \leq 0.01$).

Table 4: p-values for all GAM models analysed. Significant models ($p \leq 0.01$) are shown in bold and with an (*). **Mudflat area excludes saltmarsh and macroalgal mats.

Natural capital stocks	Drivers								
	Fishing pressure	Mudflat (area)	Macroalgal mats (area)	Nitrates	Phosphates	Saltmarsh (area)	Sediment shoaling	River flow	Water temperature
Mudflat (area)**	N/A	N/A	0.0032*	N/A	N/A	0.377	0.0051*	0.002*	0.265
Manila clam Arne (Stocks)	0.003*	N/A	N/A	0.06	0.09	N/A	0.262	0.308	0.38
Manila clam Seagull Island (Stocks)	0.002*	N/A	N/A	0.05	0.10	N/A	0.231	0.301	0.41
Manila clam Round Island (Stocks)	0.067	N/A	N/A	0.16	0.18	N/A	0.276	0.453	0.24
Saltmarsh (area)	N/A	0.377	0.0017*	0.747	0.472	N/A	0.0027*	0.0021*	0.497
Waders and wildfowl	N/A	0.072	0.0061*	0.678	0.965	0.0051*	0.1390	N/A	N/A

We found that macroalgal mats (area), sediment shoaling and river flow were the most important predictors for explaining the variability in area of both mudflats and saltmarsh. This finding is confirmed based on the r^2 evidence ratio (Figure 4) with the three covariates explaining 91% and 85% of the total variance of each model respectively. Macroalgal mats and saltmarsh were the most important predictors of wader and wildfowl stocks with a relative importance of 0.21% and 0.18% and were significant at $p \leq 0.01$. Although mudflat area and sediment shoaling were not significant for determining wader and wildfowl stocks, they had a high relative importance in explaining the variability of the final models (0.13-0.15%). Fishing pressure was the only significant ($p \leq 0.01$) predictor of Manila clam stocks in Arne Bay and Seagull Island with a relative importance of ~84% respectively. As no environmental variables were significant in influencing the Round Island Manila clam populations, we removed this time series from the next step of full GAM analysis. Other variables were less important for all indices, ranging from 0.01 to a relative importance of 0.13 (see Figure 4).




Figure 4: Relative importance of different pressures for each of the natural capital stock models. The proportion of variance explained by the final model(s) was: mudflat area (99.16%), Manila clam stocks for Arne and Seagull Island (~97.6%), saltmarsh area (86.90%) and waders/wildfowl (76.54%). *Mudflat area excludes saltmarsh and macroalgal mats.

The full GAM analyses allowed identification of relationships between NC status and significant pressures. Macroalgal mats showed evidence for negative nonlinear relationships (Figure 5) with three NC proxies namely mudflat area, saltmarsh area and numbers of wading birds. Sediment shoaling generally increased with mudflat area and a significant positive trend was observed at a value of ~ -0.9 (SD). Saltmarsh vs sediment shoaling also showed an increasing trend before crossing a threshold at ~ -0.9 (SD) and then decreasing to below its initial value. Mudflat area also showed a negative nonlinear relationship with river flow, with a clear threshold observed $\sim 0.2-0$ (SD). The relationship between saltmarsh area and river flow was best described as a hockey stick, such that

saltmarsh area was negatively associated with river flow at values < -0.2 (SD), but then inverted to a positive trend when river flow was not significantly different from zero. As macroalgal mat area increased wader and wildfowl numbers decreased, particularly at higher values of the former, with a threshold response evident at $\sim 0.08-0.05(\text{SD})$ for both pressure-states. Similarly there was a generally negative relationship between wader and wildfowl numbers and saltmarsh area, with a threshold again detected at around -0.5 (SD). There was no evidence for nonlinear responses or thresholds in Manila clam stocks in response to fishing pressure in either Arne Bay or Seagull Island, suggesting a purely linear relationship between the variables. Overall, of the three proxies for NC stocks with nonlinear responses, all three showed evidence for thresholds in relation to more than one pressure.




Figure 5 GAMs of the four normalised natural capital stocks response to pressures ($p \leq 0.01$), where the horizontal black line represents significant positive or negative trends, representing a significant deviation from zero (i.e. the mean). The grey polygon represents 95% confidence intervals and red dotted arrow indicates the best estimate of the location of a threshold (i.e., where the second derivative is most different from zero within the threshold range). *Mudflat area excludes saltmarsh and macroalgal mats.

5 Discussion

In this study, we employed STARS and generalised additive models (GAMs) to identify trends and thresholds in pressure-time series relationships. Using this analysis we identified distinct points where four NC assets of the harbour (Manila clam stocks, mudflat area, saltmarsh area and waders/wildfowl numbers) have been substantially changed in the past, and the potential drivers of that may have caused such variabilities. Although the STARS technique has been previously been used to identify thresholds in ecological time series data (Moellmann *et al.*, 2009; Conversi *et al.*, 2010), the present study is the first to employ this method to empirically identify thresholds within a

NC framework and one of only a few studies to use such analysis in a transitional estuarine system (e.g, Chevillot *et al.*, 2016).

5.1 Trends, thresholds, and fundamental features from STARS analysis

In applying STARS to available drivers for the Poole Harbour ecosystem, the following picture emerges. The 1980-2015 period was categorised by three steadily increasing endogenous pressures (i.e. emanating from the surrounding catchment and within the system; Elliott, 2011) including nitrate concentrations, macroalgal mats and river flows. With respect to these drivers, nitrate loading, a common driver of algal growth and water quality (McGlathery *et al.*, 2007; Lyons *et al.*, 2014), has shifted the estuarine watershed beyond the long-term safe loading limits determined by the Water Framework Directive for the catchment, leading towards an “unfavourable-bad” eutrophic status (Howarth & Marino, 2006; Conley *et al.*, 2009). The current Nitrogen Reduction Strategy (Kite *et al.*, 2012) for the catchment identifies the main source of nitrogen to be diffuse agricultural inputs (73%) with nitrogen entering the harbour forecast to rise further over the next few decades. This is owing to a lag effect of nitrogen leaving the riparian soil zone of surrounding agricultural land and entering the harbour. The consequences of crossing this threshold are likely to be the continued expansion of macroalgal mats fuelled by rising concentrations of nitrate and other inorganic nitrogen compounds in harbour waters. These effects could be compounded by the observed rise in river flow levels since the 1980’s, which may act to convey more nitrogen into the harbour owing to the poor flushing characteristics of the Harbour (Dyrynda, 2005). In contrast, phosphate concentrations entering the harbour have decreased substantially since the 1980’s. This is likely due to substantial land use changes and improvements to phosphorous stripping sewage treatment processes in the catchments of the two main rivers (Frome and Piddle) discharging into Poole Harbour. Evidence of a shift in sediment shoaling in the Wareham channel and fishing pressure on Manila clam populations occurred about the time as the dramatic declines in Manila clam landings (2004-2007). Results from the STARS analysis suggest that the magnitude of the changes in sediment in the Wareham channel were relatively minor, concurring with reports that since 1980 many channels have deepened in most parts of the harbour (May, 2005). In this study we only considered one exogenous pressure (i.e. those emanating from outside the system; Elliott 2011), in the form of water temperature, which showed evidence of a shift to warmer waters around 1989. Over recent decades, an increase in temperature and associated changes in precipitation and sea level rise have been observed in Europe as well as other parts of the world (Pachauri *et al.*, 2014) and it expected such trends will continue in the future.

Among the NC proxies of the harbour, several significant thresholds were identified in the time series data. Relating these changes back to our conceptual framework outlined in Figure 1, STARS results here show saltmarsh area of the harbour to have declined linearly (Type V) between 1980-1988 before stabilising since 1994 at ~400 ha. Longer term trends (1890-2013) in the saltmarsh species *Spartina anglica* by Gardiner (2015) describe the rapid colonisation of the perennial grass over the mudflats between 1890 and 1924 before passing a threshold, and since then there has been much loss of *Spartina* across the harbour. Despite evidence here that this degradation may have now ceased, there is local evidence (e.g. in Holes Bay) that show *Spartina* is still receding in some locations (Gardiner *et al.*, 2007).

Trends of waders/wildfowl and mudflat area in the harbour both exhibited abrupt thresholds (Type VI) at the estuarine scale with the most abrupt threshold response taking place in bird numbers between 2007 and 2012. Irrespective of such abrupt shifts, as of 2012-2015, bird numbers of the harbour were higher with those of the 1980’s but lower than the beginning of the 1990’s. One possible reason for a general increase in bird numbers in the early 1990’s as suggested by Raybould

(2005) could be the larger invertebrate prey base opened up in the form of increasing area of mudflats as saltmarsh receded. Evidence from STARS analysis also suggest that the decline in bird numbers since the early 1990's could be related to the decline in total mudflat area around the same time (1994), likely as a direct result of mudflats becoming increasingly covered by macroalgal mats. The spread of macroalgae on mudflats has been implicated in the decline of wader/wildfowl populations in many British estuaries (Tubbs & Tubbs, 1980; Anders *et al.*, 2009) including Poole Harbour (Jones & Pinn, 2006), owing to its impact on invertebrates when macroalgal wet weight biomass reaches 2 kg m^{-2} (Raffaelli *et al.*, 1991; 1999). Indeed, recent evidence presented by Thornton (2016) based on field experiments conducted in Poole Harbour, suggests that bird species preferred prey under lower macroalgal mat biomass ($\sim 800 \text{ g m}^{-2}$ wet weight), supporting a lowering of the current legislative threshold of 2 kg m^{-2} to 1 kg m^{-2} . As the condition of mudflats, wading birds and the extent of algal mats are sanctions under current legislation (JNCC 2004) for Poole Harbour, is important to be able to reliably assess the impact from macroalgal mats on these NC assets.

In the Manila clam fishery of the harbour, free-living stocks in the harbour were shown to have generally increased considerably since their introduction in the late 80's. However, an abrupt decline in the densities of clams was observed at two sites of the harbour between 2006 and 2008, and since the values have only recently (2015) shown signs of recovery. These changes were also the greatest in magnitude of all the threshold responses observed in this study and fit the criteria outlined in Figure 1 for a tipping point transition (i.e. type VII). However, at this point STARS analysis could only provide qualitative evidence of the impact of drivers and potential feedback mechanisms on the time series data. To quantitatively unravel the relative importance of different drivers as well as potential feedback mechanisms (which are a prerequisite of a tipping point), we considered the results from the GAMs, as explored further below. Moreover, in attempting to reconcile the difference between clam stocks at various locations of the harbour we recognise that the three sites investigated here are only a snapshot of the local populations. However, our results corroborate with other larger studies (e.g. Herbert *et al.*, (2018)) that have looked at several sites across the harbour (including our sites, albeit with different protocols), which suggest a fall in the number of Manila clams at the harbour level between 2008-2009 (26 to 11 ind per m^2), but a relatively stability in clam numbers from 2011-2015 at approximately 25-28 ind per m^2 , based on a resurvey of a selected number of the original sites.

5.2 The impact of multiple stressors on natural capital stocks

By means of multi-model inference, we were able to determine statistically the relative contribution of fishing pressure, macroalgal mats, nitrates, phosphates, river flows, sediment shoaling and elevated water temperatures to the dynamics of four NC assets of Poole Harbour. This is important information for the management of the harbour, because any thresholds identified by asset-driver-state interactions indicate where particular management interventions might be needed to avoid abrupt changes occurring. However, the models that we generated in this research did not take into account the complex interactions that may occur between driver variables (e.g. Crain *et al.*, 2008), and we may have missed important drivers from the analysis (e.g. sea level rise, disease, heavy metals and other pollutants). Hence, future studies could usefully account for interactions between a larger suite of drivers and NC relationships.

The area of macroalgal mats was a significant predictor of mudflat area and saltmarsh area. For example when algal mats increased above $\sim -1(\text{SD})$, we noted significant decreasing trends in the area of both NC stocks. This is coherent with existing evidence that the smothering effect of excessive macroalgal growth and the concentrations of nitrates causing them are damaging to the

habitats of this internationally important site (Herbert *et al.*, 2010). As such, these results support recently proposed algal harvesting measures (Taylor, 2015) that have been suggested as a means to reduce and recycle nitrogen, as well as to reduce the volume of green macro-algae, thus protecting saltmarsh and mudflat habitats. While little information is available about the impacts that the macroalgal mats have on the businesses of the harbour, there are a number of studies in other estuaries (e.g. Troell *et al.*, 2005; Ferreira *et al.*, 2010) that indicate frequent macroalgal blooms can cause significant biodiversity loss, aesthetic impacts and public health problems, effectively eroding the benefit flows provided by NC stocks (as described in Figure 1, I).

As suggested in the STARS analysis above, areas of macroalgal mats and saltmarsh were shown to have significant negative but mostly linear effect (II, Figure 1) on wader and wildfowl numbers, with a threshold observed in both cases ~ -0.5 (SD). While mudflat area was not a significant predictor in our bird models, it did have a high relative importance in explaining the variation within models. Thus, as suggested by Bowgen *et al.* (2015) it is likely that waders/wildfowl in Poole Harbour are able to adapt to changes in their environment (e.g. increasing algal mats and reduced mudflat area) by switching to alternative habitats with different prey species and size classes, and may only undergo true tipping point transitions (i.e. VII, Figure 1) under extreme scenarios (e.g. the total removal of invertebrates from a system). However, this generalisation was developed based on analysis of the wader/wildfowl populations as a whole, and it is likely that individual species may have responded very differently to the environmental changes documented here (e.g. Durell *et al.*, 2006).

Two other environmental pressure variables, sediment shoaling and river flow, both responded to changes in mudflat and saltmarsh area in a deterministic manner. This is consistent with the fact that feedbacks between hydrodynamic forces and sediment accretion are key processes in shaping mudflats and saltmarshes (Kirwan & Murray, 2007; Wesenbeeck *et al.*, 2008). Here we show that sediment shoaling rates had a generally positive effect on mudflat area but mainly a negative impact on saltmarsh area. *Spartina* has been well documented as affecting the sediment regime of the harbour (Raybould, 2005), acting to consolidate sediment by rhizome growth in periods of expansion and releasing sediment into the harbour as it dies back, in a density dependent negative feedback manner. While many different biogeochemical mechanisms and drivers can lead to saltmarsh change (Crooks & Pye, 2000), there is evidence that the loss of *Spartina* in the harbour is mainly attributable to physical mechanisms such as direct human destruction (urbanisation) and erosion caused by changes in hydrodynamics and/or morphology (Gardiner, 2015). The optimal river flow rates predicted by the smoothing functions (Figure 5) suggest an abrupt threshold (III, Figure 1) for mudflat area ~ -0.5 (SD) and a negative linear effect on saltmarsh, with a shift in both variables towards net accretion trend at the current mean values for these assets at the harbour level. Accumulating evidence already suggests that many of the ecosystem services provided by saltmarshes have been jeopardized by the dieback of *Spartina* including the ability of the marshes to (1) reduce water flows and retain sediment (Raybould, 2005), (2) remediate nutrients and store heavy metals (Hübner *et al.*, 2010), (3) provide habitat for a variety of animals (Gardiner, 2015).

Finally, we identified fishing pressure to be the only significant driver to have influenced the abrupt time series trends in Manila clam stocks at two of the long term monitoring sites of the harbour. As expected, the relationship between fishing pressure and clam stocks was entirely linear (II, Figure 1), suggesting there was no definitive threshold where reducing fishing pressure could prevent the collapse of clam stocks. As fishing effort is controlled by the density of clams (the minimum landing size of Manila clams in Poole Harbour is 35 mm), this means that if the density of large sized clams increases so does fishing effort, and when the density decreases so does fishing effort (Humphreys *et al.*, 2007). This is analogous to a predator-prey system, whereby fishing effort increases after the population density increases, before reducing again once the population of "legal" sized clams has

reduced. However, Unregulated and Unsustainable (IUU) fishing has been noted as a particular problem for the fishery over the study period and before the introduction of the Permit Byelaw in 2015, there were significant illegal landings, the magnitude of which are unknown (Harris 2016). While IUU fishing activities almost certainly would have affected the value of landings being delivered into the harbour, the stock data used here (rather than landings) should highlight the densities of clams available in the harbour indiscriminately of legal or illegal fishing. In response to IUU fishing, enhanced enforcement by the local inshore fisheries and conservation authority (IFCA) has led to a significant reduction in illegal fishing and there are signs from this survey and some more recent stock assessments (SIFCA, 2017) that the new bylaws have had a positive impact on stocks of clams in the harbour, with a recently awarded Marine Stewardship Council (MSC) accreditation for Manila clams being designated as of 2018 (Williams & Davies 2018).

While fishing pressure is clearly a key driver in the population status of this species, it is also important to consider other mechanisms that could have been responsible for the abrupt shifts in the stocks seen in this study. For example, there is evidence that Manila clams cultured on the lease beds in the harbour were subject to recurring bouts of mass mortalities around 2006-2008 (Bateman *et al.*, 2012), resulting in many lease holders switching to other aquaculture species such as oysters (Othniel Oysters Ltd, Personal Communication, June 2018). From the literature it is unclear what caused such events but viral infection combined with low winter temperatures and food availability are the most likely possibilities (Humphreys *et al.*, 2007; Bateman *et al.*, 2012; Franklin *et al.*, 2012). Such occurrences provide an example of a potential positive feedback mechanism and possible evidence for a tipping point in the stocks of clams in the harbour (i.e. type VII, Figure 1). As viral infection reduces the fitness of the population (e.g. gamete release may be related to the metabolic depletion caused by the virus (Uddin *et al.*, 2010)), the carrying capacity of the population is also lowered owing to a decreased resistance to disease, causing a powerful positive feedback that further decreases shellfish stocks. This in turn has socio-economic consequences, with local aquaculture businesses and regulators potentially switching to more lucrative species as the condition of the NC stock is reduced. Therefore, while the environmental conditions of Poole Harbour are currently favourable for Manila clam proliferation (as evidenced by the recent increase in the wild stocks of the harbour), different types of disturbance may have acted together to cause an abrupt decline in the Manila clam aquaculture fisheries of the harbour and therefore the stocks of clams that was observed. In accordance with theory (Scheffer *et al.*, 2001), if a critical value of a press disturbance is exceeded, this may lead to a tipping point driven by a positive feedback mechanism, which could be triggered by a pulse disturbance. In this case study, fishing pressure (legal or illegal) and increasing water temperature can both be considered as press disturbances, the latter potentially increasing the risk of viral infections outbreaks, which represent a form of pulse disturbance. Such processes are not likely to be specific to Poole Harbour, with at least eleven estuaries in southern England currently accommodating naturalised populations of Manila clam (Humphreys *et al.*, 2015) and mass mortality events of Manila clam now being reported in other locations around the world (Pretto *et al.*, 2014; Nam *et al.*, 2018).

The increase in the wild stocks of a commercially attractive species such as Manila clam is also likely to have substantial consequences on the wider ecology and economy of the harbour. For instance, there is evidence that the introduction of the clams in the late 1980's has potentially had a positive effect on the over-winter mortality of several wader/wildfowl species such as oystercatchers in the Harbour (Caldow *et al.*, 2007). Thus, it could be suggested that if clam stocks were to continue to increase this would have the potential to provide an indirect benefit to several European shorebird populations *via* a spill-over effect increasing wild populations. There is also evidence that when

cultured at high densities Manila clams can provide other indirect benefits to humans such as altering biogeochemical cycles, thereby reducing the effects of nutrient pollution and the deployment of algal mats (Rose *et al.*, 2015), both of which are key issues for managers in Poole Harbour. Furthermore, in terms of direct economic value, a recent report by Williams & Davies (2018) suggests that although the overall landed weight of Manila clams and the value of landings have decreased by 50% and 25% respectively since 2010, the direct Gross Value Added (GVA) added to the local economy by the Manila clam is by a wide margin the highest of any species landed into the harbour (£838,911per annum vs the next highest species: whelks £249,562, based on 2016/2017 data). This suggests there is a local economic interest in ensuring that clam stocks remain high in the harbour. Nonetheless, such financial benefits must be balanced against the potential problems of removing commercial quantities of Manila clams from Poole Harbour. For example, there is evidence that the use of pump-scoop dredges can have significant impacts on the benthic community by reducing fine sediment and some prey species available to wintering birds (Clarke *et al.*, 2017). Managing fisheries and aquaculture development in a way that does not lead to deleterious ecosystem change is considered as a serious governance challenge not just in Poole Harbour but in many marine protected areas around the world (Edgar *et al.*, 2014). One way to avoid ecological tipping points as advocated by the FAO (The Food and Agriculture Organization of the United Nations), could be through prudent application of the precautionary principle (Carvalho *et al.*, 2006).

6 Conclusions

Given the growing evidence that coastal and shallow marine ecosystems are increasingly experiencing multiple disturbances, based on the numbers of studies reporting strong anthropogenic impacts resulting from multiple drivers (Crain *et al.*, 2008; Halpern *et al.*, 2008; Hewitt *et al.*, 2015; Gunderson *et al.*, 2016), both scientists and resource managers must confront the potential challenges of nonlinear shifts in ecosystem structure and function (Crain *et al.*, 2009; Côté *et al.*, 2016). Yet, despite the ecological literature being replete with terms related to ecological thresholds, tipping points and other concepts relating to multiple stable states (e.g. regime shifts), there is currently very little empirical evidence that such transitions actually occur in estuaries and other nearshore ecosystems (Nally *et al.*, 2014). Practical application of such concepts in a policy or management context are impeded by several factors such as 1) terminological inconsistency; 2) inadequacy of the temporal and spatial datasets for evaluating abrupt trends; 3) insufficient demonstration of mechanistic links between human or natural factors that cause ecosystem change (Capon *et al.*, 2015). In this study we have considered all three criteria and demonstrate that abrupt nonlinear thresholds in NC assets may occur in transitional protected systems such as harbours. The ecological thresholds that we have identified are driven by interactions among biophysical, ecological, and socioeconomic mechanisms mainly at the catchment scale. As we often lack robust ecological information in most systems to make *a priori* mechanistic predictions of where thresholds will occur (Dodds *et al.*, 2010), we believe that the methods outlined in this paper could be used to help local managers evaluate and articulate strategies to detect thresholds and tipping points in a way that can be incorporated in resource management frameworks (*sensu* Selkoe *et al.*, 2015). This would support global efforts by the United Nations Intergovernmental Oceanographic Commission (IOC) and other international initiatives to improve the long term sustainability of resources within large marine protected areas and their associated watersheds, with a particular focus on ecosystem based approaches to deliver healthy marine ecosystems and sustained ES. Further research could also usefully combine information on temporal trends with spatial data on status of NC and/or

multiple interacting drivers to create conceptual and dynamic modelling tools to support management decision-making.

Acknowledgements

This publication is part of a larger effort from the Mechanisms and Consequences of Tipping Points in Lowland Agricultural Landscapes (TPAL) project (www.tpalvaluing-nature.co.uk). The Valuing Nature Programme (www.valuing-nature.net) is funded by the Natural Environment Research Council, the Economic and Social Research Council, the Biotechnology and Biological Sciences Research Council, the Arts and Humanities Research Council and the Department for Environment, Food and Rural Affairs. This work was supported by NERC grant reference number: NE/P007716/1. We would like to thank the four anonymous reviewers for their helpful and constructive comments that greatly contributed to improving the final version of the paper.

References

- Anders, N.R., Churchyard, T. and Hiddink, J.G., (2009). Predation of the shelduck *Tadorna tadorna* on the mud snail *Hydrobia ulvae*. *Aquatic ecology*, 43(4), pp.1193.
- Andersen, T., Carstensen, J., Hernandez-Garcia, E. and Duarte, C.M., (2009). Ecological thresholds and regime shifts: approaches to identification. *Trends in Ecology and Evolution*, 24(1), pp.49-57.
- Ashwin, P., Wieczorek, S., Vitolo, R. and Cox, P., (2012). Tipping points in open systems: bifurcation, noise-induced and rate-dependent examples in the climate system. *Philosophical Transactions of the Royal Society*, 370(1962), pp.1166-1184.
- Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C. and Silliman, B.R., (2011). The value of estuarine and coastal ecosystem services. *Ecological monographs*, 81(2), pp.169-193.
- Bateman, K. S., White, P., and Longshaw, M. (2012). Virus-like particles associated with mortalities of the Manila clam *Ruditapes philippinarum* in England. *Diseases of aquatic organisms*, 99(2), pp.163-167.
- Beaumont, N.J., Austen, M.C., Mangi, S.C. and Townsend, M., (2008). Economic valuation for the conservation of marine biodiversity. *Marine pollution bulletin*, 56(3), pp.386-396.
- Bestelmeyer, B. T., Ellison, A. M., Fraser, W. R., Gorman, K. B., Holbrook, S. J., Laney, C. M., Sharma, S. (2011). Analysis of abrupt transitions in ecological systems. *Ecosphere*, 2(12), pp.129.
- Biggs, R., Schlüter, M., Biggs, D., Bohensky, E.L., BurnSilver, S., Cundill, G., Dakos, V., Daw, T.M., Evans, L.S., Kotschy, K. and Leitch, A.M., (2012). Toward principles for enhancing the resilience of ecosystem services. *Annual review of environment and resources*, 37, pp.421-448.
- Bowes, M.J., Smith, J.T., Neal, C., Leach, D.V., Scarlett, P.M., Wickham, H.D., Harman, S.A., Armstrong, L.K., Davy-Bowker, J., Haft, M. and Davies, C.E., (2011). Changes in water quality of the River Frome (UK) from 1965 to 2009: Is phosphorus mitigation finally working?. *Science of the Total Environment*, 409(18), pp.3418-3430.
- Bowgen, K.M., Stillman, R.A., and Herbert, R.J.H, (2015). Predicting the effect of invertebrate regime shifts on wading birds: Insights from Poole Harbour, UK. *Biological Conservation*, 186, pp.60-68.

- 651 Bryan, G., Kite, D., Money, R., Jonas, P. and Barden, R. (2013). Strategy for Managing Nitrogen in
652 the Poole Harbour catchment to 2035. Environment Agency; Natural England.
- 653 Caldwell, R.W., Stillman, R.A., dit Durell, S.E.L.V., West, A.D., McGrorty, S., Goss-Custard, J.D., Wood,
654 P.J. and Humphreys, J., (2007). Benefits to shorebirds from invasion of a non-native
655 shellfish. *Proceedings of the Royal Society of London B: Biological Sciences*, 274(1616), pp.1449-1455.
- 656 Cantarello, E., Newton, A.C., Martin, P.A., Evans, P.M., Gosal, A. and Lucash, M.S., (2017).
657 Quantifying resilience of multiple ecosystem services and biodiversity in a temperate forest
658 landscape. *Ecology and evolution*, 7(22), pp.9661-9675.
- 659 Capon, S.J., Lynch, A.J.J., Bond, N., Chessman, B.C., Davis, J., Davidson, N., Finlayson, M., Gell, P.A.,
660 Hohnberg, D., Humphrey, C. and Kingsford, R.T., (2015). Regime shifts, thresholds and multiple
661 stable states in freshwater ecosystems; a critical appraisal of the evidence. *Science of the Total*
662 *Environment*, 534, pp.122-130.
- 663 Carpenter, S.R., Cole, J.J., Pace, M.L., Batt, R., Brock, W.A., Cline, T., Coloso, J., Hodgson, J.R., Kitchell,
664 J.F., Seekell, D.A. and Smith, L., (2011). Early warnings of regime shifts: a whole-ecosystem
665 experiment. *Science*, 332(6033), pp.1079-1082.
- 666 Carvalho, F.P., (2006). Agriculture, pesticides, food security and food safety. *Environmental science &*
667 *policy*, 9(7-8), pp.685-692.
- 668 Chevillot, X., Pierre, M., Rigaud, A., Drouineau, H., Chaalali, A., Sautour, B., and Lobry, J. (2016).
669 Abrupt shifts in the Gironde fish community: an indicator of ecological changes in an estuarine
670 ecosystem. *Marine Ecology Progress Series*, 549, pp.137-151.
- 671 Clarke, L.J., Esteves, L.S., Stillman, R.A. and Herbert, R.J.H, (2017). Impacts of a novel shellfishing gear
672 on macrobenthos in a marine protected area: pump-scoop dredging in Poole Harbour, UK. *Aquatic*
673 *Living Resources*, 31, pp.5.
- 674 Conley, D.J., Paerl, H.W., Howarth, R.W., Boesch, D.F., Seitzinger, S.P., Havens, K.E., Lancelot, C. and
675 Likens, G.E., (2009). Controlling eutrophication: nitrogen and phosphorus. *Science*, 323(5917),
676 pp.1014-1015.
- 677 Conversi, A., Umani, S. F., Peluso, T., Molinero, J. C., Santojanni, A., and Edwards, M. (2010). The
678 Mediterranean Sea regime shift at the end of the 1980s, and intriguing parallelisms with other
679 European basins. *Plos one*, 5(5), e10633.
- 680 Côté, I.M., Darling, E.S. and Brown, C.J., (2016). Interactions among ecosystem stressors and their
681 importance in conservation. *Proceedings of the Royal Society B*, 283(1824), pp.20152592.
- 682 Crain, C.M., Halpern, B.S., Beck, M.W. and Kappel, C.V., (2009). Understanding and managing human
683 threats to the coastal marine environment. *Annals of the New York Academy of Sciences*, 1162(1),
684 pp.39-62.
- 685 Crain, C.M., Kroeker, K. and Halpern, B.S., (2008). Interactive and cumulative effects of multiple
686 human stressors in marine systems. *Ecology letters*, 11(12), pp.1304-1315.
- 687 Crooks, S., and Pye, K. (2000). Sedimentological controls on the erosion and morphology of
688 saltmarshes: implications for flood defence and habitat recreation. *Geological Society*, London,
689 Special Publications, 175(1), pp.207-222.

- deYoung, B., Barange, M., Beaugrand, G., Harris, R., Perry, R. I., Scheffer, M., and Werner, F. (2008). Regime shifts in marine ecosystems: detection, prediction and management. *Trends in Ecology and Evolution*, 23(7), pp.402–409.
- Dodds, W. K., Clements, W. H., Gido, K., Hilderbrand, R. H., and King, R. S. (2010). Thresholds, breakpoints, and non-linearity in freshwaters as related to management. *Journal of the North American Benthological Society*, 29(3), pp.988-997.
- Donohue, I., Hillebrand, H., Montoya, J.M., Petchey, O.L., Pimm, S.L., Fowler, M.S., Healy, K., Jackson, A.L., Lurgi, M., McClean, D. and O'Connor, N.E., (2016). Navigating the complexity of ecological stability. *Ecology letters*, 19(9), pp.1172-1185.
- Durell, S.E.L.V., Stillman, R.A., Caldow, R.W., McGrorty, S., West, A.D. and Humphreys, J., (2006). Modelling the effect of environmental change on shorebirds: a case study on Poole Harbour, UK. *Biological Conservation*, 131(3), pp.459-473.
- Dyrynda, P., (2005). 8. Sub-tidal ecology of Poole Harbour—An overview. In *Proceedings in marine science* (7), pp.109-130.
- Edgar, G.J., Stuart-Smith, R.D., Willis, T.J., Kininmonth, S., Baker, S.C., Banks, S., Barrett, N.S., Becerro, M.A., Bernard, A.T., Berkhout, J. and Buxton, C.D., (2014). Global conservation outcomes depend on marine protected areas with five key features. *Nature*, 506(7487), p.216.
- Elliott, M. (2011). Marine science and management means tackling exogenic unmanaged pressures and endogenic managed pressures—a numbered guide. *Marine Pollution Bulletin*, 62, 651e655.
- Ferreira, J.G., Andersen, J.H., Borja, A., Bricker, S.B., Camp, J., Cardoso da Silva, M., Garcés, E., Heiskanen, A.S., Humborg, C., Ignatiades, L. and Lancelot, C., (2010). Marine strategy framework directive—task group 5 report eutrophication. *EUR*, 24338, pp.49.
- Foley, M.M., Martone, R.G., Fox, M.D., Kappel, C.V., Mease, L.A., Erickson, A.L., Halpern, B.S., Selkoe, K.A., Taylor, P. and Scarborough, C., (2015). Using ecological thresholds to inform resource management: current options and future possibilities. *Frontiers in Marine Science*, 2, pp.95.
- Folke, C., Jansson, Å., Rockström, J., Olsson, P., Carpenter, S.R., Chapin III, F.S., Crépin, A.S., Daily, G., Danell, K., Ebbesson, J. and Elmqvist, T., (2011). Reconnecting to the biosphere. *AMBIO: A Journal of the Human Environment*, 40(7), pp.719-738.
- Franklin, D.J., Humphreys, J., Harris, M., Jensen, A.C., Herbert, R.J.H. and Purdie, D.A. (2012). An investigation into the annual cycle of phytoplankton abundance in Poole Harbour and its relationship with Manila clam nutrition. Report for the Marine Management Organisation.
- Gardiner, S. (2015). Physical drivers of saltmarsh change in enclosed microtidal estuaries (Doctoral dissertation, University of Southampton).
- Gardiner, S., S. Hanson, R. J. Nicholls, Z. Zhang, S. Jude, A. Jones, J. Richards, A. Williams, T. Spencer, S. Cope (2007). *The Habitats Directive, Coastal Habitats and Climate Change - Case Studies from the South Coast of the UK* In Tyndall Centre Working Paper pp.108.
- Gladwell, M. (2000). *The Tipping Point*. Abacus, London.

- 727 Goss-Custard, J. D., Stillman, R. A., West, A. D., Caldow, R. W. G., Triplet, P., Dit Durell, S. L. V., and
 728 McGrorty, S. (2004). When enough is not enough: shorebirds and shellfishing. *Proceedings of the*
 729 *Royal Society of London B: Biological Sciences*, 271(1536), pp.233-237.
- 730 Goss-Custard, J.D. and Moser, M.E., (1988). Rates of change in the numbers of dunlin, *Calidris alpina*,
 731 wintering in British estuaries in relation to the spread of *Spartina anglica*. *Journal of Applied Ecology*,
 732 pp.95-109.
- 733 Groffman, P.M., Baron, J.S., Blett, T., Gold, A.J., Goodman, I., Gunderson, L.H., Levinson, B.M.,
 734 Palmer, M.A., Paerl, H.W., Peterson, G.D. and Poff, N.L., (2006). Ecological thresholds: the key to
 735 successful environmental management or an important concept with no practical
 736 application?. *Ecosystems*, 9(1), pp.1-13.
- 737 Grömping, U. (2006). Relative importance for linear regression in R: the package relaimpo. *Journal of*
 738 *statistical software*, 17 (1), pp.1-27.
- 739 Gunderson, A.R., Armstrong, E.J. and Stillman, J.H., (2016). Multiple stressors in a changing world:
 740 the need for an improved perspective on physiological responses to the dynamic marine
 741 environment. *Annual Review of Marine Science*, 8, pp.357-378.
- 742 Halpern, B. S., McLeod, K. L., Rosenberg, A. A., and Crowder, L. B. (2008). Managing for cumulative
 743 impacts in ecosystem-based management through ocean zoning. *Ocean and Coastal Management*,
 744 51(3),pp.203–211.
- 745 Harris, M. R. (2016). A study of the naturalisation and dispersal of a non-native bivalve, the Manila
 746 clam, *Ruditapes philippinarum* (Adams and Reeve 1850) in estuaries along the South coast of
 747 England (Doctoral dissertation, University of Portsmouth).
- 748 Herbert, R.J., Grandfield F., Redford M., Rivers E., Hall A. and Harrison A. (2018) Littoral sediment
 749 condition of Poole Harbour SSSI. BU Global Environmental Solutions (BUG) report (BUG2777) to
 750 Natural England.pp 34.
- 751 Herbert, R.J., Ross, K., Huebner, R. and Stillman, R.A., (2010). Intertidal invertebrates and biotopes of
 752 Poole Harbour SSSI and survey of Brownsea Island Lagoon.
- 753 Herbert, R.J., Willis, J., Jones, E., Ross, K., Hübner, R., Humphreys, J., Jensen, A. and Baugh, J., (2012).
 754 Invasion in tidal zones on complex coastlines: modelling larvae of the non-native Manila clam,
 755 *Ruditapes philippinarum*, in the UK. *Journal of Biogeography*, 39(3), pp.585-599.
- 756 Hewitt, J. E., and Thrush, S. F. (2010). Empirical evidence of an approaching alternate state produced
 757 by intrinsic community dynamics, climatic variability and management actions. *Marine Ecology*
 758 *Progress Series*, 413, pp267–276.
- 759 Hewitt, J. E., Ellis, J.I. and Thrush, S.F., (2015). Multiple stressors, nonlinear effects and the
 760 implications of climate change impacts on marine coastal ecosystems. *Global Change Biology*, 22(8),
 761 pp.2665-2675.
- 762 Howarth, R.W. and Marino, R., (2006). Nitrogen as the limiting nutrient for eutrophication in coastal
 763 marine ecosystems: evolving views over three decades. *Limnology and Oceanography*, 51(part2),
 764 pp.364-376.

- Hübner, R., Herbert, R.J.H, and Astin, K.B., (2010). Cadmium release caused by the die-back of the saltmarsh cord grass *Spartina anglica* in Poole Harbour (UK). *Estuarine, Coastal and Shelf Science*, 87(4), pp.553-560.
- Huggett, A.J., (2005). The concept and utility of 'ecological thresholds' in biodiversity conservation. *Biological conservation*, 124(3), pp.301-310.
- Hughes, T.P., Linares, C., Dakos, V., van de Leemput, I.A. and van Nes, E.H., (2013). Living dangerously on borrowed time during slow, unrecognized regime shifts. *Trends in Ecology & Evolution*, 28(3), pp.149-155.
- Humphreys, J. and May, V., (2005). Introduction: Poole Harbour in context. *Proceedings in Marine Science*, 7, pp.1-7.
- Humphreys, J., (2005). Salinity and tides in Poole Harbour: Estuary or lagoon?. *Proceedings in Marine Science*, 7, pp.35-47.
- Humphreys, J., Caldow, R.W., McGrorty, S., West, A.D. and Jensen, A.C., (2007). Population dynamics of naturalised Manila clams *Ruditapes philippinarum* in British coastal waters. *Marine Biology*, 151(6), pp.2255.
- Humphreys, J., Harris, M.R., Herbert, R.J.H, Farrell, P., Jensen, A. and Cragg, S.M., (2015). Introduction, dispersal and naturalization of the Manila clam *Ruditapes philippinarum* in British estuaries, 1980–2010. *Journal of the Marine Biological Association of the United Kingdom*, 95(6), pp.1163-1172.
- Jasim Uddin, M., Yang, H.S., Choi, K.S., Kim, H.J., Hong, J.S. and Cho, M., (2010). Seasonal changes in *Perkinsus olseni* infection and gametogenesis in Manila clam, *Ruditapes philippinarum*, from Seonjaedo Island in Incheon, off the west coast of Korea. *Journal of the world aquaculture society*, 41(s1), pp.93-101.
- Jensen, A. C., Humphreys, J., Caldow, R. W. G., Grisley, C., and Dyrinda, P. E. J. (2004). Naturalization of the Manila clam (*Tapes philippinarum*), an alien species, and establishment of a clam fishery within Poole Harbour, Dorset. *Journal of the Marine Biological Association of the United Kingdom*, 84(5), pp.1069-1073.
- JNCC, (2004). Common Standards Monitoring Guidance for Littoral Sediment Habitats. Peterborough: Joint Nature Conservation Committee.
- Johnson, C.J., (2013). Identifying ecological thresholds for regulating human activity: Effective conservation or wishful thinking?. *Biological Conservation*, 168, pp.57-65.
- Jones, M. and Pinn, E., (2006). The impact of a macroalgal mat on benthic biodiversity in Poole Harbour. *Marine pollution bulletin*, 53(1-4), pp.63-71.
- Kirwan, M. L., and Murray, A. B. (2007). A coupled geomorphic and ecological model of tidal marsh evolution. *Proceedings of the National Academy of Sciences*, 104(15), pp.6118-6122.
- Kite, D.J., Bryan, G., Jonas, P. (2012). Strategy for Managing Nitrogen in Poole Harbour to 2035 (final version 6 June 2013); Environment Agency Report.
- Large, S. I., Fay, G., Friedland, K. D., and Link, J. S. (2013). Defining trends and thresholds in responses of ecological indicators to fishing and environmental pressures. *ICES Journal of Marine Science*, 70(4), pp.755-767.

- Large, S.I., Fay, G., Friedland, K.D. and Link, J.S., (2015). Critical points in ecosystem responses to fishing and environmental pressures. *Marine Ecology Progress Series*, 521, pp.1-17.
- Lenton, T. M., Held, H., Kriegler, E., Hall, J. W., Lucht, W., Rahmstorf, S., and Schellnhuber, H. J. (2008). Tipping elements in the Earth's climate system. *Proceedings of the National Academy of Sciences*, 105(6), pp.1786–1793.
- Lenton, T.M., (2013). Environmental tipping points. *Annual Review of Environment and Resources*, 38, pp.1-29.
- Levin, P.S. and Möllmann, C., (2015). Marine ecosystem regime shifts: challenges and opportunities for ecosystem-based management. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 370(1659), pp.20130275.
- Lyons, D.A., Arvanitidis, C., Blight, A.J., Chatzinikolaou, E., Guy-Haim, T., Kotta, J., Orav-Kotta, H., Queirós, A.M., Rilov, G., Somerfield, P.J. and Crowe, T.P., (2014). Macroalgal blooms alter community structure and primary productivity in marine ecosystems. *Global change biology*, 20(9), pp.2712-2724.
- Mac Nally, R., Albano, C. and Fleishman, E., (2014). A scrutiny of the evidence for pressure-induced state shifts in estuarine and nearshore ecosystems. *Austral Ecology*, 39(8), pp.898-906.
- Mace, G. M., Hails, R. S., Cryle, P., Harlow, J., and Clarke, S. J. (2015). Towards a risk register for natural capital. *Journal of Applied Ecology*, 52(3), pp.641–653.
- Martin, J., Runge, M.C., Nichols, J.D., Lubow, B.C. and Kendall, W.L., (2009). Structured decision making as a conceptual framework to identify thresholds for conservation and management. *Ecological Applications*, 19(5), pp.1079-1090.
- May, V., (2005). Geomorphology of Poole Harbour. In *Proceedings in Marine Science* (7), pp.25-34. Elsevier.
- McGlathery, K.J., Sundbäck, K. and Anderson, I.C., (2007). Eutrophication in shallow coastal bays and lagoons: the role of plants in the coastal filter. *Marine Ecology Progress Series*, 348, pp.1-18.
- McGrorty, S., and Goss-Custard, J. D. (1987). A review of the rehabilitation of areas cleared of *Spartina*. *Institute of Terrestrial Ecology (Natural Environment Research Council)*, Dorset, UK.
- McLusky D.S, Elliott M., Transitional Waters: a new approach, semantics or just muddying the waters? (2007) *Estuary Coast Shelf Science*, 71, pp.359-363.
- Meyer, K. (2016). A mathematical review of resilience in ecology. *Natural Resource Modeling*, 29(3), pp.339-352.
- Mollmann, C., Diekmann, R., Müller-Karulis, B., Kornilovs, G., Plikshs, M. and Axe, P. (2009). Reorganization of a large marine ecosystem due to atmospheric and anthropogenic pressure: a discontinuous regime shift in the Central Baltic Sea. *Global Change Biology*, 15, pp.1377–1393
- Nam, K.W., Jeung, H.D., Song, J.H., Park, K.H., Choi, K.S. and Park, K.I., (2018). High parasite burden increases the surfacing and mortality of the Manila clam (*Ruditapes philippinarum*) in intertidal sandy mudflats on the west coast of Korea during hot summer. *Parasites & vectors*, 11(1), p.42.
- Natural Capital Committee. (2014). The state of natural capital: restoring our natural assets. Second Report to the Economic Affairs Committee. Natural Capital Committee, HM Government UK.


- 844 Newton, A. C. (2016). Biodiversity Risks of Adopting Resilience as a Policy Goal. *Conservation Letters*,
845 9(5), pp.369–376.
- 846 Office of National Statistics (2010) Revised Annual Mid-year Population Estimates: 2001 to
847 2010: www.ons.gov.uk. [Accessed 22/10/2017]
- 848 Pachauri, R.K., Allen, M.R., Barros, V.R., Broome, J., Cramer, W., Christ, R., Church, J.A., Clarke, L.,
849 Dahe, Q., Dasgupta, P. and Dubash, N.K., (2014). Climate change 2014: synthesis report. Contribution
850 of Working Groups I, II and III to the fifth assessment report of the Intergovernmental Panel on
851 Climate Change, *IPCC*, pp.151.
- 852 Petraitis, P. (2013). Multiple stable states in natural ecosystems. *OUP*, Oxford.
- 853 Piet, G. J., Quirijns, F. J., Robinson, L., and Greenstreet, S. P. R. (2006). Potential pressure indicators
854 for fishing, and their data requirements. *ICES Journal of Marine Science*, 64(1), pp.110-121.
- 855 Pretto, T., Zambon, M., Civettini, M., Caburlotto, G., Boffo, L., Rossetti, E. and Arcangeli, G., (2014).
856 Massive mortality in Manila clams (*Ruditapes philippinarum*) farmed in the Lagoon of Venice, caused
857 by *Perkinsus olseni*. *Bulletin of the European Association of Fish Pathologists*, 34(2), pp.43-53.
- 858 Quinlan, A.E., Barbés-Blázquez, M., Haider, L.J. and Peterson, G.D., (2016). Measuring and assessing
859 resilience: broadening understanding through multiple disciplinary perspectives. *Journal of Applied*
860 *Ecology*, 53(3), pp.677-687.
- 861 R Core Team. R: A language and environment for statistical computing. Version [3.3.5"]. Vienna: R
862 Foundation for Statistical Computing. Available from: URL <https://www.R-project.org/>.
- 863 Raffaelli, D., (1999). Nutrient enrichment and trophic organisation in an estuarine food web. *Acta*
864 *Oecologica*, 20(4), pp.449-461.
- 865 Raffaelli, D., Limia, J., Hull, S. and Pont, S., (1991). Interactions between the amphipod *Corophium*
866 *volutator* and macroalgal mats on estuarine mudflats. *Journal of the Marine Biological Association of*
867 *the United Kingdom*, 71(4), pp.899-908.
- 868 Raybould, A. (2005). History and ecology of *Spartina anglica* in Poole Harbour; In Humphreys, J. and
869 May, V. eds. *The Ecology of Poole Harbour*. Oxford: Elsevier, pp.71-90.
- 870 Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F.S., Lambin, E.F., Lenton, T.M., Scheffer,
871 M., Folke, C., Schellnhuber, H.J. and Nykvist, B., (2009). A safe operating space for
872 humanity. *Nature*, 461(7263), pp.472-475.
- 873 Rodionov, S., and Overland, J. E. (2005). Application of a sequential regime shift detection method to
874 the Bering Sea ecosystem. *ICES Journal of Marine Science*, 62(3), pp. 328-332.
- 875 Rodionov, S.N., (2004). A sequential algorithm for testing climate regime shifts. *Geophysical*
876 *Research Letters*, 31(9).
- 877 Rose, J.M., Bricker, S.B. and Ferreira, J.G., (2015). Comparative analysis of modeled nitrogen removal
878 by shellfish farms. *Marine pollution bulletin*, 91(1), pp.185-190.
- 879 Samhouri, J.F., Andrews, K.S., Fay, G., Harvey, C.J., Hazen, E.L., Hennessey, S.M., Holsman, K.,
880 Hunsicker, M.E., Large, S.I., Marshall, K.N. and Stier, A.C., (2017). Defining ecosystem thresholds for
881 human activities and environmental pressures in the California Current. *Ecosphere*, 8(6).

- 882 Samhuri, J.F., Levin, P.S. and Ainsworth, C.H., (2010). Identifying thresholds for ecosystem-based
883 management. *PLoS One*, 5(1), pp.e8907.
- 884 Scheffer, M., Bascompte, J., Brock, W.A., Brovkin, V., Carpenter, S.R., Dakos, V., Held, H., Van Nes,
885 E.H., Rietkerk, M. and Sugihara, G., (2009). Early-warning signals for critical
886 transitions. *Nature*, 461(7260), pp.53-59.
- 887 Scheffer, M., Carpenter, S., Foley, J.A., Folke, C. and Walker, B., (2001). Catastrophic shifts in
888 ecosystems. *Nature*, 413(6856), pp.591-596.
- 889 Scheffer, M., Carpenter, S.R., Lenton, T.M., Bascompte, J., Brock, W., Dakos, V., Van de Koppel, J.,
890 Van de Leemput, I.A., Levin, S.A., Van Nes, E.H. and Pascual, M., (2012). Anticipating critical
891 transitions. *Science*, 338(6105), pp.344-348.
- 892 Selkoe, K.A., Blenckner, T., Caldwell, M.R., Crowder, L.B., Erickson, A.L., Essington, T.E., Estes, J.A.,
893 Fujita, R.M., Halpern, B.S., Hunsicker, M.E. and Kappel, C.V., (2015). Principles for managing marine
894 ecosystems prone to tipping points. *Ecosystem Health and Sustainability*, 1(5), pp.1-18.
- 895 Selkoe, K.A., Blenckner, T., Caldwell, M.R., Crowder, L.B., Erickson, A.L., Essington, T.E., Estes, J.A.,
896 Fujita, R.M., Halpern, B.S., Hunsicker, M.E. and Kappel, C.V., (2015). Principles for managing marine
897 ecosystems prone to tipping points. *Ecosystem Health and Sustainability*, 1(5), pp.1-18.
- 898 SIFCA, (2017). Poole Harbour Bivalve Stock Survey 2017. Southern Inshore Fisheries and
899 Conservation Authority. Available: [http://www.southern-ifca.gov.uk/research-and-evidence-](http://www.southern-ifca.gov.uk/research-and-evidence-reports)
900 [reports](http://www.southern-ifca.gov.uk/research-and-evidence-reports)[Accessed 29/08/2018]
- 901 Simpson., M (2004). Poole Harbour Approach Channel Deepening and Beneficial Use of Dredged
902 Material EIA. Non-Technical Summary. Overview of the existing environment. Fishing activity.
903 Published report, Royal Haskoning, Exeter, Devon. pp.275-292.
904 https://www.phc.co.uk/downloads/channeldeepening/es10_fishing.pdf [Accessed 02/04/2018]
- 905 Taylor, D., (2015). Review of potential to remove harmful algae and reduce nitrogen load in Poole
906 Harbour. Commissioned by Dorset Local Nature Partnership. Available:
907 [www.wessexwater.co.uk/About-us/Environment/Catchment-partnerships/Poole-harbour/Algal-](http://www.wessexwater.co.uk/About-us/Environment/Catchment-partnerships/Poole-harbour/Algal-Harvesting-workshop)
908 [Harvesting-workshop](http://www.wessexwater.co.uk/About-us/Environment/Catchment-partnerships/Poole-harbour/Algal-Harvesting-workshop) [Accessed 02/04/2018]
- 909 Thornton, A. (2016). The impact of green macroalgal mats on benthic invertebrates and
910 overwintering wading birds (Doctoral dissertation, Bournemouth University).
- 911 Toms, J. and Villard, M.A., (2015). Threshold detection: matching statistical methodology to
912 ecological questions and conservation planning objectives. *Avian Conservation and Ecology*, 10(1).
- 913 Troell, M., Pihl, L., Rönnbäck, P., Wennhage, H., Söderqvist, T., and Kautsky, N. (2005). Regime shifts
914 and ecosystem services in Swedish coastal soft bottom habitats: when resilience is
915 undesirable. *Ecology and Society*, 10(1).
- 916 Tubbs, C. R. and Tubbs, J. M., (1980). Wader and Shelduck feeding distribution in Langstone Harbour,
917 Hampshire. *Bird Study*, 27 (4), 239-248
- 918 Tubbs, C.R., Tubbs, J.M. and Kirby, J.S., (1992). Dunlin *Calidris alpina alpina* in the Solent, southern
919 England. *Biological Conservation*, 60(1), pp.15-24.
- 920 Underhill-Day, J. C. (2006). A condition assessment of Poole Harbour European Marine Site.
921 Unpublished report, Footprint Ecology/Natural England. Dorset. England.

- 922 van Nes, E.H., Arani, B.M., Staal, A., van der Bolt, B., Flores, B.M., Bathiany, S. and Scheffer, M.,
 923 (2016). What Do You Mean, 'Tipping Point'? *Trends in Ecology & Evolution*, 31(12), pp.902-904.
- 924 Vinod, H.D. and López-de-Lacalle, J., (2009). Maximum entropy bootstrap for time series: the
 925 meboot R package. *Journal of Statistical Software*, 29(5), pp.1-19.
- 926 Wesenbeeck, B.K., van de Koppel, J., Herman, P.M., Bertness, M.D., van der Wal, D., Bakker, J.P. and
 927 Bouma, T.J., (2008). Potential for sudden shifts in transient systems: distinguishing between local
 928 and landscape-scale processes. *Ecosystems*, 11(7), pp.1133-1141.
- 929 Williams, C and Davies, W., (2018). A Tale of Three Fisheries The value of the small-scale commercial
 930 fishing fleet , aquaculture and the recreational charter boat fleet , to the local economy of Poole.
 931 NEF Consulting Limited. A Report to Southern IFCA. Available: [https://www.nefconsulting.com/a-](https://www.nefconsulting.com/a-tale-of-three-fisheries)
 932 [tale-of-three-fisheries](https://www.nefconsulting.com/a-tale-of-three-fisheries) [Accessed 29/08/2018]
- 933 Wood, S.N. and Augustin, N.H., (2002). GAMs with integrated model selection using penalized
 934 regression splines and applications to environmental modelling. *Ecological Modelling*, 157(2),
 935 pp.157-177.
- 936 Wood, S.N., (2004). mgcv: GAMs with GCV smoothness estimation and GAMMs by REML/PQL. R
 937 package version 1.1-8. *R Foundation for Statistical Computing*, Vienna, Austria.
- 938 Wood, S.N., (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of
 939 semiparametric generalized linear models. *Journal of the Royal Statistical Society: Series B (Statistical*
 940 *Methodology)*, 73(1), pp.3-36.

941

Appendix 1: Normalised time series of environmental drivers, in Poole Harbour for the period 1980-2015. From top to bottom: fishing pressure (no. boats), macroalgal mats (area (ha), nitrates ($\text{mg NO}_3 \text{ N l}^{-1}$), phosphates ($\mu\text{g l}^{-1}$), riparian water flows ($\text{m}^3 \text{ s}^{-1}$), sediment shoaling (m) and water temperature ($^{\circ}\text{C}$). Vertical black lines represent statistically significant ($p \leq 0.01$) breakpoints for individual trends from sequential Student's t-test.



Highlights: 3-5 bullet points, each max. 85 characters

- Addressing tipping points leads to improved management outcomes in MPAs.
- We identified nonlinear thresholds in several of the natural capital assets of Poole harbour.
- Abrupt nonlinear trends were the most common threshold identified from our time series analysis.
- Tipping points were detected most strongly in the Manila clam fisheries of the harbour
- Restoration targets need to consider multiple drivers not just recognisable drivers