

12-2019

## Caesarean Section rates in South Asian cities: Can midwifery help stem the rise?

Sulochana Dhakal Rai

*Bournemouth University, UK, sdhakalrai@bournemouth.ac.uk*

Amudha Poobalan

*University of Aberdeen, UK, a.poobalan@abdn.ac.uk*

Rafat Jan

*Aga Khan University, rafat.jan@aku.edu*

Malin Bogren

*University of Gothenburg, Sweden, malin.bogren@gu.se*

Juliet Wood

*Bournemouth University, UK, jwood@bournemouth.ac.uk*

*See next page for additional authors*

Follow this and additional works at: <https://ecommons.aku.edu/jam>



Part of the [Nursing Midwifery Commons](#)

### Recommended Citation

Rai, S D, Poobalan, A, Jan, R, Bogren, M, Wood, J, Dangal, G, Regmi, P, Teijlingen, E, Dhakal, K B, Badar, S J, & Shahid, F. Caesarean Section rates in South Asian cities: Can midwifery help stem the rise?. *Journal of Asian Midwives*. 2019;6(2):4–22.

---

## Caesarean Section rates in South Asian cities: Can midwifery help stem the rise?

### Authors

Sulochana Dhakal Rai, Amudha Poobalan, Rafat Jan, Malin Bogren, Juliet Wood, Ganesh Dangal, Pramod Regmi, Edwin van Teijlingen, Keshar Bahadur Dhakal, Sahib Jan Badar, and Farhana Shahid

## Caesarean Section rates in South Asian cities: Can midwifery help stem the rise?

<sup>1</sup>Sulochana Dhakal Rai, <sup>2</sup>Amudha Poobalan, <sup>3</sup>Rafat Jan, <sup>4</sup>Malin Bogren, <sup>5</sup>Juliet Wood, <sup>6</sup>Ganesh Dangal, <sup>7</sup>Pramod Regmi, <sup>8\*</sup>Edwin van Teijlingen, <sup>9</sup>Keshar Bahadur Dhakal, <sup>10</sup>Sahib Jan Badar, <sup>11</sup>Farhana Shahid

1. RGN, PhD Student, Centre for Midwifery, Maternal & Perinatal Health, Faculty of Health & Social Sciences, Bournemouth University, UK. Email: [sdhakalrai@bournemouth.ac.uk](mailto:sdhakalrai@bournemouth.ac.uk)
2. Senior Lecturer in Public Health, Institute of Applied Health Sciences, School of Medicine, Medical Sciences & Nutrition, University of Aberdeen, UK. Email: [a.poobalan@abdn.ac.uk](mailto:a.poobalan@abdn.ac.uk)
3. Professor, Aga Khan University, Email: [rafat.jan@aku.edu](mailto:rafat.jan@aku.edu)
4. RN, RM, PhD, Institute of Health Care Sciences, Sahlgrenska Academy, University of Gothenburg, Sweden. Email: [malin.bogren@gu.se](mailto:malin.bogren@gu.se)
5. Lecturer in Midwifery Centre for Midwifery, Maternal & Perinatal Health, Faculty of Health & Social Sciences, Bournemouth University, UK. Email: [jwood@bournemouth.ac.uk](mailto:jwood@bournemouth.ac.uk)
6. Professor, National Academy of Medical Sciences (NAMS): Senior Consultant in Department of Gynecology & Obstetrics, Kathmandu Model Hospital, Kathmandu, Nepal. Email: [ganesh.dangal@gmail.com](mailto:ganesh.dangal@gmail.com)
7. Lecturer in International Health, Faculty of Health & Social Sciences, Bournemouth University, UK. Email: [pregmi@bournemouth.ac.uk](mailto:pregmi@bournemouth.ac.uk); Visiting Fellow: Datta Meghe Institute of Medical Sciences, India
8. Professor of Reproductive Health Research, Centre for Midwifery, Maternal & Perinatal Health, Bournemouth University, UK. Email: [evteijlingen@bournemouth.ac.uk](mailto:evteijlingen@bournemouth.ac.uk); Visiting Professor, Manmohan Memorial Institute of Health Sciences, Kathmandu, Nepal.
9. Chief Consultant, Department of Gynecology & Obstetrics, Karnali Province Hospital, Nepal; Professor in Karnali Academy of Health Science, Nepal. Email: [drkeshar\\_dhakal@yahoo.com](mailto:drkeshar_dhakal@yahoo.com)
10. Sahib Jan Badar, Program Coordinator, Accelerated Action Plan (AAP) on Reduce Stunting and Malnutrition in mothers and children under the 5 years, Pak. Email: [sjb\\_dr@yahoo.com](mailto:sjb_dr@yahoo.com)
11. Assistant Professor, APNNA Institute of Public Health, Jinnah Sindh Medical University, Karachi. Pakistan. [farhana.shahid@jsmu.edu.pk](mailto:farhana.shahid@jsmu.edu.pk)

**\*Corresponding Author: Edwin van Teijlingen**

---

### Abstract

**Introduction:** Caesarean section (CS) is a life-saving surgical intervention for delivering a baby when complications arise in childbirth. World Health Organization recommends a rate of CS from 10% to 15%. However, CS rates increased steadily in recent decades and have almost doubled from 12.1% in 2000 to 21.1% in 2015. Therefore, this has become a global public health problem. The main purpose of the scoping review article is to give an overview and analysis of the rising CS use in four South Asian countries: Bangladesh, India, Nepal and Pakistan.

**Methods:** A scoping review was carried-out using several bibliographic electronic databases (MEDLINE, EMBASE, SCOPUS, CINAHL and Web of Science), organizational websites and open access journal databases. Literature was searched from December 2011 to December 2018 for articles reporting hospital-based CS rates. Inclusion criteria were primary studies conducted in institutional setting in Bangladesh, India, Nepal and Pakistan and published in the English language.

**Results:** We have included 43 studies. Together these studies show that the rate of CS is increasing in all four countries: Nepal, Bangladesh, Pakistan and India. However, this is uneven with very low rates in rural and very high rates in urban settings, the co-existence of ‘Too Little Too Late & Too Much Too Soon’. Hospital based studies have shown that the CS rate is higher in urban and private hospitals. Age, education and socio-economic status of women, urban residence and distance from health facility are associated with CSs. CS is higher among highly educated affluent urban women in private hospitals in South Asian Countries.

**Conclusion:** Rising CS rates in South Asian cities, particularly in specific groups of women, present a challenge to hospital staff and managers and policy-makers. The challenge is to avoid ‘Too Much Too Soon’ in otherwise healthy urban women and avoid ‘Too Little Too Late’ in women living in remote and rural area and in poor urban women.

**Keywords:** *Caesarean Section, South Asia, Scoping review, midwifery*

---

## **Introduction**

Caesarean section (CS) is a surgical procedure which is performed to reduce the risk of mortality or morbidity in the mother and foetus. The World Health Organization (WHO) recommends a CS rate of 10 to 15%, irrespective of geographical region, because, from a health point of view, there is no justification to have a rate of CS higher than this.<sup>1</sup> Experts emphasize that, because of the risks associated with it, CS should be performed only based on medical indications<sup>2–5</sup>. In spite of this, CSs are regularly carried out without clear medical indications.<sup>5</sup> At a population level, a rate of CS higher than 10-15% is not associated with a reduction in maternal and new-born mortality rate.<sup>2</sup> Nevertheless, it is advanced obstetric care which has been gaining popularity in the modern world particularly in urban settings.<sup>3-7</sup>

On the one hand, a huge rise in CS use, often for non-medical reasons, is harmful for both mother and baby. On the other hand, inadequate access to CS in most low-income countries and several middle-income countries is a major health issue. Underuse of CS (<

10%) leads to perinatal morbidity and mortality.<sup>2-7</sup> Thus, the WHO statement on caesarean section rates emphasises that CS should be undertaken when medically essential, rather than striving to attain a specific rate and every effort should be focused on providing CS to all women in need.<sup>2</sup>

### ***Health and social problems associated with Caesarean Birth***

CS is a lifesaving intervention, when medically indicated for specific complications during childbirth. However, the WHO states that CS can cause significant or permanent complications, death and disability especially in setting where the infra-structure is not well-developed and material resources are short.<sup>2</sup>

*The Lancet* series (2018) on optimising caesarean section use has emphasized the significance of increased understanding of short-term and long-term health effects of CS on women and children.<sup>9</sup> These include severe maternal outcomes of CS such as admission to intensive care units, severe haemorrhage/ blood transfusion, hysterectomy, complications associated with anaesthesia, obstetric shock, thromboembolism, major puerperal infection, cardiac failure, acute renal failure, in-hospital wound disruption, intubation and haematoma.<sup>9</sup> High rates of CS are positively associated with postpartum antibiotic treatment and severe maternal morbidity and mortality.<sup>4</sup>

There are many adverse outcomes for women after CS in subsequent pregnancies, which is particularly significant in areas with limited access to emergency obstetric care. These include increased risk of abnormal placentation (placenta previa), hysterectomy, uterine rupture, ectopic pregnancy, stillbirth, preterm birth and miscarriage.<sup>9,10-15</sup> Multiple CS birth increases the risk of severe maternal morbidity in subsequent pregnancies.<sup>10-11</sup> Although there are some benefits of CS identified, such as less frequency of urinary incontinence and urogenital prolapse, multiple CSs can cause undesirable long-term health effects such as pelvic adhesions, small bowel obstruction, menorrhagia, dysmenorrhoea, sexual dysfunction, chronic pain and subfertility.<sup>9,12,14</sup> Furthermore, a study revealed that CS can have negative effects on the quality of life of primiparous women.<sup>15</sup>

In addition to negative maternal outcomes, infants born by CS incur adverse health outcomes due to exposure to a different medical, hormonal, bacterial and physical environment as compared to infants born vaginally. The *Lancet* series (2018) has highlighted many short-term health risks for children born by a caesarean birth, such as allergy, atopy, asthma, alteration of immune development and reduced intestinal gut microbiome diversity.<sup>9</sup>

Children born by CS are more prone to several chronic health conditions such as leukaemia, bowel diseases, juvenile arthritis, asthma, systematic connective tissue disorders and inflammatory immune deficiencies.<sup>16</sup> Additionally, CS is negatively associated with successful breast feeding initiation.<sup>17</sup> A study reported that infants born by elective CS had significantly higher rates of mortality, respiratory morbidity and risk of special care admission as compared to planned vaginal delivery.<sup>18</sup> Similarly, other studies showed associations with childhood obesity,<sup>12,19</sup> asthma<sup>12</sup> and type 1 diabetes<sup>20</sup>.

### ***Global rising of Caesarean Section Rate***

The rate of CS has risen steadily worldwide, and has now reached unprecedented levels.<sup>3-7</sup> Betran et al. (2016) reported the rising global CS rate with 12.4% increase from 1990 (6.7%) to 2014(19.1%) with the highest CS rate being in the Caribbean (40.5%), followed by Northern America (32.3%), Oceania (31.1%), Europe (25.0%), Asia (19.1%) and the lowest in Africa (7.3%).<sup>6</sup> *The Lancet* series (2018) highlighted the global rising trend of CS rate of 21.1% in 2015 (29.7 million births), which was almost double that of 2000 (12.1%, 16.0 million births). According to this report, the CS rate was the highest in Latin America and the Caribbean (44.3%) and lowest in West and Central Africa (4.1%).<sup>7</sup>

CS utilisation was higher among rich and educated women.<sup>7</sup> Huge inequality in use of CS exists between, within countries and higher and lower resources settings.<sup>3-7</sup> CSs was very low in South Sudan (0.6%) and very high in the Dominican Republic (58.1%).<sup>7</sup> Out of 169 countries, CS rate was still less than 10% in 47 countries with women are struggling to receive lifesaving emergency obstetric care. The wide variations in use of CS between high-income and middle/low-income countries as well as urban and rural settings exist. The rising rate of CS world-wide has become an alarming issue in public health.<sup>3-7</sup> In the growing economies of South Asia, the number of hospitals and especially private ones is growing rapidly and therefore, the risk of increased CS birth. This scoping review aims to highlight the prevalence of CS in Bangladesh, India, Nepal and Pakistan.

### **Methods**

A scoping review of the published literature of CS rates was carried-out using bibliographic electronic databases such as MEDLINE, EMBASE, SCOPUS, CINAHL and Web of Science. In addition, organizational websites such as WHO, and open access journal databases such as Nepal journals on-line (NepJOL) and Bangladesh journals on-line (BanglaJOL) were also searched. Databases were searched from January 2011 to December

2018 especially for articles reporting hospital-based CS rates. Inclusion criteria were primary studies reporting the CS rates, conducted in institutional setting in Bangladesh, India, Nepal and Pakistan and published in English language. Mesh terms and Key words for ‘caesarean’; ‘cesarean’; C-section’ was combined with the specific country using Boolean operators (and/or). Titles and abstracts of the identified citations were initially scanned for the rates of caesarean sections to assess eligibility. Full text articles of eligible studies were appraised, and relevant data was extracted, and proportions of C- sections were reported. Articles highlighting the issues around the increase in CSs and the potential reasons were also included, and a simple content analysis was conducted.<sup>21</sup>

**Results**

We initially found 1518 primary studies but most articles were on obstetric issues and only mentioned CS as a cause or consequence. After scanning of titles and abstracts we included 43 papers, five were from Bangladesh, 18 from India, 12 from Nepal, and eight were from Pakistan. Most prevalence studies were retrospective studies of hospital records. The results showed that the CS is steadily rising in all four South Asian countries (especially in urban settings).<sup>6,13-15</sup> However, as in Africa, there is a huge inequality in access to CS use between rich and poor women.<sup>22</sup> CS use is higher among highly educated urban women in private hospitals in South Asian Countries.<sup>23</sup> There are concerns about the rising rate of CS with wide variation between urban and rural, private and public hospital, and with no evidence of associated reductions in morbidities or mortalities.

***Rising Caesarean Section Rate in Bangladesh***

Caesarian sections have become increasingly common in Bangladesh (Table 1) and have increased alarmingly in recent decades. Despite the rising trend of institutional births from 23% in 2010 to 47% in 2016, maternal deaths are still as high as 194 per 100,000 live births.<sup>24</sup> In Bangladesh, CS rate increased from 3.5% in 2004 to 23.0% in 2014 with wide variation in access to CS use between urban (40%) and rural women (18.7%).<sup>25</sup> Among women, older age and high educational status;<sup>25-28</sup> residing in urban area and obesity;<sup>25,26</sup> high socio-economic status;<sup>25,27</sup> access to antenatal care<sup>25,26,28</sup> and birth order<sup>25,27</sup> were found to be associated with CS delivery.

Table 1: CS rates in hospital-based studies in Bangladesh

<b>Authors &amp; year</b>	<b>Hospitals in Bangladesh</b>	<b>CS rate (%)</b>
---------------------------	--------------------------------	--------------------

Nazneen et al., 2011 <sup>29</sup>	Holy Family Red Crescent Medical Collage Hospital, Dhaka	70.55*
Aminu et al., 2014 <sup>30</sup>	Five Rural Public Hospitals, Thakurgoan Districts	21.2 <sup>+</sup>
Husan et al., 2015 <sup>31</sup>	Four Hospitals, Rajshahi City	77.3*
Islam et al., 2015 <sup>32</sup>	Six Hospitals, Narsingdi	32.3 <sup>#</sup>
Ara et al. 2018 <sup>33</sup>	Combined Military Hospital (CMH), Savar, Dhaka	70.3*

\*Urban hospital; # semi-urban hospital; + rural hospital

### **Rising Caesarean Section Rate in India**

CS rates are steadily increasing in India with 2.9% in 1992/93, 7.1% in 1998/99, 8.5% in 2005/6 and 17.2% in 2015/16.<sup>34</sup> Similar to other South Asian countries, there was disparity in CS rate between urban and rural areas as well as different states of India. In 2015/16, CS rate was 12.9% in rural India as compared to 28.3% in urban areas. The lowest CS rate was in Nagaland (5.8%) and the highest rate in Telangana (58.0%). CS rate was high in Andhra Pradesh (40.1%), Kerala (35.8%), Tamil Nadu (34.1%), Jammu & Kashmir (33.1%) and Goa (31.1%). The proportion of institutional birth has increased from 38.7% in 2005/06 to 78.9% in 2015/16.<sup>34</sup> It is also worth noting that the CS rate has increased from 26.6% in 2007/08 to 40.7% in 2010/11 in the state of Madhya Pradesh in India, where the National Janani Swastha Yojana (YSY) scheme was implemented. YSY is a conditional cash transfer paid to the mother when they present to deliver their baby in a health facility.<sup>35</sup> Interestingly, in 2003, Sreevidya and Sathuyasekaran reported a high CS rate (32.6%) in urban India with higher CS rate in the private sector (47%) as compared to charitable (38%) and public (20%).<sup>36</sup> That trend has continued, with increased CS rate in private hospitals from 27.7% in 2005/6 to 40.9% in 2015/16 but it declined in public hospitals from 15.2% in 2005/6 to 11.9% in 2015/16.<sup>36</sup> Similarly, Mittal and colleagues also reported a rising trend of CS rate in tertiary care hospital in Western India where CS rate has increased from 69.03% in 2001 to 72.62% in 2011.<sup>37</sup> Age and education of mother, parity, previous history of CS and distance from the health facility were found to be important determinants of CS.<sup>38</sup> Furthermore, hospital-based studies stress the rise of CS rate in India (Table 2).

Table 2: CS rates in hospital-based studies in India

<b>Authors &amp; year</b>	<b>Hospitals in India</b>	<b>CS rate (%)</b>
Padamleela et	Government Teaching Hospital, Andra Pradesh	31*



---

al.,2013 <sup>39</sup>		
Santhanalakshmi et al., 2014 <sup>40</sup>	Karpaga Vinayaga Inst Med Sciences, Kancheepura	12.5 <sup>#</sup>
Bade et al.,2014 <sup>41</sup>	Government Medical College, Latpur	23.97 <sup>#</sup>
Yadav RG, 2015 <sup>42</sup>	Baroda Medical Collage, Vadadara	25.17 <sup>#</sup>
Nikhil et al., 2015 <sup>43</sup>	GMERS Medical College, Ahmedabad	25.18*
Subhashini et al., 2015 <sup>44</sup>	Adhra Medical Colleage, Andra Pradesh	25.66*
Yadav et al., 2016 <sup>45</sup>	MMIMSR Mullana, Haryana	21.60 <sup>#</sup>
Sarma et al., 2016 <sup>46</sup>	Tezpur Medical Colleage & Hospital, Assam	27.60*
Saxena et al., 2016 <sup>47</sup>	Shri Guru Ram Rai Inst Med & Health Sci, Dehradun	31.18*
Jawa et al., 2016 <sup>48</sup>	Mahatma Gandhi Medical Sciences & hospital, Jaipur	31.80*
Das et al., 2016 <sup>49</sup>	NEIGRIHMS, Shillong	33.1*
Preetkamal et al., 2017 <sup>50</sup>	Sri Guru Das Inst Med Sci Research, Punjab	33.2*
Chavada et al., 2017 <sup>51</sup>	P. D. U Medical Colleage, Gujrat	19.9 <sup>#</sup>
Gupta et al., 2017 <sup>52</sup>	RUSH Colleage of Medical Sciences, Jaipur	31.46*
Bala et al., 2017 <sup>53</sup>	JK Ion Hospital & Medical Colleage, Kota	39.6*
Kaul et al., 2017 <sup>54</sup>	Defence Zonal hospital, Varanasi	45.5*
Banergee et al.,2018 <sup>55</sup>	Silchar Medical Collage, Assam	34.1*
Das et al.,2018 <sup>56</sup>	IMS& SUM Medical Colleage, Odisha	34.45*

\*Urban hospital; # semi-urban hospital; + rural hospital

### ***Rising Caesarean Section Rate in Nepal***

The prevalence of CS increased more than four times from 2001 to 2011 in Nepal.<sup>57</sup> Similarly, the rate of CS was 4.6% in 2011,<sup>58</sup> 8.6% in 2014<sup>59</sup> and 9.0% in 2016.<sup>60</sup> However, there is wide inequality in access to CS between urban and rural women. Studies have reported that CS rate is significantly higher (12-19%) in urban areas than in rural Nepal (3.5-7.1%).<sup>57-60</sup> Additionally, the most recent Nepal Demographic and Health Survey (2016) discovered that CS rate is substantially higher in private hospitals (35.5%) than public hospitals (12.5%).<sup>59</sup> Some studies showed that older women, women with good education, residing in city and rich women are more likely to undergo CS delivery.<sup>57-58</sup> The Government of Nepal has been promoting safe motherhood through initiatives such as offering free

delivery care and transportation incentive schemes to women who give birth in hospital. The percentage of institutional birth has increased from 35% in 2011 to 57% in 2016.<sup>60</sup> The growing trend of birth in hospital is contributing to rising rate of CS in Nepal.<sup>7</sup> Amatay et al reported an increase in CS rates at Tribhuvan University Teaching hospital from 16.6% in 2005 to 25.4% in 2009.<sup>61</sup> Furthermore, individual hospital-based studies on CS in different hospitals in Nepal show the rising CS rate (Table 3).

Table 3: CS rates in hospital-based studies in Nepal

Authors & year	Hospitals in Nepal	CS rate (%)
Chhetri et al., 2011 <sup>62</sup>	B P Koirala Health Science Teaching Hospital, Dharan	33.7*
Subedi, 2012 <sup>63</sup>	Nobel Medical College, Biratnagar	19.89*
Suwal et al., 2013 <sup>64</sup>	Nepal Medical College Hospital, Kathmandu	22.30*
Amatya et al., 2014 <sup>65</sup>	Khadbari District Hospital, Sankhuwasabha	18.5 <sup>#</sup>
Pradhan et al., 2014 <sup>66</sup>	Kirtipur Hospital, Kirtipur	50.9*
Pradhan et al., 2015 <sup>67</sup>	Patan hospital, Lalitpur	41.9*
Samdal et al., 2016 <sup>68</sup>	Okhaldunga Community Hospital, Okhaldunga	9.5+
Singh et al., 2017 <sup>69</sup>	Western Regional Hospital, Pokhara	25*
Prasad et al., 2017 <sup>70</sup>	Kathmandu Medical College Teaching hospital	48.81*
Dhakalet al., 2018 <sup>71</sup>	Mid-Western Regional Hospital, Surkhet	18.9 <sup>#</sup>
Vaidya Malla et al., 2018 <sup>72</sup>	Nepalese Army Shree Birendra Hospital, Kathmandu	22.57*
Chaudhary et al., 2018 <sup>73</sup>	Paschimanchal Community hospital, Pokhara	63.2*

\*Urban hospital; <sup>#</sup> semi-urban hospital; <sup>+</sup> rural hospital

### ***Rising Caesarean Section Rate in Pakistan***

Similar trends and inequalities of CS rates are observed in Pakistan, where the CS rate increased from 2.7% in 1990–'91 to 15.8% in 2012–'13 with a big difference between urban (25.6%) and rural (11.5%) as well as the poorest (5.5%) and the richest (35.3%).<sup>74</sup> Similarly, highly educated women (40.3%) are more likely to have access to CS delivery than non-

educated women (7.7%).<sup>74</sup> Hospitals based studies reflect rising CS rate in Pakistan (Table 4).

Table 4: CS rates in hospital-based studies in Pakistan

Authors & year	Hospitals in Pakistan	CS rate (%)
Karim et al., 2011 <sup>75</sup>	Civil hospital & Dow university health science, Karachi	27.94*
Jabeen et al., 2013 <sup>76</sup>	CMH. Rawalpindi	56*
Hafeez et al., 2014 <sup>77</sup>	Sharif Medical & Dental collage, Lahore	21.4 <sup>#</sup>
Bano et al., 2015 <sup>78</sup>	Aga Khan Hospital for women, Karachi	31.26*
Baig et al., 2016 <sup>79</sup>	CMH, Hyderabad	41.96*
Latif et al., 2017 <sup>80</sup>	Nawas Sharif Social Security hospital, Lahore	81%*
Naeem et al., 2018 <sup>81</sup>	Government Sardar Begam Hospital, Sialkot	40%*
Tahir et al., 2018 <sup>82</sup>	Combined Military hospital, Abbottabad	46.7%*

\*Urban hospital; <sup>#</sup> semi-urban hospital; <sup>+</sup> rural hospital

There are wide variations also in the CS rates across South Asia, comparing urban and rural populations/hospitals. The private health providers/hospitals may be partly influencing the rise of CS rate in South Asian countries.<sup>23</sup> The rising rates in the cities are possibly linked to ‘unnecessary intervention of CS. However, the actual reasons behind this and the wide variations reported across facilities are unclear.

## Discussion

The data presented above show a range of CS rates within each country and with a trend of increases over time. There are a range of possible explanations as listed in the next section.

### Reasons of Caesarean Section

There are many medical and non-medical reasons for performing CS delivery. In South Asian countries, the most common indications of CS are foetal distress<sup>33,39,40,42-47,49,50,52-56,62-64,66,67,75,76,78-81</sup>, foetal malpresentation/malposition<sup>29-32,39,40,42-44,46,47,49-56,62,63,65-</sup>

73,75,76,78-81, previous CS<sup>29-31,39,40,42-56,62-64,65,73,75-78</sup>, non-progressing labour/prolong/obstructed labour<sup>29-33,40,42,44-50,51,54,56,62-67,68-73,75-81</sup>, failed induction<sup>29,30,42,44-46,50,52,64-72,75,79,80</sup>, cephalo-pelvic disproportion (CPD)<sup>39,30,39,40,42,43,44,47,49,40,52-56,62,63,65,66,68,70,72,73,75,78</sup>, pregnancy induced hypertensive disorder including pre-eclampsia/eclampsia<sup>29-31,40,42,44,45,47,49,50,56,63,72,78-81</sup>, and oligohydramnios<sup>30,31,40,42,43,47,49,50,56,65,66,69,70,72,73,75</sup>.

There are several other indicators for CS which are noteworthy such as antepartum haemorrhage<sup>29,31,33,40,42,45,50,53,63,69,71,75,78</sup>, multiple pregnancy<sup>39,40,49,52-56,62,63,69,70,72</sup>, intra uterine growth retardation (IUGR)<sup>39,40,43,44,46,49,51,56,75</sup>, placenta previa/abruptio<sup>29,31,33,44,49,72,73,81</sup>, postdate pregnancy<sup>30,31,43,46,47,53</sup>, meconium stained liquor<sup>32,62,65,67,70,71</sup>, bad obstetric history/complicating pregnancy<sup>39,47,53,56,69,72,81</sup> and big baby<sup>33,43,44,52,56</sup>. Similarly, some studies show premature rupture of membrane (PROM)<sup>30,31,68</sup>, cord prolapse<sup>33,66</sup>, chorio-amnionitis<sup>79,80</sup>, cervical dystocia<sup>32,33</sup>, obesity<sup>33</sup>, previous traumatic delivery<sup>68</sup> and congenital malformation as causes for CS.<sup>52</sup>

Most importantly, some studies have highlighted that CSs are performed for non-medical indications such as demand of women and family.<sup>39,43,53,56,63,66,75,79-</sup> Some of the demand for CSs are due to either bad obstetric history or precious pregnancy due to the subfertility.<sup>39,75,79,80</sup><sup>81</sup> This trend is increasing in South Asia.

### ***The role of midwifery in keeping childbirth normal***

There is international consensus that midwifery care is the most cost effective way of supporting normal childbirth. Midwives are trained to use minimal intervention but to refer timely when complications arise. Midwifery care can decrease maternal and new-born mortality in low and lower-middle income countries.<sup>83</sup>,<sup>84</sup> To ensure positive maternal and newborn health outcomes, high quality intrapartum care, especially through the promotion of spontaneous vaginal births with a minimum of medico-technical interventions is central.<sup>85,86</sup> The outcome of the care for women and newborns around the time of birth in health facilities reflects the evidence-based practices used and the overall quality of services provided. The quality of care during childbirth in health facilities depends on the physical infrastructure, human resources, knowledge, skills and capacity to deal with both normal pregnancies and complications that require prompt, life-saving interventions.<sup>87</sup> There is evidence to suggest that midwives who work in the communities in which they live and are known are most effective at delivering effective care.<sup>88</sup>

### ***Combating 'Too little, too late' (TLTL) and 'Too much too soon' (TMTS)***

TLTL refers to lack of resources, low standard of quality of care/services and unavailability of care until too late. TMTS describes the over use of emergency obstetric care as well as unnecessary use of non-evidence-based intervention and over-medicalisation of normal pregnancy and childbirth.<sup>89</sup> The rate of CS has been increasing in developing countries with socio-economic disparity.<sup>22</sup> Massive inequality exists in access to CS among poor and middle-income developing countries.<sup>6,7,21</sup> Similarly, there is wide variation in CS rate between rural and urban areas as well as between private and public hospitals.<sup>23</sup> In South Asia, there is a double burden with low numbers of CSs in poor remote rural areas and very high in wealthier urban areas. The former is due to underdeveloped local health systems in remote areas with limited provision of timely and safe CS procedure to save the lives of mother and fetus.<sup>22</sup> Provision of adequate access to skilled care is essential in rural areas. In contrast, emergency obstetric care is easily accessible in urban areas. Easily available private facilities (oversupply of hospital beds) and education of women have been suggested as factors for the rising rates of CS in urban settings.<sup>23</sup> Dhakal et al recently highlighted the rising rates of CS in urban Nepal.<sup>89</sup> Overuse of CS may cause not only severe maternal and foetal outcome but also financial burden to low-income countries and low and middle-income families in those countries.

The rising CS rate in South Asian countries is an alarming public health problem. The issue of growing CS rate in South Asia need to be reduced in effective ways by implementing evidence-based interventions to reduce unnecessary caesarean sections in healthy women. *The Lancet* series (2018) highlights the need for multicomponent and locally tailored interventions which address both women's and professionals' concerns as well as health system and financial factors.<sup>90</sup> Similarly, same fee for CS and vaginal delivery, comprehensive information on benefits and risk of CS to women, uniform classification system for CS and annual CS rate publication by hospitals are also needed to reduce the high CS rate.<sup>91</sup> Routine information on all aspect of childbirth and adoption of standard classification systems (Robson's 10 groups classification) are needed for comparison and improvement of CS use.<sup>92</sup>

## **Conclusion**

The CS rates are growing at an alarming rate in urban settings in South Asian countries warranting immediate attention. Further research needs to be conducted to explore the factors/reasons associated with rising CS in urban settings both from the patient's and the health system perspectives. This will help to develop appropriate tailored interventions.

Improving the quality of intrapartum care through midwifery care in health facilities is an important focus in the pursuit to reducing unnecessary CS and end preventable mortality and morbidity among mothers and newborns.

### Conflict of Interest

None declared by authors

### References

1. World Health Organization (1995). Appropriate technology for birth. *Lancet*, 2(8452):436-7.
2. World Health Organization (2015). *WHO statement on Caesarean section rates*. Available: [http://apps.who.int/iris/bitstream/handle/10665/161442/WHO\\_RHR\\_15.02\\_eng.pdf;jsessionid=37B37C57EB23CA66A19CAC0098A553E9?sequence=1](http://apps.who.int/iris/bitstream/handle/10665/161442/WHO_RHR_15.02_eng.pdf;jsessionid=37B37C57EB23CA66A19CAC0098A553E9?sequence=1), accessed on 01/01/2019.
3. Lumbiganon P, Laopaiboon M, Gülmezoglu AM, et al. (2010). World Health Organization Global Survey on Maternal and Perinatal Health Research Group: Method of delivery and pregnancy outcomes in Asia: the WHO global survey on maternal & perinatal health 2007-08. *The Lancet*, 375: 490-98.
4. Villar J, Valladares E, Woidyla D, et al. (2006). WHO 2005 global survey on maternal and perinatal health research group. Caesarean delivery rates and pregnancy outcomes: the 2005 WHO global survey on maternal & perinatal health in Latin America. *The Lancet*, 367(9525):1819-29.
5. Souza J P, Gulmezoglu A M, Lumbiganon P, et al. (2010). The WHO global Survey on maternal health and perinatal health research group. Caesarean section without medical indications is associated with an increased risk of adverse short-term maternal outcomes: the 2004-2008 WHO Global Survey on Maternal and Perinatal Health. *BMC Medicine*, 8:71.
6. Betran AP, Ye J, Moller AB, Zhang J, et al. (2016). The increasing trend in caesarean section rates: Global, Regional and National estimates: 1990-2014. *PLOS ONE*, 5;11(2): e0148343.
7. Boerma T, Ronsmans C, Barros AJ, et al. (2018). Global epidemiology of use of disparities in caesarean sections. *The Lancet*, 392(10155):1341-1348.
8. Popay P, Roberts H, Sowden A, et al. (2006) *Guidance on the Conduct of Narrative Synthesis in Systematic Reviews*. A Product from the ESRC Methods Programme. Lancaster, UK.

9. Sandall J, Tribe RM, Avery L, et al. (2018). Short-term and long-term effects of caesarean section on the health of women and children. *The Lancet*, 392(10155):1349-1357.
10. Marshall NE, Fu R, Guise JM (2011). Impact of multiple cesarean deliveries on maternal morbidity: a systematic review. *American Journal of Obstetrics & Gynecology*, 205 (3):262.
11. Keag OE, Norman JE, Stock SJ (2018). Long-term risks and benefits associated with cesarean delivery for mother, baby, and subsequent pregnancies: systematic review and meta-analysis. *PLoS Medicine*, 15: e1002494.
12. O'Neill SM, Kearney PM, Kenny L, et al. (2013). Caesarean delivery and subsequent stillbirth or miscarriage: systematic review and meta-analysis. *PLOS ONE*, 8: e5458.
13. O'Neill SM, Kearney PM, Kenny LC, et al. (2013). Caesarean delivery and subsequent pregnancy interval: a systematic review and meta-analysis. *BMC Pregnancy & Childbirth*, 13:165.
14. Guise JM, Denman MA, Emeis C, et al. (2010). Vaginal birth after cesarean: new insights on maternal and neonatal outcomes. *Obstetrics & Gynecology*, 115: 1267-1278.
15. Mousavi SA, Mortazavi F, Chaman R, Khosravi A. (2013). Quality of life after caesarean section and vaginal delivery. *Oman Medical Journal*, 284:245-251.
16. Sevelsted A, Stokholm J, Bonnelykke K, Bisgaard H (2015). Cesarean Section and Chronic Immune Disorders. *Pediatrics (online)*, 135(1): e92-8.
17. Perez-Rios N, Ramos-Valencia G, Ortiz AP. (2006). Caesarean section delivery as a barrier for breast feeding initiation: The Puerto Rican experience. *Journal of Human Lactation* 24(3):293-302.
18. De Luca R, Boulvain M, Irion O, Berner M, Pfister RE (2009). Incidence of early neonatal mortality and morbidity after late-preterm and term caesarean delivery. *Pediatrics (online)* 123(6): e1064-71.
19. Li HT, Zhou YB, Liu JM (2013). The impact of cesarean section on offspring overweight and obesity: a systematic review and meta-analysis. *International Journal of Obesity* 37(7):893-69.
20. Bonifacio E, Warncke K, Winker C, et al. (2011). Caesarean section and inferno-induced helicase gene polymorphisms combine to increase childhood type 1 diabetes risk. *Diabetes* 60(12);3300-3306.

21. Popay, J., Roberts, H., Sowden, A., et al. (2006). Guidance on the conduct of narrative synthesis in systematic reviews. A product from the ESRC methods programme Version, 1:b92
22. Ronsmans C, Holtz S, Stanton C (2006). Socioeconomic differentials in caesarean rates in developing countries: a retrospective analysis. *The Lancet* 368: 1516-1523.
23. Newman M, Alcock G, Azad K et al. (2014). Prevalence and determinants of caesarean section in private and public health facilities in underserved south Asia communities: cross-sectional analysis of data from Bangladesh, India and Nepal. *BMJ Open* (online), 4: e005982.
24. National Institute of Population Research and Training (NIPORT), B.I. International Centre for Diarrhoeal Disease Research, b), and MEASURE Evaluation (2016). *Bangladesh Maternal Mortality and Health Care Survey (BMMS): Preliminary Report*: Dhaka, Bangladesh, and Chapel Hill, NC, USA: NIPORT, icddr,b, and MEASURE Evaluation.
25. Rahaman MM, Haider MR, Moinuddin M et al. (2018). Determinants of caesarean section in Bangladesh; Cross-sectional analysis of Bangladesh Demographic and health survey 2014 data. *PLOS ONE (Online)*, 13(9): e0202879.
26. Khan MM, Islam MM, Sariff AA, et al. (2017). Socio-demographic predictors and average annual rates of caesarean section in Bangladesh between 2004 and 2014. *PLOS ONE (Online)*, 12(5): e0177579.
27. Rahman M, Shariff AA, Shafie A, et al. (2014). Determinants of caesarean risk factor in northern region of Bangladesh: a multivariate analysis. *Iranian Journal of Public Health*, 43(1), 16.
28. Begum T, Rahman A, Nababan H et al. (2017). Indication and determinants of caesarean section delivery: Evidence from a population-based study in Matla, Bangladesh. *PLOS ONE(Online)*,12(11): e0188074.
29. Nazneen R, Begum RA, Sultana K (2011). Rising trend of caesarean section in a tertiary hospital a decade. *Journal of Bangladesh College of Physicians & Surgeons*, 29(3):126-132.
30. Aminu M, Utz B, Halim A et al. (2014). Reasons for performing a caesarean section in public hospitals in rural Bangladesh. *BMC Pregnancy & Childbirth*, 14:130.
31. Hasan F, Sabiruzzaman M, Joardar C et al. (2015). Maternal socio-demographic factors and nutritional status as predictors of caesarean delivery at hospitals in Rajshahi city Bangladesh. *Malaysian Journal of Nutrition*, 21(3):345-353.



32. Islam MT, and Yoshimura Y (2015). Rate of caesarean section delivery at hospitals providing emergency obstetrics care in Bangladesh. *International journal of Gynaecology & Obstetrics*, 128(1):40-43.
33. Ara I, Sultan R, Solaiman SM, Hassain MMS, Sultana R1 (2018). Current Trend of Caesarean Section in a Tertiary Care Military Hospital. *Bangladesh Medical Research Council Bulletin*, 44):15-22
34. Radhakrishna T, Vasantahakumari KP, Babu PK (2017). Increasing trend of caesarean section rates in India: Evidence from NFHS – 4. *Journal of Medical Science & Clinical Research*, 5(8):26167-176
35. Bogg L, Diwan V, Vora KS, DeCosta A (2016). Impact of alternative maternal demand-side finance support program in India on the caesarean section rate: Indications of supplier-induced demand. *Maternal and Child Health Journal*, 20(1):11-15.
36. Sreevidya S, Sathiyasekaran BWC (2003). High caesarean section rate in Madras (India): a population based cross sectional study. *BIGOJ*, 110:106-111.
37. Mittal S, Pardeshi S, Mayades N, Mane J (2014). Trend in caesarean section: Rate and Indications. *Journal of Obstetrics and Gynaecology of India*, 64(4):251-254.
38. Desai G et al. (2017). Rates, indications and outcomes of caesarean section deliveries: A comparison of tribal and non-tribal women in Gujarat, India. *PLOS ONE*, 12(12): e0189260.
39. Padmaleela K, Thomas V, Prasad KV (2013). An analysis of the institutional deliveries and their outcomes in government teaching hospitals of Andhra Pradesh, India. *International Journal of Health Science & Research (IJHSR)*. 3(5):76-81.
40. Santhanalakshmi C, Gnanasekaran V, Chakravarthy AR (2015). A retrospective analysis of cesarean section in a tertiary care hospital. *International Journal of Science & Research*, 4(9):2097-9.
41. Bade P, Kendre V, Jadhav Y, Wadagale A (2014). An analysis of indications for caesarean section at government medical college, Latur. *International Journal Recent Trends in Science & Technology*, 11(1):6-8.
42. Yadav RG, Maitra N (2015). Examining cesarean delivery rates using the Robson's ten group classification. *Journal of Obstetrics & Gynecology of India*, 66(Suppl 1):1.
43. Nikhil A, Desai A, Vijay K, et al. (2015). Analysis of trends in LSCS rate and indications of LSCS: a study in a Medical College Hospital GMERS, Sola, Ahmedabad. *International Journal of Pharmacy & Biological Sciences*, 2(1):1-5.

44. Subhashini R,Uma N (2015). Changing trends in Caesarean delivery. *International Archives of Integrated Medicine (IAIM)*, 2(3):96-102.
45. Yadav S, Kaur S, Yadav SS, Thakur B (2016). Analysis of caesarean rate, indications and complications: review from medical college Ambala, Haryana, India. *International Journal of Reproduction, Contraception, Obstetrics & Gynecology*, ,5:3326-9.
46. Sarma P, Boro RC, Acharjee PS (2016). An analysis of indications of caesarean sections at Tezpur medical college and hospital, Tezpur (a government hospital). *International Journal of Reproduction, Contraception, Obstetrics & Gynecology*, , 5:1364-7.
47. Saxena N, Sharma B, Gupta V, Negi KS (2016). A six-year appraisal of caesarean delivery at a teaching hospital in Uttarakhand. *International Journal of Reproduction, Contraception, Obstetrics & Gynecology*, 5:4369-72.
48. Jawa A, Garg S, Tater A, Sharma U (2016). Indications and rates of lower segment caesarean section at tertiary care hospital-an analytical study. *International Journal of Reproduction, Contraception, Obstetrics & Gynecology*,5:3466-9.
49. Das A, Panda S, Singh SA (2016). An attempt to the control the increasing trend of caesarean section. *Obstetrics & Gynecology International Journal*, 5(6);00178.
50. Kaur H and Nagpal M (2017). Is current rising trend of cesarean sections justified? *International Journal of Reproduction, Contraception, Obstetrics & Gynecology*,, 6:872-6.
51. Chavda D, Goswam K, Dudhrejiya K (2017). A cross sectional study of 1000 lower segment cesarean section in obstetrics and gynecology department of P.D.U Medical College, Rajkot, Gujarat, India. *International Journal of Reproduction, Contraception, Obstetrics & Gynecology*, 6(4):1186-91.
52. Gupta M and Garg V (2017). The rate and indications of caesarean section in a teaching hospital at Jaipur. *India. Int J Reprod Contracept Obstet Gynecol*, 6:1786-92.
53. Bala S, Patidar BL, Gupta B (2017). A retrospective analysis of annual caesarean section rate in a tertiary care hospital, Kota. *Journal of Medical Science and clinical Research*, 05(07): 25325-25329.
54. Kaul KK and Singh OP (2017). Rising trends of C-section: In a defence zonal hospital: A retrospective study. *International Journal of Scientific Research*, 6(10):1514-1517.
55. Banergee A, Bhadra B, Dey KR (2018). Analysis of caesarean section in a tertiary care hospital, Assam, India. *International Journal of Reproduction, Contraception, Obstetrics & Gynaecology*, 7(4):1514-1517.

56. Das RK, Subudhi KT, Mohanty RK (2018). The rate and indication of caesarean section in a tertiary care teaching hospital eastern India. *International Journal of Contemporary Paediatrics*, 5(50):1733-1739.
57. KC P and Neupane S (2014). Caesarean deliveries among Nepalese mothers: changes over time 2001–2011 and determinants. *Archives of Gynaecology and Obstetrics*, 289(2): 421–427.
58. Ministry of Health and population Nepal (2012). *Nepal Demographic and Health Survey 2011*. Ministry of Health and Population Nepal. Available from:
59. [https://dhsprogram.com/pubs/pdf/FR257/FR257\[13April2012\].pdf](https://dhsprogram.com/pubs/pdf/FR257/FR257[13April2012].pdf) [Accessed on 22/10/2018].
60. UNICEF Nepal, Central Bureau of Statistics (CBS), 2018. *Nepal Multiple Indicator Cluster Survey 2014*. Available from: [gov.np/image/data/2018/Nepal%20Multiple%20Indicator%20Cluster%20Survey%202014%20Final%20Report.pdf](http://gov.np/image/data/2018/Nepal%20Multiple%20Indicator%20Cluster%20Survey%202014%20Final%20Report.pdf). Accessed on 03/01/2019.
61. Ministry of Health and population (2017). *Nepal Demographic and Health Survey 2016*. Available from: <https://www.dhsprogram.com/pubs/pdf/fr336/fr336.pdf>. Accessed on 24/10/2018.
62. Amatya A, Poudel R, Poudel A, et al. (2013). Examining stratified caesarean section rates using Robson classification system at Tribhuvan University Teaching hospital. *Journal of Nepal Health Research Council*, 11(25):255 – 258.
63. Chhetri S and Singh U (2011). Caesarean section: its rate and indications at tertiary referral centre in Eastern Nepal. *Health Renaissance*,9(3):179-183.
64. Subedi S (2012). Rising rate of caesarean section – a year review. *Journal of Nobel Medical College*, 2:72–76.
65. Suwal A, Shrivastav VR, Giri A (2013). Maternal and foetal outcome in elective versus emergency caesarean section. *Journal of Nepal Medical Association*, 52(192):563-6.
66. Amatya YR and Sapkota D (2014). Patterns of Delivery and Perinatal Outcomes Among Women Delivered at District Hospital of Rural Nepal. *Online Journal of Health Allied Sciences*, 3(3):1.
67. Pradhan P, Shrestha S, Rajbhandari P, Dangal G (2014). Profile of Caesarean Section in Kirtipur Hospital. *Nepal Journal of Obstetrics and Gynaecology*, 9(2):51-4.
68. Pradhan B, Duwal Shrestha S RC L, et al. (2015). Increasing Trend of Caesarean Section in Patan Hospital. *Journal of General Practice & Emergency Medicine in Nepal*, 4(6):3-5.

69. Samdal LJ, Steinsvik KR, Pun P, et al. (2016). Indication of cesarean section in rural Nepal. *Journal of Obstetrics and Gynaecology India*, 66(Suppl 1):284-8.
70. Singh D, Regmi R, Gurung T, Sunar L (2018). Cesarean Delivery and its indication: a Cross Sectional Study in a Tertiary Care Hospital, Pokhara, Nepal. *Nepal Journal of Obstetrics & Gynaecology*, 12(2), 79-82.
71. Prasad A, Prasad A, Bhandari G, Saha R. (2017). Profile of caesarean section in Kathmandu Medical Collage. *Journal of Nepal Health Research Council*, 15(36):110-13.
72. Dhakal KB, Dhakal S, Bhandari S (2018). Profile of caesarean section in Mid – Western Regional hospital in Nepal. *Journal of Nepal Health Research Council*, 16(38):84-8.
73. Malla-Vaidya R, Hamal C, Neupane B, Khatri R (2018). Analysis of caesarean section using Robson’s 10-group classification at a tertiary level hospital in Nepal. *Med J Shree Birendra Hospital*,17(2):4-11.
74. Chaudhary R, Raut KB, Pradhan K (2018). Prevalence and Indications of Cesarean Section in a Community Hospital of Western Region of Nepal. *Journal of the Nepal Medical Association* 56(213).
75. Mumtaz S, Bahk J, Khang YH (2017). Rising trends and inequalities in caesarean section rates in Pakistan:Evidence from Pakistan Demographic and Health Surveys, 1990-2013. *PLOS ONE (Online)*,12(10): e0186563.
76. Karim F, Ghazi A, Ali T,Aslam R (2011). Trends and determinants of caesarean section. *Journal of Surgery Pakistan (International)*, 16 (1):22-27.
77. Jabeen J, MansoorM. H, Mansoor A. (2013). Analysis of Indications of Caesarean Sections. *Journal of Rawalpindi Medical College (JRMCI)*; 2013;17(1):101-103.
78. Hafeez M, Yasin A, Badar N et al. (2014). Prevalence and indications of caesarean section in a teaching hospital. *Journal of International Medical Sciences Academy (JIMSA)*, 27(1): 15-16.
79. Bano R, Mushtaq A, Adhi M et al. (2015) Rates of caesarean section and trials and success of vaginal birth after caesarean sections in secondary care hospital. *Journal of Pakistan Medical Association*, 65(1): 81-83.
80. Baig JR, Jamal MM, Ashfaq T (2016). Two-year analysis of caesarean section delivery at CMH Hyderabad. *Pakistan Armed Forces Medical Journal*, 66(1):25-29.
81. Latif R, Farfiq S, Ashfaq M et al. (2017). An analysis of prevalence and indication of caesarean section in primigravida. *Pakistan Journal of Medical & Health Science*, 11(1):9-11.

82. Naeem A, Answer A, Sajjad S (2018). Caesarean section: Introduction and rate of caesarean section in Primigravida. *Professional Medical Journal*, 25(9):1338-1341.
83. Tahir N, Adil M, Fatima S, Khan S (2018). Caesarean sections: Frequency and indications at peripheral tertiary care hospital. *Pakistan Armed Forces Medical Journal*, 68(2):273-79.
84. Renfrew, M.J., McFadden, A., Bastos, M.H. et al. (2014) Midwifery and quality care: findings from a new evidence-informed framework for maternal and newborn care. *Lancet*, 384(9948):1098.
85. Homer CS, Friberg IK, Dias MA et al. (2014). The projected effect of scaling up midwifery. *The Lancet*, 384(9948):1146-57.
86. Koblinsky M, Moyer CA, Calvert C et al. (2016). Quality maternity care for every woman, everywhere: a call to action. *Lancet*, 388(10057):2307-2320.
87. Sharma G, Powell-Jackson T, Halder K et al. (2016) Quality care during labour and birth: a multi-country analysis of health system bottlenecks and potential solutions. *BMC Pregnancy Childbirth*, 15 Suppl 2: S2.
88. Miller S, Abalos E, Chamillard M, Ciapponi A et al. (2016). Beyond too little, too late and too much too soon: a pathway towards evidence-based, respectful maternity care worldwide. *The Lancet*, 388(10056):2176-92.
89. Hoopé-Bender P, Bernis L, Campbell J, Downe, S, Fauveau V et al. (2014) Improvement of maternal and newborn health through midwifery. *The Lancet*, 384 (9949), 1226-35.
90. Dhakal Rai, S, Regmi, P, van Teijlingen, E, et al.. (2018). Rising Rate of Caesarean Section in Urban Nepal. *Journal of Nepal Health Research Council*, 6(41): 479-80.
91. Betran AP, Temmerman M, Kingdon C, et al. (2018). Interventions to reduce unnecessary caesarean section in healthy women and babies. *The Lancet*, 392(10155):1358-1368.
92. Visser GHA, Ayer-de-Compos D, Barnea ER, et al.( 2018). FIGO position paper: how to stop the caesarean section epidemic. *The Lancet*, 392(10155):1286-1297.
93. Robson MS (2001). Can we reduce the cesarean section rate? *Best Practice & Research Clinical Obstetrics & Gynecology*, 15:179-94.