

COVID-19 vaccine hesitancy concerns: Findings from a Ghana Clinical Radiography Workforce Survey

Authors:

¹Benard Ohene Botwe, PhD: Sirbenard13@gmail.com /bebotwe@ug.edu.gh:
ORCID: 0000-0002-0477-640X

¹William Kwadwo Antwi, PhD: wkantwi@ug.edu.gh:
ORCID: 0000-0002-6726-2342

¹Jennifer Amponsem Adusei, BSc: jenniadu21@gmail.com

²Raphael Nicholas Mayeden, MD: nobility12@yahoo.com

³Theophilus N. Akudjedu, PhD: takudjedu@bournemouth.ac.uk:
ORCID: 0000-0003-2423-6897

¹Seyram Derick Sule (PhD): sdsule@chs.edu.gh

Affiliations:

¹Department of Radiography, School of Biomedical and Allied Health Sciences, College of Health Sciences, University of Ghana, Box KB143, Korle Bu, Accra, Ghana.

²University of Health and Allied Sciences, Ho, Ghana.

³Institute of Medical Imaging & Visualisation, Department of Medical Science & Public Health, Faculty of Health & Social Sciences, Bournemouth University, Bournemouth, UK.

Corresponding Author:

Corresponding author: ¹Benard Ohene Botwe: Sirbenard13@gmail.com /bebotwe@ug.edu.gh
¹Department of Radiography, School of Biomedical and Allied Health Sciences, College of Health Sciences, University of Ghana, Box KB143, Korle Bu, Accra, Ghana. Tel: +233244029365

Abstract

Introduction: Vaccination is a key global strategy to mitigate the clinical impact of the COVID-19 virus. As part of local efforts to manage the outbreak, the government of Ghana announced its intention to vaccinate its population starting with essential and high-risk workers including radiographers. However, there were reports of hesitance to receiving the vaccine among the radiography workforce. This study was undertaken prior to the intended vaccination exercise to assess the willingness and concerns of radiographers to undergo the COVID-19 vaccination and to suggest recommendations to improve the vaccine uptake.

Method: An ethically-approved online survey strategy was employed for this cross-sectional study conducted between 24th–28th February 2021. The survey employed quantitative questions and open text response options. Quantitative and open text responses were analysed using statistical and thematic analyses, respectively.

Results: There were 108 responses (response rate of 46.3%). The majority ($n=64$, 59.3%) were willing to have the vaccine, however, some ($n=44$, 40.7%) were not. The main reason behind their willingness to have the vaccine was its ability to reduce the spread of infections and lower mortality ($n=35$, 54.7%). However, doubts about the vaccine's efficacy and side effects ($n=26$, 56.8%), conspiracy theory concerns about its effects on the Ghanaian race ($n=4$, 9.1%), and fertility concerns ($n=2$, 4.5%) were some reasons for their hesitance to receive the vaccine. The open text commentary further revealed that the vaccine was thought of as a lifesaving medication, however, clinical safety concerns, lack of education/information and religious beliefs were affecting peoples' willingness to be vaccinated.

Conclusion: Our findings demonstrate the need for an urgent public health educational intervention to address the COVID-19 vaccine hesitancy concerns raised by radiographers to help increase the vaccine uptake.

Implication for practice: The study provides pertinent information to improve COVID-19 vaccine uptake among radiographers to limit the spread of infections.

COVID-19 vaccine hesitancy concerns: Findings from a Ghana Clinical Radiography Workforce Survey

Abstract

Introduction: Vaccination is a key global strategy to mitigate the clinical impact of the COVID-19 virus. As part of local efforts to manage the outbreak, the government of Ghana announced its intention to vaccinate its population starting with essential and high-risk workers including radiographers. However, there were reports of hesitance to receiving the vaccine among the radiography workforce. This study was undertaken prior to the intended vaccination exercise to assess the willingness and concerns of radiographers to undergo the COVID-19 vaccination and to suggest recommendations to improve the vaccine uptake.

Method: An ethically-approved online survey strategy was employed for this cross-sectional study conducted between 24th–28th February 2021. The survey employed quantitative questions and open text response options. Quantitative and open text responses were analysed using statistical and thematic analyses, respectively.

Results: There were 108 responses (response rate of 46.3%). The majority ($n=64$, 59.3%) were willing to have the vaccine, however, some ($n=44$, 40.7%) were not. The main reason behind their willingness to have the vaccine was its ability to reduce the spread of infections and lower mortality ($n=35$, 54.7%). However, doubts about the vaccine's efficacy and side effects ($n=26$, 56.8%), conspiracy theory concerns about its effects on the Ghanaian race ($n=4$, 9.1%), and fertility concerns ($n=2$, 4.5%) were some reasons for their hesitance to receive the vaccine. The open text commentary further revealed that the vaccine was thought of as a lifesaving medication, however, clinical safety concerns, lack of education/information and religious beliefs were affecting peoples' willingness to be vaccinated.

Conclusion: Our findings demonstrate the need for an urgent public health educational intervention to address the COVID-19 vaccine hesitancy concerns raised by radiographers to help increase the vaccine uptake.

Implication for practice: The study provides pertinent information to improve COVID-19 vaccine uptake among radiographers to limit the spread of infections.

Keywords: COVID-19; vaccine hesitancy; Ghana; radiographer; Survey

Introduction

The first case of Coronavirus disease 2019 (COVID-19), caused by the severe acute respiratory syndrome Coronavirus 2 (SARS CoV-2)¹ was reported in December 2019.² The disease has since spread to 220 countries and territories across the globe.³ The first cases of COVID-19 in Ghana were recorded on 13th March, 2020.⁴ As at June, 2021, Ghana had recorded a total of 95,259 cases, with a total of 974 deaths.³ About the same time, a total of 3, 899, 172 deaths had been recorded worldwide out of 179, 686, 071 infections.³ Healthcare workers are amongst those at a high risk for infection with COVID-19. They are a potential source of virus transmission to patients, colleagues, family and friends.^{5,6} Studies from Spain and Ireland, all conducted in 2020 showed very high infection rates amongst healthcare workers, despite provision of adequate personal protective equipment [PPEs] and implementation of suppression strategies for community spread of the virus.^{7,8} In Ghana (as of May 2021), a total of 2079 health workers had been infected with 11 deaths.^{9,10}

COVID-19 predominantly affects the respiratory system, resulting in pneumonia.¹¹ Medical imaging is crucial in the early diagnosis, monitoring disease progression and severity of COVID-19.^{12,13} Chest computed tomography (CT), lung ultrasound and chest radiographs, performed by radiographers, were critical adjunctive tools in the early diagnosis and management of COVID-19.¹⁴ For example, in a Singapore hospital, chest radiography formed part of the triage protocol in the emergency department, designated screening areas and on the wards for monitoring symptoms (e.g., among patients presenting with acute respiratory symptoms) as well as non-COVID-19 related emergency conditions.^{15,16} This puts radiographers at particularly increased risk of exposure to the virus, because by the nature of their clinical practice they have physical contacts with both confirmed positive and suspected infected patients.¹⁷ The complexities for risk of infection among this workforce is further increased in situations where radiological examinations are to be conducted in places not originally designed for this purpose, e.g., in newly created isolation centres with limited PPEs.^{18,19} For example, Foley and colleagues reported that a high proportion of Irish radiographers (45%) were inadvertently exposed to COVID-19-positive patients without adequate PPE.¹⁷ A similar trend was reported during the Middle East Respiratory Syndrome (MERS) outbreak in Saudi Arabia to indicate that radiographers had the highest infection rate of 29.4% relative to other healthcare workers.²⁰

1
2
3
4 Vaccination has been shown to be effective in preventing infection and reducing mortality
5 of most infectious pathologies.²¹ There has been a rapid development, approval and emergency
6 use authorisation of several brands of COVID-19 vaccines and vaccination commenced by
7 December 2020 in several European countries.^{22,23} Vaccination is confirmed as the key global
8 strategy to protect and mitigate the clinical impact of the COVID-19 virus.²⁴ Meanwhile,
9 acceptability of vaccines is critical, because it underpins the success of any vaccination
10 programme.²⁵ According to the WHO, vaccine hesitancy (defined as the delay in acceptance,
11 reluctance, or refusal of vaccination despite the availability of vaccination services) is one of the
12 top ten threats to global health in 2019.²⁶ Factors influencing vaccine hesitancy are complex,
13 showing varying dynamics among different population groups and locations and to specific
14 vaccines.²⁵ Some of these factors have been identified to be trust in governmental policies and
15 communications, socio-cultural/socioeconomic status, media, educational status, perception of
16 individual and community risk, acquaintance with persons infected by COVID-19 and being a
17 health worker.^{25,26-29}

18
19
20
21
22
23
24
25
26
27
28
29
30 International population-wide studies provided predictions of up to 70% of vaccine
31 acceptability of current potential vaccines.²⁹ Health personnel are said to be a reliable source of
32 vaccine-related information for the public. They are better placed to explain to their patients'
33 concerns they may have about the importance and safety of vaccines.³⁰ Although the frontline
34 healthcare workforce is prioritised for vaccination against COVID-19,³¹ vaccine acceptance rates
35 and perceptions are diverse. For example, studies among Congolese healthcare workers and Israeli
36 doctors in March and April of 2020 reported acceptance rates of 27.7% and 78.1%, respectively.³²
37 Other studies conducted in Canada and Belgium as well as parts of USA in the last quarter of 2020
38 reported vaccine acceptance rates in the range of 36% to 72.4%.³³⁻³⁵ Of note, these studies were
39 conducted either before vaccines were developed or before the actual commencement of the
40 vaccination programmes.^{27,28} The lack of information about COVID-specific vaccines and their
41 potential side effects at the time of the studies have the potential of influencing people's
42 responses.²⁹

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2
3
4 willingness and concerns of radiographers to undergo the COVID-19 vaccination and to suggest
5 recommendations to improve the vaccine uptake among radiographers.
6
7
8
9

10 **Method**

11 *Study Design, Development and Distribution*

12
13
14 A cross-sectional research design employing an online survey strategy that uses
15 quantitative questions and some open text response options were utilised to obtain data. The
16 research instrument was a self-reported questionnaire (Appendix 1) developed by two of the
17 researchers (BOB and WKA). This approach was also useful because rapid data/information was
18 needed to develop strategies that would enhance the COVID-19 vaccination process and to
19 promptly address any concern that radiographers may have about the vaccination exercise.
20
21
22
23
24

25 The online survey was active between the 24th-28th February 2021 prior to the
26 commencement of the COVID-19 vaccination in Ghana on 1st March 2021. This gave all the
27 registered radiographers ($n=234$) with the professional council (Allied Health Professions Council,
28 AHPC) the opportunity to participate in study. To prevent physical contact with the respondents
29 while limiting the spread of COVID-19 infection during the study period, an online survey strategy
30 was employed. Respondents were given the opportunity to provide their general perspectives about
31 the COVID-19 vaccination as a sequel to the closed-ended questions to gain better understanding
32 and in-depth insight of the research problem. Prior to the data collection process, expert review
33 analysis was used to assess and approve the content and face validity of the questionnaire. This
34 process involved two academic reviewers with experience in the research area. They also assessed
35 the questionnaire functionality and presentation of the survey (across different types of devices,
36 operating system platforms etc) using a 2-point content rating scale (0=non important;
37 1=important) of which none of the questions was rejected. This process helped in evaluating
38 whether each of the measuring items match the given conceptual domain of the study objectives.³⁶
39
40 The questionnaire was also piloted among three radiographers and the result was satisfactory.
41 Moreover, a retest (second pilot) was conducted on the closed ended questions and a test-retest
42 statistical analysis yielded a kappa score of 0.80 to imply a strong reliability.
43
44
45
46
47
48
49
50
51
52
53
54
55

56 Google Forms (Google, Mountain View, CA) was used to host the research instrument
57 online along with the study information sheet and the associated link was shared and promoted on
58 various platforms of the Ghana Society of Radiographers. The platform allowed respondents to
59
60
61
62
63
64
65

1
2
3
4 give their consent before they could participate in the study. Each participants' response was coded
5
6 to ensure anonymity and were digitally stored on google drive with an encrypted password. Prior
7
8 to commencement, the study received ethical approval from the Ethics and Protocol Review
9
10 Committee of the School of Biomedical & Allied Health Science
11
12 (SBAHS/AA/RAD/10679769/2020-2021) and gatekeepers' permission was obtained from the
13
14 Ghana Society of Radiographers to engage its membership for this study.

15 16 17 **Data analysis**

18
19 At the end of the study, data was downloaded from Google Forms and prepared for
20
21 analyses. For the quantitative responses Microsoft Excel 2013 was used to generate graphs for
22
23 easy presentations of the results. Subsequently, the data were transferred into Statistical Package
24
25 for the Social Sciences (SPSS) version 23 (IBM, Armonk, NY) for inferential analysis. Spearman
26
27 correlation was used to assess the level of association between respondent's knowledge levels
28
29 about COVID-19 vaccines and age. Mann-Whitney U test was used to compare the knowledge
30
31 levels about COVID-19 vaccine among the gender groups. Furthermore, Wilcoxon signed-rank
32
33 test was used to assess statistical differences in knowledge levels between those who indicated
34
35 (un)/willingness to receive a COVID-19 vaccine. Since the intention to vaccinate or not (which is
36
37 the dependent variable) is a dichotomous data, multivariate binary logistic regression analysis was
38
39 used to determine the impact of the demographic variables on the intention to vaccinate or receive
40
41 the COVID-19 vaccine. Statistical significance level was set at $p < 0.005$. For the open text
42
43 commentary, thematic analysis was used to generate major themes as recommended.³⁷

44 45 **Results**

46 47 **1. Quantitative Results**

48
49 At the end of the study, 108 responses (representing a response rate of 46.1%) were received
50
51 from radiographers in Ghana about their views on the acceptance of COVID-19 vaccine. The
52
53 majority ($n=73$, 67.6%) of the respondents were males and a good number ($n=66$, 61.1%) had
54
55 worked for 10 or less years. Details of the demographic characteristics of the study respondents
56
57 are presented in **Table 1**. Almost all, except two (1.9%), were aware of intention of the government
58
59 of Ghana to embark on COVID-19 vaccination. When respondents were asked to self-describe
60
61
62
63
64
65

1
2
3
4 their knowledge levels (on a scale of 1-5, where 1 is poorly knowledgeable and 5 is highly
5 knowledgeable) about COVID-19 vaccines, the majority ($n=40$, 37.0%) indicated that they were
6 moderately (averagely) knowledgeable about them, only a few ($n=4$, 3.7%) described their
7 knowledge as poor (**Figure 1**). However, the knowledge levels of the respondents did not
8 significantly correlate with age ($rs=0.152$, $p=0.117$), neither did they differ among the gender
9 groups (*Mann Whitney U*, $p=0.253$). The results further showed that the respondents were aware
10 of many types of COVID-19 vaccines (**Table 2**). Only one (0.9%) did not know about any of the
11 COVID-19 vaccines.
12
13
14
15
16
17
18

19
20 When questioned on whether they will receive a COVID-19 vaccine, the majority ($n=64$,
21 59.3%) said yes, however, a significant number ($n=44$, 40.7%), said no. Respondents who
22 indicated their willingness to receive a COVID-19 vaccine revealed that the ability of the vaccine
23 to help curb the spread and reduce the number of cases and deaths ($n=35$, 54.7%) was their key
24 reason among others (**Table 3**). However, the main reason provided for not receiving a COVID-
25 19 vaccine was the fact that they were not convinced about its effectiveness, efficiency, and side
26 effects ($n=26$, 56.8%). Others mentioned the lack of adequate research evidence to back the
27 potency of the vaccines ($n=5$, 11.4%), limited evidence and concerns about its effects on the
28 African race and Ghanaian population ($n=4$, 9.1%), lack of adequate information and education
29 ($n=3$, 6.8%) about COVID-19 vaccines, lack of interest ($n=2$, 4.5%) and a very small number of
30 respondents appeared to be influenced by a range of conspiracy theories including the effects of
31 the vaccine on fertility ($n=2$, 4.5%). Further reasons are provided in **Table 4**. **Figure 2** shows the
32 key demographic characteristics of respondents who indicated their (un)/willingness to receive a
33 COVID-19 vaccine. Given the opportunity for them to prescribe interventions that could be used
34 to encourage them to have a COVID-19 vaccine, some said they would accept a COVID-19
35 vaccine if there were more than enough evidence to show that it provides immunity against the
36 infection and it's variants ($n=14$, 31.8%) and no adverse reactions happen to people of their race
37 after they have taken it for a while ($n=10$, 22.7%). Others also indicated that they would receive a
38 vaccine if it were proven beyond reasonable doubt that it is generally safe and would not affect
39 their spiritual wellbeing ($n=3$, 6.8%). However, some ($n=17$, 38%) indicated that nothing will
40 encourage them except God. Inferential statistics further showed that the knowledge levels about
41 COVID-19 vaccines between those who indicated their (un)/willingness to receive a vaccine were
42 not statistically significant (*Wilcoxon signed-rank*, $p=0.132$). Moreover, the intention to receive
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 the vaccine did not statistically associate with educational level (AOR=1.425; CI 95%: 0.596-
5 3.404; p=0.426), age (AOR=0.927; CI 95%: 0.388-2.212; p=0.864), gender (AOR=0.494; CI 95%:
6 0.214-1.141; p=0.099), years of practice (AOR= 0.942; CI 95%: 0.485-1.831; p=0.861) and work
7 setting (AOR=0.842; CI 95%:0.337-2.016; p=0.671).
8
9

10 11 12 13 14 15 **2. Open Text Results:** Perception about COVID-19 vaccination 16

17 The various themes developed from the open text responses on the perception of respondents
18 about COVID-19 vaccination were (i) lifesaving medication, (ii) safety doubts and concerns, (iii)
19 lack of education/information and (iv) religious beliefs. Further details are presented below and is
20 schematically presented in **Figure 3**.
21
22

23 24 25 **(i) Lifesaving medication.** 26

27 Some respondents perceived COVID-19 vaccines as lifesaving medication that would fight
28 against the COVID-19.
29

30 *“I think it can prevent the spread of the virus” (Rad 3)*

31 *“it’s been tested and proven to be effective against the pandemic” (Rad 56)*

32 *“It is a preventive way to control the spread and effect of the virus and reduce the death*
33 *rates” (Rad 63).*
34
35
36

37 38 **(ii) Safety doubts and concerns** 39

40 Several of the respondents also had some doubts and concerns about the safety of the vaccines.
41

42 *“Efficacy of the vaccines have not been adequately confirmed” (Rad 31).*

43 *“We should really consider the efficacy rate of the various vaccines taking into*
44 *consideration our race and setting and not because it's considered safe for other races or*
45 *settings” (Rad 22).*
46
47

48 *“I don’t fully trust it” (Rad 98).*
49

50 Again, some also held onto the opinion that since it is statistically evident that the Western world
51 were more affected by COVID-19 than African countries there was no reason why African leaders
52 should rush in getting its citizens vaccinated for a pandemic of which they suffer the least. Thus,
53 some did not trust the government’s intentions in the fight against COVID-19.
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 *“I don’t trust our leaders. Africans are less affected compared with the western world.*
5
6 *Why should the government go for a vaccine and expose the citizens to be used for*
7
8 *clinical trials for vaccination of a drug which has not been fully tried and tested. It has*
9
10 *only pass emergency approval” (Rad 40).*

11
12
13 **(iii) Lack of education**
14

15
16 A few of the respondents also felt that education about the vaccines was not enough to
17 help them make an informed decision on whether to receive the vaccines or not, this would help
18 to allay the fears of people. Others also felt that lack of proper education created a vacuum or the
19 platform for people to peddle falsehood information which has “polluted” people’s mind about
20 the vaccines.
21
22

23
24 *“I think there should be enough education about the vaccine to allay people's fears about*
25 *the vaccine” (Rad 50).*

26
27 *“I think it's a step in the right direction. It's unfortunate however that Nay Sayers and*
28 *Charlatans have been given the platform to pollute people’s minds” (Rad 17).*
29

30
31 **(iv) Religious beliefs**
32

33
34 Due to the religious belief, some respondents perceived that it was not necessary to accept
35 the vaccine because God would protect them from getting infected by the COVID-19 virus. One
36 respondent, in particular, indicated that:
37

38 *“I have God” (Rad 10).*
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1: Demographic characteristics of the respondents.

Demographic parameters	Frequency <i>n</i> (%)
Gender	
Male	73(67.6%)
Female	35(32.4%)
Age (yrs)	
20-29	55(50.9%)
30-39	35(32.4%)
40-49	16(14.8%)
50-59	2(1.9%)
≥60	0 (0%)
Years of practice	
< 10	66(61.1%)
10-20	35(32.4%)
>20	7(6.5%)
Sector of practice	
Government sector	97(73.1%)
Private sector	20(18.5%)
Quasi-government	9(8.3%)
Highest educational level	
Certificate/diploma qualification	5(4.6%)
Bachelor's degree (BSc)	81(75.0%)
Postgraduate certificate/diploma	4(3.7%)
Master's degree	15(13.9%)
Doctor of Philosophy	3(2.8%)

Key: yrs = years




Figure 1: Respondents self-described knowledge level about COVID-19 vaccines. (*Note: Moderate is average and Adequate is satisfactory*)

Table 2: Types of vaccines known by respondents.

Types of vaccines	Frequency(n)	Percentage (%)
Oxford-Astra Zeneca	64	59.3
Pfizer-BioNTech	71	65.7
Moderna	26	24.1
Sputnik V	29	26.9
Novavax (NVX-CoV2373)	16	14.8
Sinopharm	8	7.4
Sinovac	21	19.4
Johnson and Johnson's Janssen	2	1.9
I do not know	1	0.9

Table 3: Reasons why respondents will accept the COVID-19 vaccine ($n=64$).

Reason	Frequency	
	<i>n</i>	%
Vaccination will help curb the spread and reduce the number of cases and deaths.	35	54.7
It will help us to achieve herd immunity.	9	14.1
I trust the effectiveness, efficiency of the vaccine which comes with minimal side effects.	2	3.1
Taking of the vaccine is inevitable, it is the only main solution to the pandemic.	8	12.5
I have enough knowledge of it.	2	3.1
It has been scientifically tested and approved by the food and drugs authority (FDA) and WHO.	4	6.3
It is quite safe, and the benefits outweigh the risk.	3	4.7
I think it will work like other vaccines worked in the past.	1	1.6

Table 4: Reasons for not accepting COVID-19 vaccines ($n=44$).

Reason	Frequency	
	<i>n</i>	%
Not convinced about its effectiveness and side effects.	25	56.8
Not interested.	2	4.5
No evidence on effects on my type of race and population (Ghanaians).	4	9.1
Due to adverse reactions to previous vaccines eg. H1N1 vaccine in 2012.	1	2.3
Lack of adequate information and education.	3	6.8
Lack of enough research evidence to back the potency of the vaccines.	5	11.4
Not sure if it affects fertility.	2	4.5
I have heard of some severe side effects.	1	2.3
I do not know whether people with underlying conditions can take it.	1	2.3

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 2: Key demographic characteristics of respondents who indicated their unwillingness and willingness to have a COVID-19 vaccine.




Figure 3: Respondents’ perception of about COVID-19 vaccination

Discussion

Medical imaging is imperative in evidence-based choice of action in the diagnosis, management and treatment of COVID-19 patients.^{14,18,38-47} Despite the dreadful nature of the COVID-19 pandemic and post lock down effects in Ghana, it took some time for the government to secure vaccines for its people. In an attempt to combat COVID-19, radiographers were identified as part of the front-line healthcare experts who are at a high risk to be infected.^{17,19,37-39} This is due to the nature of the profession and the role medical imaging plays in the diagnosis and care management of COVID-19 patients.

The current study sought to identify the concerns and readiness of radiographers in Ghana to take the COVID-19 vaccination. Out of 234 registered radiographers in Ghana, the study had a response rate of 46.1%, which is lower than the values obtained in quite recent studies among radiographers in Ghana^{48,49} but is within a reasonable range to yield comparable results for similar inferences.⁵⁰ The awareness among the radiographers about the intent of the government to pursue COVID-19 vaccination was found to be encouraging which is attributed to the promotion of the vaccination process in the media. This was relatively higher than the COVID-19 vaccination

1
2
3
4 awareness level (40.8%) observed in a similar Ethiopian cohort⁵¹ and other countries such as in
5 Democratic Republic of Congo⁵² and Saudi Arabia.⁵³ Similarly, Shaw et al³⁰ received good
6 response to the uptake of the vaccination in the USA due to a high promotional approach of the
7 vaccination programme which created more awareness among the people. In contrast to a related
8 survey in Egypt, a greater number of respondents had inadequate information about the vaccination
9 programme.⁵⁴ Besides, the current study identified reasonable knowledge among respondents
10 about various COVID-19 vaccines. It was also encouraging that the majority of respondents in the
11 current study agreed to be vaccinated with several reasons including the capability of the COVID-
12 19 vaccine to limit the spread of infections and reduce the number of deaths.

13
14
15
16
17
18
19
20
21 Notwithstanding, appreciable number of radiographers, who were youthful and self-
22 reported their COVID-19 vaccine knowledge levels as mainly moderate/average (38.6%) and
23 adequate/satisfactory (27.3%) (**Figure 1**), were against the vaccination and gave noteworthy
24 explanations. In particular, a few believed that they have God to protect them and so will not
25 vaccinate, while others wanted to wait till when no adverse reaction happens to people of their
26 race or population after they have taken it for a while. This was found to be interesting as one
27 would have expected that as health professionals, they would rather take the vaccine to encourage
28 family members to vaccinate. This attitude could be ascribed to misinformation (conspiracy
29 theories) on social media regarding the vaccines and previous knowledge of vaccine risks which
30 some respondents might have been exposed to. Again, some also held onto the opinion that since
31 it is statistically evident that the western world is more affected by COVID-19 than African
32 countries there was no reason why African leaders should rush in getting its citizens vaccinated
33 for a pandemic of which they suffer the least. Thus, some did not trust the government's intentions
34 in the fight against COVID-19 with the vaccination exercise. A study in Italy had respondents
35 showing similar misconceptions about the vaccine which led to hesitancy to being vaccinated.⁵⁵
36 Moreover, a reasonable number of the respondents questioned the intentions of the vaccination
37 and indicated that unless it is established beyond reasonable uncertainty that the COVID-19
38 vaccine is generally safe and would not affect their spiritual well-being, they will not undergo the
39 vaccination; a belief in illogical forces which appears to be endemic among some Ghanaian
40 radiographers.⁵⁶ Vaccine safety has been a concern among several healthcare workers which was
41 demonstrated in the current study. A few of the respondents cited the known adverse reactions
42 experienced with previous vaccines such as the H1N1 vaccine in 2012. These findings support
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 previous conclusions about healthcare workers anxieties regarding the safety of vaccines.⁵⁷ The
5 unwillingness to be acquiescent to COVID-19 vaccine was as a result of suspicions about the
6 efficacy and besides, the side effects of the vaccine, as was the case in the findings of Dini et al.⁵⁵
7
8 Some respondents were also particularly concerned about the vaccine's tolerability among the
9 Ghanaian race. This was not surprising as different racial/ethnic depictions in COVID-19 vaccine
10 trials is imperative because drugs and vaccines can diversely affect groups reflecting disparity in
11 underlying experiences and environmental exposure.⁵⁸ Others felt there were not enough evidence
12 to establish the potency of the vaccines. Although COVID-19 vaccines are highly effective,
13 particularly against hospitalisation and severe disease, no vaccine is 100% protective.^{59,60} Based
14 on some of these misgivings about the vaccines in general, some respondents reported that nothing
15 will inspire them to take the vaccine apart from God. This is not surprising as Ghana's population
16 is made up of about 71% Christians and 17% Muslims.⁶¹ Therefore, there would be a need for
17 religious leaders to play a critical role in dispelling the myths for the success of the vaccination
18 programmes. This is important because vaccination of healthcare personnel is a critical aspect of
19 occupational medicine strategies and infection control programmes within the healthcare
20 services.⁶²

21
22
23
24
25
26
27
28
29
30
31
32
33
34 The survey further revealed that the awareness level about COVID-19 vaccines between
35 those who showed their inclination to receive it and those otherwise, was not statistically
36 significant. This implies that the unwillingness to partake in the vaccination was not because of
37 lack of awareness. Moreover, the performed statistical tests demonstrated that the self-reported
38 knowledge levels of the respondents did not significantly correlate with age, neither did they differ
39 among the gender groups. This is contrary to the findings of Agyekum and colleagues⁶³ which
40 demonstrated that certain demographic parameters impact on the intention to vaccinate. The
41 findings of the current study demonstrated that the demographic variables of the radiographers
42 were not a determinant of an intention to vaccinate, and may further suggest that the positions
43 taken by the radiographers could be due to their self-conviction about the vaccine.

44
45
46
47
48
49
50
51
52 Since 1st March 2021, when the vaccination programme (Oxford–AstraZeneca COVID-
53 19 vaccines) started in Ghana, only 1.3% of the entire Ghanaian population (30+ million) had been
54 fully vaccinated and 2.8% had received a single dose as of July 2021.^{64,65} Even though this study
55 could not obtain data on the total number of radiographers who have been vaccinated so far, it is
56 anticipated that an innovative strategy would be needed to address the concerns raised by
57
58
59
60
61
62
63
64
65

1
2
3
4 participants of the current study to improve the vaccine acceptability among radiographers in
5
6 Ghana. In tackling these concerns, public health education would be important. Particularly, there
7
8 should be more proof to dispel the conspiracy theories being spread about the COVID-19
9
10 vaccination. As a proactive method to support the vaccination, this work has already led to the
11
12 initiation of educational strategies aimed at encouraging vaccine uptake by radiographers.
13

14 15 **Conclusion**

16
17
18 The study showed that a large proportion of the Ghanaian radiography workforce were
19
20 willing to receive the vaccine, however, a significant number were not. The observed results call
21
22 for an urgent public health educational intervention from stakeholders to promptly address the
23
24 COVID-19 vaccine hesitancy concerns raised by the radiographers to help increase vaccine uptake.
25
26 In tackling these concerns, more education about the COVID-19 vaccines is needed to encourage
27
28 radiographers to develop interest in accepting them. Particularly, when offering the education
29
30 (through e.g. specialised CPD, seminars, conferences, talk shows etc), there should be more proof
31
32 of the efficacy of the various vaccines in helping to limit the spread of infections of the various
33
34 variants. Furthermore, there is a need for more information and evidence to be shared on the safety
35
36 of the vaccines, particularly, among their race and tribes (e.g. among the Ghanaian population).
37
38 This safety education must also try to allay the fears of those who have reacted adversely to other
39
40 vaccines previously, those who think the vaccination is an attempt to make people infertile, and
41
42 those with underlying medical conditions but were unsure they could take the vaccines. There
43
44 should also be more than enough evidence to suggest that the vaccination exercise is generally
45
46 safe. Moreover, education to dispel myths and conspiracy theories about the COVID-19 vaccine
47
48 is highly needed.
49

50 **Conflict of interests**

51
52 None
53
54
55
56
57
58
59
60
61
62
63
64
65

References

1. World Health Organization. Naming the coronavirus disease (COVID-19) and the virus that causes it. Available from: [https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-\(covid-2019\)-and-the-virus-that-causes-it](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it)
2. Cucinotta D, Vanelli M. WHO declares COVID-19 a pandemic. *Acta Bio-Medica Atenei Parm*, 91 (1) (2020 19): 157-160
3. Worldometers.info [Internet]. Reported Cases and Deaths by Country or Territory. Available from: <https://www.worldometers.info/coronavirus/>
4. COVID-19 Updates /Ghana. First confirmed COVID-19 cases in Ghana 12 Mar 2020 (ghanahelthservice.org). Accessed May 08, 2021.
5. Wang J, Zhou M, Liu F. Reasons for healthcare workers becoming infected with novel coronavirus disease 2019 (COVID-19) in China. *J Hosp Infect*. 2020 May;105(1):100-101. doi: 10.1016/j.jhin.2020.03.002.
6. Asad H, Johnston C, Blyth I, Holborow A, Bone A, Porter L, et al. Health care workers and patients as Trojan horses: a COVID19 ward outbreak. *Infect Prevent Pract*. 2020;2:100073. <https://doi.org/10.1016/j.infpip.2020.100073>.
7. Suárez-García I, Martínez de Aramayona López MJ, Sáez Vicente A, Lobo Abascal P. SARS-CoV-2 infection among healthcare workers in a hospital in Madrid, Spain. *J Hosp Infect*. 2020;106(2):357-363. doi:10.1016/j.jhin.2020.07.020
8. Government of Ireland (2020) Health (Preservation and Protection and other Emergency Measures in the Public Interest) Act 2020. Irish Statute Book, Dublin Available at <http://www.irishstatutebook.ie/eli/2020/act/1/enacted/en/html>
9. MyJoyOnline, *789 Nurses and Midwives Test Positive for Covid-19 in 2021 – General Secretary, GRNMA*, Available from: https://www.myjoyonline.com/789-nurses-and-midwives-test-positive-for-covid-19-in-2021-general-secretary-grnma/?utm_source=rss&utm_medium=rss&utm_campaign=789-nurses-and-midwives-test-positive-for-covid-19-in-2021-general-secretary-grnma.
10. Kwofi M, *COVID-19 Hits 450 Doctors, kills 7*, Available from: <https://www.graphic.com.gh/news/general-news/covid-19-hits-450-doctors-kills-7.html>.
11. Cellina M, Orsi M, Oliva G. How to Reorganize the Radiology Departments to Face the 2019 Coronavirus Disease Outbreak. *Disaster Med Public Health Prep*. 2020;14(6):789-791. doi:10.1017/dmp.2020.159
12. Kooraki S, Hosseiny M, Myers L, Gholamrezanezhad A. Coronavirus (COVID-19) outbreak: what the department of radiology should know. *J Am Coll Radiol*, 2020; 17 (4): 447-451. doi: 10.1016/j.jacr.2020.02.008.
13. Xu X, Yu C, Qu J, Zhang L, Jiang S, Huang D, et al. Imaging and clinical features of patients with 2019 novel coronavirus SARS-CoV-2. *Eur J Nucl Med Mol Imaging* **47**, 1275–1280 (2020). <https://doi.org/10.1007/s00259-020-04735-9N>.
14. Stogiannos N, Fotopoulos D, Woznitza N, Malamateniou C. COVID-19 in the radiology department: What radiographers need to know. *Radiography (Lond)*. 2020 Aug;26(3):254-263. doi: 10.1016/j.radi.2020.05.012.
15. Chia A, Cheng LT, Wijaya L, Png MA, Sim WY, Hong WL, Chen RC. Chest radiographs and CTs in the era of COVID-19: indications, operational safety

- 1
2
3
4 considerations and alternative imaging practices. *Acad Radiol.* 2020;27(9):1193–1203.
5 doi:10.1016/j.acra.2020.06.022.
- 6
7 16. Kooraki S, Hosseiny M, Myers L, Gholamrezanezhad A (2020) Coronavirus (COVID-
8 19) Outbreak: What the Department of Radiology Should Know. *J Am Coll Radiol*
9 17(4):447–451
- 10 17. Foley, S.J., O’Loughlin, A. & Creedon, J. Early experiences of radiographers in Ireland
11 during the COVID-19 crisis. *Insights Imaging* **11**, 104 (2020).
12 <https://doi.org/10.1186/s13244-020-00910-6>
- 13
14 18. Pereira JM, Silva C, Freitas D, Salgado A. Burnout among Portuguese radiographers
15 during the COVID-19 pandemic. *Radiography (Lond)*. 2021 May 18:S1078-
16 8174(21)00054-7. doi: 10.1016/j.radi.2021.05.001.
- 17
18 19. Lewis S, Mulla F. Diagnostic radiographers' experience of COVID-19, Gauteng South
19 Africa. *Radiography (Lond)*. 2021 May;27(2):346-351. doi: 10.1016/j.radi.2020.09.009.
- 20
21 20. Alraddadi BM, Al-Salmi HS, Jacobs-Slifka K, et al. Risk factors for Middle East
22 respiratory syndrome coronavirus infection among healthcare personnel. *Emerg Infect*
23 *Dis* 2016; 22:1915–1920
- 24
25 21. Osterholm, N.S.; Kelley, A.; Sommer, E.A.; Belongia, E.A. Efficacy and effectiveness of
26 influenza vaccines: A systematic review and meta-analysis. *Lancet Infect. Dis.* **2012**, *12*,
27 36–44.
- 28
29 22. European Centre for Disease Prevention and Control (ECDC). First COVID-19 Vaccine
30 Authorized for Use in the European Union. Stockholm. Available
31 online: [https://www.ecdc.europa.eu/en/news-events/first-covid-19-vaccine-
32 authorised-use-european-union](https://www.ecdc.europa.eu/en/news-events/first-covid-19-vaccine-authorized-use-european-union).
- 33
34 23. EMA Recommends COVID-19 Vaccine Moderna for Authorisation in the EU. Available
35 online: [https://www.ema.europa.eu/en/news/ema-recommends-covid-19-vaccine-
36 moderna-authorisation-eu](https://www.ema.europa.eu/en/news/ema-recommends-covid-19-vaccine-moderna-authorisation-eu)
- 37
38 24. Anderson RM, Vegvari C, Truscott J, Collyer BS. Challenges in creating herd immunity
39 to SARS-CoV-2 infection by mass vaccination. *Lancet*. 2020 Nov 21;396(10263):1614-
40 1616. doi: 10.1016/S0140-6736(20)32318-7.
- 41
42 25. MacDonald, N.E.; Eskola, J.; Liang, X.; Chaudhuri, M.; Dube, E.; Gellin, B. Vaccine
43 hesitancy: Definition, scope and determinants. *Vaccine* **2015**, *33*, 4161–4164.
- 44
45 26. World Health Organization (WHO). Ten Threats to Global Health in 2019. 2019.
46 Available online: [https://www.who.int/news-room/spotlight/ten-threats-to-global-
47 health-in-2019](https://www.who.int/news-room/spotlight/ten-threats-to-global-health-in-2019)
- 48
49 27. Harapan H, Wagner AL, Yufika A, Winardi W, Anwar S, Gan AK, Setiawan AM,
50 Rajamoorthy Y, Sofyan H, Mudatsir M. Acceptance of a COVID-19 Vaccine in
51 Southeast Asia: A Cross-Sectional Study in Indonesia. *Front Public Health*. 2020 Jul
52 14;8:381. doi: 10.3389/fpubh.2020.00381.
- 53
54 28. Reiter PL, Pennell ML, Katz ML. Acceptability of a COVID-19 vaccine among adults in
55 the United States: How many people would get vaccinated? *Vaccine*. 2020 Sep
56 29;38(42):6500-6507. doi: 10.1016/j.vaccine.2020.08.043.
- 57
58 29. Lazarus JV, Ratzan SC, Palayew A, Gostin LO, Larson HJ, Rabin K, Kimball S, El-
59 Mohandes A. A global survey of potential acceptance of a COVID-19 vaccine. *Nat Med*.
60 2021 Feb;27(2):225-228. doi: 10.1038/s41591-020-1124-9.
- 61
62 30. Shaw J, Stewart T, Anderson KB, Hanley S, Thomas SJ, Salmon DA, Morley C.
63 Assessment of U.S. health care personnel (HCP) attitudes towards COVID-19
64
65

- vaccination in a large university health care system. Clin Infect Dis. 2021 Jan 25:ciab054. doi: 10.1093/cid/ciab054.
31. WHO. Coronavirus Disease (COVID-19): Vaccine Access and Allocation. Available online: [https://www.who.int/news-room/q-a-detail/coronavirus-disease-\(covid-19\)-vaccine-access-and-allocation](https://www.who.int/news-room/q-a-detail/coronavirus-disease-(covid-19)-vaccine-access-and-allocation).
 32. Sallam M. COVID-19 vaccine hesitancy worldwide: a concise systematic review of vaccine acceptance rates. Vaccines (Basel). 2021. <https://doi.org/10.3390/vaccines9020160>.
 33. Verger P, Scronias D, Dauby N, Adedzi KA, Gobert C, Bergeat M, Gagneur A, Dubé E. Attitudes of healthcare workers towards COVID-19 vaccination: a survey in France and French-speaking parts of Belgium and Canada, 2020. Euro Surveill. 2021 Jan;26(3):2002047. doi: 10.2807/1560-7917.ES.2021.26.3.2002047.
 34. Shekhar R, Sheikh AB, Upadhyay S, Singh M, Kottewar S, Mir H, Barrett E, Pal S. COVID-19 Vaccine Acceptance among Health Care Workers in the United States. Vaccines (Basel). 2021 Feb 3;9(2):119. doi: 10.3390/vaccines9020119.
 35. Gadoth A, Halbros M, Martin-Blais R, Gray A, Tobin NH, Ferbas KG, Aldrovandi GM, Rimoin AW. Cross-sectional Assessment of COVID-19 Vaccine Acceptance Among Health Care Workers in Los Angeles. Ann Intern Med. 2021 Jun;174(6):882-885. doi: 10.7326/M20-7580.
 36. Bolarinwa OA. Principles and methods of validity and reliability testing of questionnaires used in social and health science researches. Niger Postgrad Med J. 2015 Oct-Dec;22(4):195-201. doi: 10.4103/1117-1936.173959.
 37. Hsieh HF and Shannon SE. Three Approaches to Qualitative Content Analysis Qualitative Health Research, 2005; 15(9): 1277- 1288. DOI: 10.1177/1049732305276687
 38. Akudjedu TN, Mishio NA, Elshami W, Culp MP, Lawal O, Botwe BO, Wuni AR, Julka-Anderson N, Shanahan M, Totman JJ, Franklin JM. The global impact of the COVID-19 pandemic on clinical radiography practice: A systematic literature review and recommendations for future services planning. Radiography (Lond). 2021 Jul 12:S1078-8174(21)00084-5. doi: 10.1016/j.radi.2021.07.004.
 39. Akudjedu TN, Botwe BO, Wuni AR, Mishio NA. Impact of the COVID-19 pandemic on clinical radiography practice in low resource settings: The Ghanaian radiographers' perspective. Radiography (Lond). 2021 May;27(2):443-452. doi: 10.1016/j.radi.2020.10.013.
 40. Yasin B, Barlow N, Milner R. The impact of the Covid-19 pandemic on the mental health and work morale of radiographers within a conventional X-ray department. Radiography (Lond). 2021 May 11:S1078-8174(21)00047-X. doi: 10.1016/j.radi.2021.04.008.
 41. Elshami W, Akudjedu TN, Abuzaid M, et al. The radiology workforce's response to the COVID-19 pandemic in the Middle East, North Africa and India. *Radiography (Lond)*. 2021;27(2):360-368. doi:10.1016/j.radi.2020.09.016
 42. Akudjedu TN, Lawal O, Sharma M, Elliott J, Stewart S, Gilleece T, McFadden S, Franklin JM. Impact of the COVID-19 pandemic on radiography practice: findings from a UK radiography workforce survey. BJR Open. 2020 Sep 2;2(1):20200023. doi: 10.1259/bjro.20200023.
 43. England A, Littler E, Romani S, Cosson P. Modifications to mobile chest radiography technique during the COVID-19 pandemic - implications of X-raying through side room

- 1
2
3
4 windows. *Radiography (Lond)*. 2021 Feb;27(1):193-199. doi:
5 10.1016/j.radi.2020.07.015.
6
7 44. Zanardo M, Martini C, Monti CB, Cattaneo F, Ciaralli C, Cornacchione P, et al. .
8 Management of patients with suspected or confirmed COVID-19, in the radiology
9 department. *Radiography* 2020; 26: 264–8. doi: 10.1016/j.radi.2020.04.010
10 - [DOI](#) - [PMC](#) - [PubMed](#)
11
12 45. Sim WY, Chen RC, Aw LP, Abu Bakar R, Tan CC, Heng AL, Ooi CC. How to safely
13 and sustainably reorganise a large general radiography service facing the COVID-19
14 pandemic. *Radiography (Lond)*. 2020 Nov;26(4):e303-e311. doi:
15 10.1016/j.radi.2020.05.001.
16
17 46. Ruiz C, Llopis D, Roman A, Alfayate E, Herrera-Peco I. Spanish radiographers' concerns
18 about the COVID-19 pandemic. *Radiography (Lond)*. 2021 May;27(2):414-418. doi:
19 10.1016/j.radi.2020.10.001.
20
21 47. Shanahan MC, Akudjedu TN. Australian radiographers' and radiation therapists'
22 experiences during the COVID-19 pandemic. *J Med Radiat Sci*. 2021 Jun;68(2):111-120.
23 doi: 10.1002/jmrs.462.
24
25 48. Botwe BO, Antwi WK, Arkoh S, Akudjedu TN. Radiographers' perspectives on the
26 emerging integration of artificial intelligence into diagnostic imaging: The Ghana study. *J*
27 *Med Radiat Sci*. 2021 Feb 14. doi: 10.1002/jmrs.460.
28
29 49. Ashong GGNA, Rogers H, Botwe BO, Anim-Sampong S. Effects of occupational stress
30 and coping mechanisms adopted by radiographers in Ghana. *Radiography*, 2016, 22(2),
31 pp. 112–117. <https://doi.org/10.1016/j.radi.2015.09.002>
32
33 50. Morton BMS, Bandara DK, Carr AEP. In the 21st Century, what is an acceptable
34 response rate? *Australian And New Zealand Journal Of Public Health* 2012, 36(2): 106-
35 108 doi: 10.1111/j.1753-6405.2012.00854.x
36
37 51. Zewude B and Habtegiorgis T. Willingness to Take COVID-19 Vaccine Among People
38 Most at Risk of Exposure in Southern Ethiopia. *Pragmat Obs Res*. 2021 May 27;12:37-47.
39 doi: 10.2147/POR.S313991).
40
41 52. Nzaji KM, Ngombe KL, Mwamba NG, et al. Acceptability of Vaccination against Covid-
42 19 among healthcare workers in the Democratic Republic of the Congo. *Pragmatic*
43 *Observational Res*. 2020;11:103–109. doi:10.2147/POR.S271096
44
45 53. Qattan AMN, Alshareef N, Alsharqi O, Al Rahahleh N, Chirwa GC, Al-Hanawi MK.
46 Acceptability of a COVID-19 Vaccine Among Healthcare Workers in the Kingdom of
47 Saudi Arabia. *Front Med*. 2021;8:644300. doi:10.3389/fmed.2021.644300
48
49 54. Fares S, Elmnyer MM, Mohamed SS, Elsayed R. COVID-19 Vaccination Perception and
50 Attitude among Healthcare Workers in Egypt. *Journal of Primary Care & Community*
51 *Health*, 2021, (12): 1–9
52
53 55. Gerussi V, Peghin M, Palese A, et al. Vaccine Hesitancy among Italian Patients
54 Recovered from COVID-19 Infection towards Influenza and Sars-Cov-2 Vaccination.
55 *Vaccines*. 2021;9(2):172. doi:10.3390/vaccines9020172
56
57 56. Antwi WK, Reeves P, Ferris C, Aziato L. Behavioural beliefs of Ghanaian radiographers
58 and reporting of child physical abuse. *Radiography (Lond)*. 2019 Feb;25(1):51-57. doi:
59 10.1016/j.radi.2018.10.002.
60
61 57. Dini G, Toletone A, Sticchi L, Orsi A , Bragazzi NL & Durando P. Influenza vaccination
62 in healthcare workers: a comprehensive critical appraisal of the literature, (2017):
63 *Human Vaccines & Immunotherapeutics*, DOI: 10.1080/21645515.2017.1348442
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
58. Lopez L, Hart LH, Katz MH. Racial and Ethnic Health Disparities Related to COVID-19. *JAMA*. 2021;325(8):719–720. doi:10.1001/jama.2020.26443
 59. WHO. Coronavirus disease (COVID-19): Vaccines safety. [https://www.who.int/news-room/q-a-detail/coronavirus-disease-\(covid-19\)-vaccines-safety](https://www.who.int/news-room/q-a-detail/coronavirus-disease-(covid-19)-vaccines-safety) Accessed 22/06/21
 60. Yiwen K, Hegney D, Drury V. A comprehensive systematic review of healthcare workers' perceptions of risk from exposure to emerging acute respiratory infectious diseases and the perceived effectiveness of strategies used to facilitate healthy coping in acute hospital and community healthcare settings. *JBI Libr Syst Rev*. 2010;8(23):917-971.
 61. Ghana statistical Service. Ghana population 2021. Worldpopulationreview.com/countries/ghana-population Accessed 11/08/2021
 62. Maltezou, HC, Theodoridou K, Ledda C, Rapisarda V, and Theodoridou M. (2018). Vaccination of healthcare workers: is mandatory vaccination needed? Expert Review of Vaccines. doi:10.1080/14760584.2019.1552141
 63. Agyekum Aw, Afrifa-Anane GF, Kyei-Arthur F, Addo B. Research Article Acceptability of COVID-19 Vaccination among Health Care Workers in Ghana. *Advances in Public Health* 2021;1-18. <https://doi.org/10.1155/2021/9998176>
 64. Statista. Cumulative number of COVID-19 vaccine doses administered in Ghana 2021. <https://www.statista.com/statistics/1227780/total-number-of-covid-19-vaccination-doses-in-ghana/> Accessed on 27/06/2021
 65. Google COVID-19 Vaccination update. COVID-19 Vaccine Ghana. <https://www.google.com/search?q=covid-19+vaccination+in+Ghana&rlz>. Accessed 27/06/21.

1
2
3
4 **Appendix 1: Questionnaire**

5 **COVID-19 vaccine uptake and insight by radiographers: A study in Ghana**

6
7 * Required

8 If you consent to participate in this study, please tick the box below.

9 *Check all that apply.*

10 I consent to participate in this study.

11
12
13 **DEMOGRAPHICS**

14 1. Gender

15 *Mark only one oval.*

16 Male

17
18 Female

19
20
21 2. Kindly indicate your age range * *Mark only one oval.*

22
23 20- 29 years

24 30-39 years

25 40-49 years

26 50-59 years

27 60 years and above

28
29
30
31 3. Years of practice * *Mark only one oval.*

32 Less than 10 years

33 10- 20 years

34 21 years and above

35
36
37 4. Work setting *

38 *Mark only one oval.*

39 Private sector

40 Government sector

41 Quasi- government sector

42
43
44
45 5. Highest educational level

46 **Mark only one oval.*

47 Certificate/Diploma

48 Bachelor of Science (BSc)

49 Postgraduate Certificate/Diploma

50 Master's Degree (MSc)

51 Doctor of Philosophy

52 Other please specify.....

53
54
55
56
57
58 **AWARENESS OF COVID-19 VACCINES**

59 6. Have you heard of any Covid -19 vaccine?

1
2
3
4 *Mark only one oval.

- 5 Yes
6 No
7

8
9 7. On a scale of 1-5 (where 1 is poorly knowledgeable and 5 is highly knowledgeable), how
10 much knowledge do you have about COVID-19 vaccine?
11

12 *Mark only one oval. (Note: Moderate is average and Adequate is satisfactory)

- 13
14 1- Poorly knowledgeable
15 2- Fairly knowledgeable
16 3- Moderately knowledgeable
17 4- Adequately knowledgeable
18 5- Highly knowledgeable
19
20

21 8. Which vaccine are you aware of being used globally?
22

23 Check all that apply.

- 24
25 Pfizer-BioNTech
26 Moderna
27 Sinovac
28 Sinopharm
29 Oxford-Astra Zeneca
30 Sputnik V
31 Novavax
32

33 Other: _____
34
35
36
37

38
39
40 **WILLINGNESS AND UNWILLINGNESS TO RECEIVE COVID-19 VACCINES**

41 9. Would you receive any of the COVID-19 vaccines if provided?
42

43 *Mark only one oval.

- 44 Yes
45
46 No
47

48 10. Kindly provide a reason for your answer to question 9.
49

50
51
52

53 11. If you would not receive any of the COVID-19 vaccines, please indicate what can be done
54 to get you to be vaccinated.
55

56
57
58

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

GENERAL PERCEPTION ABOUT COVID-19 VACCINATION

12. What is your general view about the COVID-19 vaccination?

.....
.....

Acknowledgement

We would like to thank the leadership of the Ghana Society of Radiographers as well as the members for supporting and participating in this study.