



## Review article

## Genomic instability and the link to infertility: A focus on microsatellites and genomic instability syndromes

Jack Wieland<sup>a,\*</sup>, Sarah Buchan<sup>a</sup>, Sioban Sen Gupta<sup>b</sup>, Anna Mantzouratou<sup>a</sup><sup>a</sup> Department of Life and Environmental Sciences, Faculty of Science and Technology, Bournemouth University, Poole BH12 5BB, UK<sup>b</sup> Institute for Women's Health, 86-96 Chenies Mews, University College London, London WC1E 6HX, UK

## ARTICLE INFO

## Keywords:

Infertility

Microsatellite instability

Recurrent miscarriage

Genomic instability syndrome

Mismatch repair

## ABSTRACT

Infertility is associated to multiple types of different genomic instabilities and is a genetic feature of genomic instability syndromes. While the mismatch repair machinery contributes to the maintenance of genome integrity, surprisingly its potential role in infertility is overlooked. Defects in mismatch repair mechanisms contribute to microsatellite instability and genomic instability syndromes, due to the inability to repair newly replicated DNA. This article reviews the literature to date to elucidate the contribution of microsatellite instability to genomic instability syndromes and infertility. The key findings presented reveal microsatellite instability is poorly researched in genomic instability syndromes and infertility.

## Introduction

Infertility is defined as either primary or secondary infertility. Primary infertility refers to the failure of pregnancy after one year without the use of birth control methods [1], whilst secondary infertility refers to the failure of conceiving another child after the first [2]. Furthermore, recurrent miscarriage (RM), is defined as three or more consecutive miscarriages [3–4]. Infertility is reported to affect approximately 15% of couples globally [5]. Genetic analyses has provided strong evidence for the role of gene variants and chromosomal aberrations [6–14]. These data suggest that individuals harbouring factors that predispose them to genomic instability have decreased fertility rates.

Genomic instability can manifest as multiple types of aberrations within the human genome. These aberrations are evident at the chromosomal or gene level as aneuploidy [15], translocations [16], inversions [17], deletions [18]. Genomic instability is most pronounced in patients with genomic instability syndromes due to variant(s) of gene(s) responsible for DNA repair. Genomic instability and genomic instability syndromes may be accompanied with microsatellite instability (MSI) [19–20,10,21].

## Microsatellites

Microsatellites, the “genetic fingerprints” of the human genome, contains variable numbers of repeated 1–6 base pair units (bp), encompass approximately 3% of the human genome [22]. Generally, microsatellites are located within the intron regions (Fig. 1) of a gene [23], the 5' and 3' untranslated regions (5'UTR and 3'UTR) [24], intergenic spaces [25], and present in exon regions [26]. Additionally, for humans specifically, microsatellites can be observed adjacent to transcriptional start sites [27], translation initiation sites [28], and around CpG DNA methylation sites [27]. High levels of variability between individuals at microsatellite sites lends microsatellites to different applications including DNA profiling [29–31], and studies of disease [32–34]. Also, these may not be actively conserved in the human genome, as microsatellites are in regions that are not vital to gene function and hence, they are prone to variation (Fig. 1). DNA replication slippage may also be one of the factors involved in MSI occurrence [35–37].

The hypermutable state of microsatellites is partially due to the mismatch repair (MMR) system, correcting DNA sequences by acting as a “proof-reader” (Fig. 2) if there is a sequence error. The other

**Abbreviations:** microsatellite instability, (MSI); recurrent miscarriage, (RM); primordial germ cells, (PGCs); double stranded breaks, (DSBs); Bloom syndrome, (BS); mosaic variegated aneuploidy syndrome, (MVA); Lynch syndrome, (LS); colorectal cancer, (CRC); mouse embryo fibroblasts, (MEFs); mismatch repair, (MMR); sister chromatid exchange, (SCE); idiopathic recurrent miscarriage, (IRM); preimplantation genetic testing, (PGT).

\* Corresponding author.

E-mail addresses: [jwieland2@bournemouth.ac.uk](mailto:jwieland2@bournemouth.ac.uk), [jack.wieland@outlook.com](mailto:jack.wieland@outlook.com) (J. Wieland), [sbuchan@bournemouth.ac.uk](mailto:sbuchan@bournemouth.ac.uk) (S. Buchan), [sioban.sengupta@ucl.ac.uk](mailto:sioban.sengupta@ucl.ac.uk) (S. Sen Gupta), [amantzouratou@bournemouth.ac.uk](mailto:amantzouratou@bournemouth.ac.uk) (A. Mantzouratou).

<https://doi.org/10.1016/j.ejogrb.2022.06.001>

Received 21 March 2022; Received in revised form 25 May 2022; Accepted 1 June 2022

Available online 3 June 2022

0301-2115/© 2022 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

contributor is the simple nature of a microsatellite sequence as discussed above. DNA errors recognised by the MMR pathway include insertions [46], deletions [43], and mis-incorporations [47] of a nucleotide base. Defects within the MMR pathway is to MSI. MSI in turn is associated with several diseases and syndromes including cancers [48–51], inflammatory diseases including Crohn's disease [32,52] and Behçet's syndrome [33]. It should also be acknowledged that “hereditary non-polyposis colorectal cancer (HNPCC) or Lynch syndrome (LS) is the most common disorder with MSI” [53–60], yet the role of MSI in infertility is poorly understood.

Research to date evidenced links between genomic instability syndromes and gene variants and infertility [61–74]. In contrast, the specific link between MSI and infertility/RM, independent of genomic instability syndromes, is poorly understood. Therefore, we aim to highlight upon the importance of MSI in infertility by reviewing the literature on 1) the role of MSI during gametogenesis, early embryogenesis, and implantation, 2) MSI in genomic instability syndromes on infertility and 3), the known impact of specific MSI sites on infertility independent of genomic instability syndromes. We show that MSI contributes to infertility in genomic instability syndromes and independent of genomic instability syndromes. However, data is limited and more research in this area is warranted.

### Microsatellite instability in germ cell development

MSI may disrupt normal foetal development through different mechanisms including prior to conception during gametogenesis, through influencing germ cell mosaicism and during early embryogenesis and implantation. We describe the role of MSI in each of these three stages in turn below.

### Gametogenesis and microsatellite instability


Gametogenesis is the process of diploid or haploid cells entering cellular division and undergoing differentiation to mature haploid gametes. In male's, primordial germ cells (PGCs) enter mitosis to produce diploid cells (2n) and meiotic events for spermatogenesis [75]. In females, PGCs are required to produce primary oocytes through a series

of meiotic events [76]. During the DNA replication stages, replicative errors may occur, leading to variation in nucleotide sequences and MSI sites.

Previous work has shown MMR genes can contribute to abnormal chromosomal synapse formations during chromosomal recombination during meiosis, leading to MSI at D6Mit59, D7Mit91, D19Mit36 and D1Mit62 in mice [77]. It was also reported that male mice deficient in PMS2, a key protein for DNA mismatch repair, had reduced spermatozoa and reduced fertility, whilst *PMS2*<sup>-/-</sup> females were fertile. Furthermore, spermatozoa were physiologically abnormal with misshaped, truncated, and irregular flagella with abnormally shaped heads. It was concluded that spermatogenesis is disrupted at the primary spermatocyte stage during meiosis I [77]. These data suggests that MSI may also disrupt human spermatogenesis and contribute to infertility.

Data relating to MMR gene variants and their impact on gametogenesis in humans are limited. However, a study by Ferrás et al. (ref 78), investigated variants of *hMLH3*, which encodes a protein key for DNA mismatch repair. Of 300 male patients studied, 13 showed spermatogenic arrest at the primary spermatocyte stage and one of the 13 had family history of HNPCC. Between them, these 13 patients had four missense variants including; 2896T/C and 2531C/T and eight intronic (IVS9 + 66G/A) within *hMLH3* [78]. Despite of these variants, MSI was not observed in the BAT-34 microsatellite.

MMR genes have also been reported to have a regulatory role within oogenesis. *Exo1*<sup>-/-</sup> female mice have been reported to be sterile due to meiotic defects, with an increased susceptibility to lymphomas and MSI [79]. The method of detecting MSI were from DNA of *EXO1*<sup>-/-</sup> mice by PCR using end-labelled primers and electrophoresis [79]. Mutation rate analysis involved studying mutation rates of the *HRPT* locus. Embryonic stem cells exhibited either *EXO1*<sup>-/-</sup>, *MSH2*<sup>-/-</sup> and wild-type genotype, the cell line studies identified a 30-fold increase ( $0.6 \times 10^{-6}$ ) of MSI occurrences in *EXO1*<sup>-/-</sup> cells. *MSH2*<sup>-/-</sup> cells exhibited a 150-fold increase ( $3.0 \times 10^{-6}$ ) for mutation rates compared to wild-type cells [79]. Cells in meiosis I entered prophase I, but chiasmata was absent in metaphase I, leading to meiotic failure. The size of oocytes was smaller in *Exo1*<sup>-/-</sup> mice, but the oocytes had normal progression through diplonema, despite of the lack of a functioning *EXO1* protein. At day 2 postpartum,



**Fig. 1. Microsatellite structure and microsatellite instability.** Within the intron, 5'UTR and 3'UTR of a gene, microsatellites can be identified which contain short (1–6 bp) regions repeated a variable number of times (orange segments). In individuals with defects in the mismatch repair pathway (MMR) (red arrows) this can lead to three outcomes; contraction because of a gene deletion [38], a nucleotide change [39], or additional repeats are added [40] (red segments). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)


oocytes were reported to be in dictyate arrest. Additionally, MSI was detected within the microsatellites; D7Mit91, D17Mit123, and at the *uPAR* gene [79]. MSI and the effects on oogenesis in humans as not been explored but from these data suggests oogenesis in humans may also be affected by MSI.

#### Microsatellite instability and germ cell mosaicism

Mosaicism can be described as the presence of two or more genetically different sets of somatic and germline cells that contain 46 chromosomes [80], or a variation in the number of chromosomes [81]. Germline mosaicism is known to be a contributing factor to infertility in males. Jaruzelska et al. (ref 81) reported a 45 X mosaicism, within 65 men with azoospermia and 23 men with oligozoospermia. However, MSI was not reported upon within this study.

Despite the study of Jaruzelska et al. [81] described above, reports investigating the impact of germline mosaicism/MSI and infertility in humans are lacking. However, *de novo* germline mosaicism within *MSH2* was identified in one patient with a Lynch-like syndrome [82]. Due to the *de novo* variant, the patient developed endometrial cancer (aged 45)

and colorectal cancer (CRC) (aged 60). Further analysis of the CRC of this patient, showed MSI and loss of expression in the genes *MSH2* and *MSH6* [82]; similar analysis was not performed for the endometrial cancer. It should be acknowledged that previous research identified that MSI can be detected with endometrial cancer [83]. One of the clinical complications with endometrial cancer is there is an increased risk of infertility. This is due to patients with endometrial cancers having a thinner endometrial wall compared to those without cancer [84] causing reduced embryo implantation in endometrial cancer patients; however, MSI was not studied in these cancer patients [84]. In other research in MSI and cancer, Black et al. (ref 85), reported 20% of 473 endometrial cancer patients contained MSI associated with defective DNA repair [85]. Whilst Kanopiene et al. (ref 86) reported on MSI status of 100 of 109 endometrial cancers showing endometrioid type histology. From the 100 endometrial cancers, 17 demonstrated MSI-high status and 83 showed MSI-stable status. Furthermore, MSI-high cancers were linked to the pathology of deep myometrial invasion and higher histological gradings of endometrioid type cancers. These findings were identified using the Promega MSI analysis system which utilises five



**Fig. 2. The DNA mismatch repair pathway.** After DNA replication has occurred the newly replicated DNA is proofread by the MMR pathway for mismatch errors. **A)** DNA mismatches are detected by the MutSa (a heterodimer of MSH2 and MSH6) or the MutSβ complex (a heterodimer of MSH2 and MSH3). These recruit MutLa (a MLH1-PMS2 heterodimer) [41,42]. **B)** PCNA interacts with MutLa for DNA MMR repair, by which PCNA binds to DNA acting as a sliding clamp to allow DNA polymerase δ (Pol δ) and EXO1 to bind to the mismatch region [41], whilst the recruitment of replication factor C (RFC) acts as the clamp loader for Pol δ and EXO1 [44]. **C)** After the binding of EXO1 to DNA, the mismatch error is excised from the DNA whilst, Pol δ replicates the DNA to complement the 3' → 5' strand. **D)** After the binding of DNA ligase to the DNA strand, the enzyme ligates the newly synthesised strand, replacing the region where the mismatch occurred. **E)** The DNA containing the mismatch error is repaired. Figure has been adapted from [45], with permission.

mononucleotide microsatellites (BAT-25, BAT-26, NR-21, NR-24, and MONO-27) [86].

In testicular cancers, germline mosaicism remains to be investigated. The presence of MSI in testicular cancer is reported in two studies. Huddart et al. (ref 87), reported multiple MSI within two tetranucleotide microsatellites (vWfA and vWfB), two trinucleotide microsatellites (TFIID and AR) and one dinucleotide marker (D16S303), in human testicular tumours [87]. Another study reported MSI within testicular cancer complemented with reduced *MLH1* and *MSH2* expression [51]. It remains unclear whether germ cell mosaicism is accompanied with the presence of MSI. Surprisingly, neither study reported upon the fertility status of the male patients, as fertility is known to be affected by testicular cancer [88].

*Embryo implantation/postzygotic microsatellite instability through mitotic and meiotic instability*

After fertilisation, the preimplantation embryo then enters the womb for implantation. Throughout the development to a fully developed foetus, a series of cellular division events occur. These events are regulated by two spindle assembly checkpoint genes, *BUB1* and *CEP57* (discussed below), ensuring chromosomal segregation is achieved [70,89–90]. Mice with defective maternally-derived *Bub1* show embryonic aneuploidy and subsequent embryo loss after implantation [91–92]. Paternal inheritance of *Bub1<sup>m/m</sup>* (*Bub1* mutant) is not associated with embryo mitotic defects in mice [91].

In relation to MMR, recent research has suggested that *MLH1* and *MLH3* are also required for normal disjunction during meiosis. Singh et al. (ref 93), used computational analysis in different human genomic databases to investigate variants associated with *MLH1* (E268G, K618E, V326A and V716M) and *MLH3* (A3914T, R93Q and R1230H). These variants were then created in mice for phenotypic analysis [93]. To summarise the results, *MLH3<sup>41230H</sup>* males were completely infertile due to spermatocyte arrest and reduced testicular size, whilst in female mice, *MLH1<sup>E268G</sup>*, *MLH1<sup>V326A</sup>*, *MLH1<sup>K618E</sup>*, *MLH1<sup>K618T</sup>*, and *MLH1<sup>V716M</sup>*, produced offspring where the size of subsequent litters reduced over time. *MLH1<sup>K618T/K618T</sup>* females had fewer follicles, thus, suggesting chromosomal crossover events during meiosis were reduced. Further effects reported included that decreased chromosomal crossover events lead to increased aneuploidy and embryo loss, however the exact mechanism remains unanswered [93]. In *MLH3<sup>R1230H/R1230H</sup>* male mice, testicular size was reduced with complete sperm absence. Screening for MSI within mutant mice did not reveal MSI occurrences but data were not shown [93]. Previous work suggests mice that do not have a *MLH1* functional gene, have MSI and show increases in morbidity, lymphoma development, and different gastrointestinal related tumours [94].

*Genomic instability syndromes*

Genomic instability syndromes arise from heritable variants within genes involved in DNA repair mechanisms which can affect the fidelity of the genome during cell replication and cell division [95], leading to different genomic aberrations as discussed above. Variants within MMR genes leads to MSI as discussed above and a feature of many genomic instability syndromes. We here, review evidence to discern whether MSI contributes to infertility/ RM or is coincident with it but not causative in the context of genomic instability syndromes. Here we have discussed three different genomic instability syndromes (Table 1); Mosaic Variegated Aneuploidy syndrome (MVA) [70], Bloom syndrome (BS) [99], and Lynch syndrome (LS) [68].

*Aneuploidy and MVA*

Aneuploidy is associated with the genomic instability syndrome MVA1 (OMIM #257300) (associated with *BUB1B*) and MVA2 (OMIM #614114) (associated with *CEP57*). MVA1 is due to variants within the *BUB1B* gene [70,89–90]. The *BUB1B* protein regulates the spindle

**Table 1**  
Summary of genomic instability syndromes.

Genomic Instability Syndrome	Genes Mutated	Gene summary or gene variant outcome	Reference
Mosaic	<i>BUB1</i> and <i>CEP57</i>	Defective chromosomal segregation leading to chromosomal aneuploidy.	[70,72]
Variegated Aneuploidy			
Lynch Syndrome	<i>MSH2</i> , <i>MLH1</i> , <i>PMS2</i> , <i>MSH6</i>	Variants in these MMR genes gives arise to a range of different cancers associated to Lynch syndrome, including colorectal cancers.	[96,59]
Bloom Syndrome	<i>BLM</i>	Essential for homologous recombination for DNA double strand break repair. Variants of <i>BLM</i> can lead to leukaemia and Burkitt lymphoma.	[97–98]

assembly checkpoint during cellular division and therefore, maintains genomic stability by delaying cell division until full chromosome segregation is achieved [100–101]. Variants within the *BUB1B* gene lead to monosomy or trisomy [102–103].

Patients who have a mutation in the *BUB1B* gene are diagnosed with MVA1 [19,104]. The research on MVA1 shows fertility status can be unaffected and patients can successfully conceive (and this syndrome is heritable) [13,105]. There is no research linking MSI to infertility in MVA1 in human patients. Baker et al. (ref 106), generated *Bub1b<sup>-</sup>* (knockout) *Bub1b<sup>H</sup>* (hypomorphic) and *Bub1b<sup>+</sup>* (wildtype) mice, investigating the role for *Bub1b*, resembling MVA1. The *Bub1b<sup>-/-</sup>* offspring died during early embryo development, *Bub1b<sup>-/H</sup>* offspring survived birth but died 12 h after birth. Furthermore, mouse embryo fibroblasts (MEFs) were also examined and revealed *Bub1b<sup>H/H</sup>* MEFs had deficient cell cycle arrest and aneuploidy, whilst *Bub1b<sup>-/H</sup>* MEFs had higher aneuploidy compared to *Bub1b<sup>H/H</sup>*. Male *Bub1b<sup>H/H</sup>* mice were infertile due to a 4-fold reduction in sperm compared to *Bub1b<sup>+/+</sup>* mice. *Bub1b<sup>+/+</sup>* mice were reported to be fully fertile. The crossing of *Bub1b<sup>H/H</sup>* with *Bub1b<sup>+/+</sup>* mice led to a reduction in the number of fertilised eggs [106]. Therefore, these data show that *Bub1b* is essential for spindle assembly checkpoints and defects cause aneuploidy; MSI was not reported upon. In humans, infertility associations with MVA1 have not been reported to date.

Mosaic variegated aneuploidy syndrome (MVA2) (OMIM #614114) is caused by variants within the *CEP57* gene, and this is a diagnostic biomarker [72]. *CEP57* protein functions to maintain microtubule stability by ensuring that spindle microtubules are attached to chromosomes for proper segregation [107–108]. A case study found secondary infertility could be a clinical feature of MVA2, but this is complicated by the consanguineous nature of the couple involved [72]. At the time of writing this review, MSI has not been detected within MVA1 or MVA2.

*Bloom syndrome*

Bloom syndrome (BS) (OMIM #210900) is the product of mutations to the causative gene, *BLM*. The *BLM* protein functions to repair DNA double stranded breaks (DSBs) via homologous recombination [97,109]. Therefore, the presence of a *BLM* variant is associated with an increased rate of MSI, due to abnormal expression patterns of the *BLM* gene [110,111]. Furthermore, other research not related to BS specifically, has associated MMR defects for DSB repair gene transcripts with MSI [112]. BS is an autosomal recessive syndrome which phenotypically presents as red rashes around the facial area, short stature and elongated limbs [113]. BS predisposes patients to other complications including type 2 like diabetes mellitus [114], sun-sensitive skin [115], moderate to severe immunodeficiency [115], and cancers including leukaemia and Burkitt lymphoma [116,110].

*BLM* gene variants in BS have also been linked to infertility. While



two case studies have revealed that female patients with BS are able to successfully achieve pregnancy and conceive [117–118], male patients with BS have been reported to be infertile with oligozoospermia or azoospermia [119]. El Ghamrasni et al. (ref 120) reported a higher occurrence of genomic instability including chromosome breaks and chromosome fragmentation of dicentric chromosomes in *Blm*<sup>-/-</sup> and *Mus81*<sup>-/-</sup> mice compared to wildtype mice. This study did not report on MSI [120]. However, this suggests MSI may be associated with BS as *BLM* is a DNA repair participating gene.

BS is also associated with higher rates of sister chromatid exchange (SCE), a process by which sister chromatids break and re-join [121]. For instance, Howell and Davis [122] reported SCE counts between 31 and 70 in BS patient, compared to normal chorionic cells that contained 6 to 8 SCE counts; these discrepancies can be used to diagnose BS [122]. Other work has suggested that the rates of SCE in BS patients are between 10 and 12 times more than those without BS [123]. In mice models of BS, telomere length was observed to be reduced due to the loss of telomerase activity and a loss in telomeric repeats in fibroblast cells [124]. Therefore, telomeric dysregulation could also be a molecular characteristic within BS patients. Previous research has found short telomere length associated with MSI [125–128]. The mechanism linking these two processes is not entirely clear.

*DNMT1* codes for DNA methyltransferase 1 which participates in DNA methylation [129–130]. DNA methylation involves the attachment of methyl groups to CpG islands, suppressing gene expression [131–133]. In relation to MSI in BS, Guo et al. (ref 20), screened for MMR deficient genes in *blm*-deficient embryonic stem cells. This analysis revealed *Dnmt1* as a putative novel MMR gene, as *dnmt1*-deficient cells exhibited MSI and *Dnmt1* expression was not detectable [20]. A similar study also reported *DNMT1* as a novel MMR gene, using *dnmt1*-deficient mouse embryonic stem cells [134]. These data also suggest that there is a potential postzygotic predisposition to MSI in BS patients; whether this contributes to infertility remains unclear.

Only one study attempted to analyse MSI in six unrelated BS patients, by PCR and cell line analysis [135]. This study used eight different cell lines, from six unrelated patients which included five BS fibroblast cell lines (GM02520A, GM03402A, GM03498C, GM03509A, GM03510), one BS SV40-transformed fibroblast cell line (GM08505B) and two BS EBV-transformed lymphoblastoid B cell lines (GM03403D, GM04408A) where the cell lines GM03498C and GM04408A came from the same patient [135]. This analysis reported that MSI could not be detected in all six patients.

Patients with BS are predisposed to an increased risk of leukaemia and Burkitt lymphoma. Both cancer types arise in childhood in BS patients [98]. Of note, MSI has been reported within a BS-associated lymphoma. In this study, two Japanese siblings aged 23 and 25 of consanguineous parents were studied [136]. Genomic analysis revealed MSI within the *p53* gene and a 3-bp deletion in the *BLM* gene [136]. MSI in non-cancerous cells was not evaluated in this report.

### Lynch syndrome

Lynch syndrome (LS) (OMIM #120435) is associated with variants in *MLH1* [137], *MSH2* [138], *MSH6* [68], and/or *PMS2* [139]. These variants contribute to the onset of multiple cancers including endometrial [96], breast [140], colorectal [141], and prostate [142]. Recently, for the first time, testicular cancer was found to be associated with LS patients [143]. Testicular cancer may not be a common cancer to occur in male LS patients.

Infertility is associated with LS. A case study revealed that females diagnosed with LS and CRC, have decreased fertility across a broad age range (15 to 47) compared to those with CRC but without LS [144]. This was expressed as “children per woman and is calculated as the sum of age-specific fertility ages 15 to 49”. This is based on a total fertility rate value defined as “an average number of children produced of a hypothetical cohort of women had by the end of their reproductive life, if

they children at the population age-specific rates during their whole life” [144]. In LS male patients decreased fertility was also observed in the same age range [144]. Even though mutations in MMR genes have been linked to male infertility contributing to oligozoospermia and azoospermia [145–148], research to date has not addressed how LS is specifically linked to male infertility.

LS is associated with other cancers as identified above in this section. Limited research thus far has reported MSI in LS. For instance, Latham et al. (ref 59), studied 15,409 patients with more than 50 different types of cancers present in LS patients. 53 of 326 (16.3%) had high levels of MSI (MSI-H), 13 of 699 (1.9%) has low MSI (MSI-L) and 37 of 14,020 (0.3%) showed microsatellite stable phenotype (MSS). Furthermore, 33 of 66 (50%) had primary tumours excluding CRC. In LS-related tumours containing MSI (MSI-H/L), 12 of 32 (37.5%) were urothelial, 26 of 137 (19%) were CRC, 2 of 13 (15.4%) were gastric cancers. In those with high or low levels of MSI (MSI-H/L) pancreatic cancers 34 of 824 (4.1%); only 5 of these 34 (14.7%) were LS related [59], involving five germline MMR genes (*MLH1*, *MSH2*, *MSH6*, *PMS2* and *EPCAM*). MSI can be detected in non-LS related breast [149], colorectal [150], prostate [151] and testicular cancers [87]. Therefore, further research is important for the investigation of MSI. However, as Latham et al. (ref 59) revealed, LS is complicated by the presence of LS and non-LS related cancers which may prove difficult to address MSI related studies in LS patients.

### Microsatellite instability in infertility and recurrent miscarriage

In this section we have discussed MSI occurrences in infertile patients independent of genomic instability syndromes. After searching the literature, we found four papers [152–155], that reported links between MSI and infertility/RM, independent of genomic instability syndromes. We have identified that from these four studies that MSI may be a common feature in idiopathic infertility. These studies reveal MSI is associated with infertility/RM and spontaneous abortions. The association of MSI within preimplantation failure in humans remains unclear. Some molecular mechanisms have recently been proposed for the potential reasons for preimplantation failure [156], discussed further below.

In the study of Kiaris et al. (ref 152) MSI was investigated at two different distal sites per spontaneously aborted embryo and compared against those of a normal birth. Thirty spontaneously aborted embryonic tissues were selected at the time of rejection and frozen until DNA analysis. Eight different microsatellite markers were selected: D6S344, D7S531, HRM, D12S94, D13S175, TCRD, THRA1 and D19S49. MSI was reported in 12 (40%) of the 30 specimens. In aborted embryonic tissue samples, in samples coming from those without a normal childbirth 61% (11 of 18 cases) had MSI. In samples that came from a normal birth, only 8% (1 of 12 cases) had MSI. THRA1 and D6S344 presented instability in 5 of the cases whilst D7S531 did not present MSI [152]. Both THRA1 and D6S344 have been found to be highly polymorphic microsatellite markers in other studies [157–160]. This provides evidence that MSI increases at the embryonic stages of pregnancy. Thus, MSI analyses should not be overlooked during genetic analysis of infertile patients and in embryos.

A second report focused on infertility associated with an MSI site in the *CTLA-4* gene [153]. The protein product of the *CTLA-4* gene (*CTLA-4* protein) is an inhibitory receptor that maintains T-cell homeostasis through suppressive mechanisms [161–162]. Tsai et al. (ref 153) hypothesised that a microsatellite residing within the 3'UTR of the *CTLA-4* gene, compromised of an AT(n) repeat region (between 16 bp and 46 bp total length), leads to mRNA instability thereby leading to altered *CTLA-4* protein expression. In turn this may alter immune tolerance of the foetus in utero and predispose to foetal rejection and infertility. To test this hypothesis, 60 couples were studied, each with a minimum of 3 unexplained spontaneous miscarriages and compromising a total cohort of 51 liveborn children and 10 abortuses. Cord blood and placental tissues from both liveborn and aborted fetuses

were genotyped for the CTLA-4 microsatellite locus. This revealed that the shorter alleles, inherited from the maternal side, were associated with increased foetal survival. Thus, mRNA transcribed from the longer alleles were shown to be intrinsically unstable, with reduced expression [153]. Other studies have also revealed a link between CTLA-4 gene expression patterns and infertility but did not report upon MSI [163–164]; whether protein expression is significantly altered by the CTLA-4 MSI site remains to be seen.

A third study aimed to evaluate MSI incidence in 35 spontaneously aborted (between 6 and 20 weeks of gestation) embryonic tissues [154]. This study compared seven different microsatellite markers between parents and embryos; D3S1234, D4S194, D10S109, HRM, TCRD, THRA1 and D17S855. In total, 19 of the patients who had a history of spontaneous abortion, had at least one instability in at least one of the microsatellite markers in aborted tissue samples [154]. This is supporting the idea that MSI in an embryo contributes to spontaneous abortions, alike to Tsai et al. [153].

Finally, a pilot study aimed to identify genetic loci that could be contributing factors for idiopathic recurrent miscarriage (IRM) [155]; 45 patients with IRM in the first trimester were included along with 44 healthy controls that were matched by sex and ethnicity. Of the 403 evaluable microsatellite markers studied, 17 were different between IRM patients and controls of which D6S446 (6q27), D9S1776 (9q31.1) and DXS1226 (Xp22.11), were found to have an association with IRM. Further analysis to identify genes around these sites, revealed 26 genes including *TNFSF8* and *TNFSF15* which are linked to immune function [155]. Of note, other members of the TNF superfamily are present at the maternal-foetal interface and likely play a key role in preventing embryonic rejection [165,166]. Despite these findings, confirmatory studies with patients with IRM of other ethnicities should be conducted to confirm if MSI is present in the same microsatellites or to reveal other unstable microsatellites.

We did not find any research assessing MSI in preimplantation embryos. A recent review suggests there are several potential molecular mechanisms including DNA damage and repair, aneuploidy and meiotic recombination that could contribute to genomic instability within a preimplantation embryo [156] and which may lead to spontaneous abortion. More research within this area is clearly needed.

### Discussion and conclusion

Genomic instabilities are a result of chromosomal aberrations serving as a genomic predisposition for genomic instability syndromes. Microsatellites are 1–6 bp units, encompassing approximately 3% of the human genome [22]. MSI is a result of mutations to genes in the DNA MMR pathway [18,167–168]. The aim of this review was to discuss research into the role of MSI and genomic instability syndromes in infertility/RM. Our findings reveal that MSI is poorly investigated in genetic analyses of patients with genomic instability syndromes and infertility, including patients with infertility/RM independent of genomic instability syndromes. Potential reasons that may account for this are that; 1) MSI is not a core component of genetic analyses for infertility or genomic instability, 2) current methods used in genetic analyses may not identify MSI as a genetic feature, 3) MSI may happen during preimplantation which may lead to technical and ethical barriers for analyses.

Preimplantation genetic testing (PGT) is used in assisted reproduction methods to determine the genetic profile of embryos and to reveal whether an embryo is at risk of inheriting a disease or syndrome. PGT can be used to identify monogenic diseases (PGT-M) [169], chromosomal structural rearrangements (PGT-SR) [170] and aneuploidy (PGT-A) [170–171]. However, these current PGT methods do not address investigate MSI within infertile patients and/or in the embryo. From the studies that have been reviewed here we suggest that MSI may be a genomic instability feature a contributor to idiopathic infertility.

To conclude, we comprehensively reviewed limited research to date

to reveal the role of microsatellite instability in genomic instability syndromes and infertility. We suggest the need for further research to define the role of microsatellite instability in infertility, including microsatellite instability in preimplantation embryos. We therefore propose the use of microsatellite instability as a biomarker of underlying DNA repair deficiencies resulting in unexplained infertility.

### Key messages

The involvement of microsatellite instability in genomic instability syndromes and infertility is poorly understood. Whether infertility is a clinical phenotype of genomic instability syndromes remains unclear.

### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### References

- [1] Thonneau P, Spira A. Prevalence of infertility: international data and problems of measurement. *Eur J Obstet Gynecol Reprod Biol* 1990;38(1):43–52.
- [2] Templeton A, Fraser C, Thompson B. The epidemiology of infertility in Aberdeen. *BMJ* 1990;301(6744):148–52.
- [3] Rai R, Regan L. Recurrent miscarriage. *The Lancet* 2006;368(9535):601–11.
- [4] Duckitt K, Quershi A. 2011. Recurrent miscarriage. *BMJ Clin Evid* 2011; 2011 (1409).
- [5] Sharlip ID, Jarow JP, Belker AM, Lipshultz LI, Sigman M, Thomas AJ, et al. Best practice policies for male infertility. *Fertil Steril* 2002;77(5):873–82.
- [6] Ellis NA, Ciocci S, Proytcheva M, Lennon D, Groden J, German J. The Ashkenazic Jewish Bloom Syndrome Mutation *blm*<sup>AsH</sup> Is Present in Non-Jewish Americans of Spanish Ancestry. *Am J Hum Gen* 1998;63(6):1685–93.
- [7] Barakat A, Ababou M, Onclercq R, Dutertre S, Chadli E, et al. Identification of a novel BLM missense mutation (2706T>C) in a Moroccan patient with Bloom's syndrome. *Hum Mutat* 2000;15(6):584–655.
- [8] Mori M, Hira A, Yoshida K, Muramatsu H, Okuno Y, Shiraishi Y, et al. 2003. Pathogenic mutations identified by a multimodality approach in 117 Japanese Fanconi anemia patients. *Haematologica* 2003; 104 (10): 1962–73.
- [9] Varon R, Dutranoy V, Weikert G, Tanzarella C, Antoccia A, Stöckl L, et al. Mild Nijmegen breakage syndrome phenotype due to alternative splicing. *Hum Mol Genet* 2006; 15 (5): 679–89.
- [10] Ameiziane N, Errami A, Léveillé F, Fontaine C, de Vries Y, van Spaendonck RML, et al. Genetic subtyping of Fanconi anemia by comprehensive mutation screening. *Hum Mutat* 2008;29(1):159–66.
- [11] Snape K, Hanks S, Ruark E, Barros-Núñez P, Elliott A, Murray A, et al. Mutations in CEP57 cause mosaic variegated aneuploidy syndrome. *Nat Genet* 2011;43(6): 527–9.
- [12] Prokofyeva D, Bogdanova N, Dubrowskaja N, Bermisheva M, Takhirova Z, Antonenkova N, et al. Nonsense mutation p. Q548X in BLM, the gene mutated in Bloom's syndrome, is associated with breast cancer in Slavic populations. *Breast Cancer Res Treat* 2013;137(2):533–9.
- [13] Cho CH, Oh M-J, Lim CS, Lee CK, Cho Y, Yoon S-Y. A Case Report of a Fetus with Mosaic Autosomal Variegated Aneuploidies and Literature Review. *Ann Clin Lab Sci* 2015;45(1):106–9.
- [14] Zareifars S, Dastsooz H, Shahriari M, Faghihi MA, Shekarkhar G, Bordbar M, et al. A novel frame-shift deletion in FANCF gene causing autosomal recessive Fanconi anemia: a case report. *BMC Medical Genet* 2019;20(1). <https://doi.org/10.1186/s12881-019-0855-2>.
- [15] Lathi RB, Westphal LM, Milki AA. Aneuploidy in the miscarriages of infertile women and the potential benefit of preimplantation genetic diagnosis. *Fertil Steril* 2008;89(2):353–437.
- [16] Stern C, Pertile M, Norris H, Hale L, Baker HWG. Chromosome translocations in couples with in-vitro fertilization implantation failure. *Hum Reprod* 1999;14(8): 2097–20101.
- [17] Gekas J, Thepot C, Turleau C, Siffroi JP, Dadoune JP, Briault S, et al. Chromosomal factors of infertility in candidate couples for ICSI: an equal risk of constitutional aberrations in women and men. *Hum Reprod* 2001; 16 (1): 82–90.
- [18] Chang PL, Sauer MV, Brown S. Y chromosome microdeletion in a father and his four infertile sons. *Hum Reprod* 1999; 14 (11): 2689–94.
- [19] Jacquemont S, Bocéno M, Rival JM, Méchinaud F, David A. High risk of malignancy in mosaic variegated aneuploidy syndrome: High Risk of Malignancy in MVA Syndrome. *Am J Med Genet* 2002;109(1):17–21.
- [20] Guo Ge, Wang W, Bradley A. Mismatch repair genes identified using genetic screens in BLM-deficient embryonic stem cells. *Nature* 2004;429(6994):891–5.
- [21] Maurer M, Hoffmann K, Sperling K, Varon R. High prevalence of the NBN gene mutation c.657-661del5 in Southeast Germany. *J Appl Genet* 2010;51(2):211–4.

- [22] Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, et al. Initial sequencing and analysis of the human genome. *Nature* 2001;409(6822):860–921.
- [23] Metzgar D, Bytof J, Wills C. Selection Against Frameshift Mutations Limits Microsatellite Expansion in Coding DNA. *Genome Res* 2000; 10 (1): 72–80.
- [24] Suraweera N, Iacopetta B, Duval A, Compont A, Tubacher E, Hamelin R. Conservation of mononucleotide repeats within 3' and 5' untranslated regions and their instability in MSI-H colorectal cancer. *Oncogene* 2001;20(51): 7472–747.
- [25] Dieringer D, Schlötterer C. Two Distinct Modes of Microsatellite Mutation Processes: Evidence From the Complete Genomic Sequences of Nine Species. *Genome Res* 2003;13(10):2242–51.
- [26] Jung B, Doctolero RT, Tajima A, Nguyen AK, Keku T, Sandler RS, et al. Loss of activin receptor type 2 protein expression in microsatellite unstable colon cancers. *Gastroenterology* 2004;126(3):64–659.
- [27] Ramamoorthy S, Garapati HS, Mishra RK. Length and sequence dependent accumulation of simple sequence repeats in vertebrates: Potential role in genome organization and regulation. *Gene* 2014;551(2):167–75.
- [28] Arabfard M, Kavousi K, Delbari A, Ohadi M. Link between short tandem repeats and translation initiation site selection. *Hum Genomics* 2018;12(1). <https://doi.org/10.1186/s40246-018-0181-3>.
- [29] Urquhart A, Kimpton CP, Downes TJ, Gill P. Variation in Short Tandem Repeat sequences — a survey of twelve microsatellite loci for use as forensic identification markers. *Int J Legal Med* 1994;107(1):13–20.
- [30] Takahashi M, Kato Y, Mukoyama H, Kanaya H, Kamiyama S. Evaluation of five polymorphic microsatellite markers for typing DNA from decomposed human tissues. *Forensic Sci Int* 1997;90(1–2):1–9.
- [31] Hoff-Olsen P, Jacobsen S, Mevåg B, Olaisen B. Microsatellite stability in human post-mortem tissues. *Forensic Sci Int* 2001;119(3):273–328.
- [32] Plevy SE, Targan SR, Yang H, Fernandez D, Rotter JI, Toyoda H. Tumor Necrosis Factor Microsatellites Define a Crohn's Disease-Associated Haplotype on Chromosome 6. *Gastroenterology* 1996;110(4):1053–60.
- [33] Mizuki N, Ota M, Yabuki K, Ando, H, Palimeris, GD, Kaklamani, E, Accorinitim M., Pivetti-Pezzi P, Ohno S, Inoko H. Localization of the Pathogenic Gene of Behçet's Disease by Microsatellite Analysis of Three Different Populations. *Invest Ophthalmol Vis Sci* 2000; 41 (12): 3702–8.
- [34] Collins JE, Heward JM, Carr-Smith J, Daykin J, Franklyn JA, Gough SCL. Association of a Rare Thyroglobulin Gene Microsatellite Variant with Autoimmune Thyroid Disease. *J Clin Endocrinol Metab* 2003;88(10):5039–42.
- [35] Ellegren H. Microsatellites: simple sequences with complex evolution. *Nat Rev Genet* 2004;5(6):435–45.
- [36] Messer PW, Arndt PF. The Majority of Recent Short DNA Insertions in the Human Genome Are Tandem Duplications. *Mol Biol Evol* 2007;24(5):1190–2117.
- [37] Leclercq S, Rivals E, Jarne P. DNA Slippage Occurs at Microsatellite Loci without Minimal Threshold Length in Humans: A Comparative Genomic Approach. *Genome Biol Evol* 2010;2(0):325–35.
- [38] Ghimenti C, Tannergård P, Wahlberg S, Liu T, Giulianotti PG, Mosca F, et al. Microsatellite instability and mismatch repair gene inactivation in sporadic pancreatic cancer and colon tumors. *Br J Cancer* 1999;80(1–2):11–6.
- [39] Nudell D, Castillo M, Turek PJ, et al. Increased frequency of mutations in DNA from infertile men with meiotic arrest. *Hum Reprod* 2000; 15 (6): 1289–94.
- [40] Liu L, Markowitz S, Gerson SL. Mismatch Repair Mutations Override Alkyltransferase in Conferring Resistance to Temozolomide but not to 1,3-Bis(2-chloroethyl)nitrosourea. *Cancer Res* 1996; 56 (23): 5375–79.
- [41] Kadyrov FA, Dzantiev L, Constantin N, Modrich P. Endonucleolytic Function of MutLα in Human Mismatch Repair. *Cell* 2006;126(2):297–308.
- [42] Warren JJ, Pohlhaus TJ, Changela A, Lyster RR, Modrich PL, Beese LS. Structure of the Human MutSα DNA Lesion Recognition Complex. *Mol Cell* 2007;46(4): 579–92.
- [43] Umar A, Risinger JI, Glaab WE, Tindall KR, Barrett JC, Kunkel TA. Functional Overlap in Mismatch Repair by Human MSH3 and MSH6. *Genetics* 1998;148(4): 1637–46.
- [44] Pavlov YI, Frahm C, McElhinny SA, Niimi A, Suzuki M, Kunkel TA. Evidence that Errors Made by DNA Polymerase alpha are Corrected by DNA Polymerase delta. *Curr Biol* 2006;16(2):202–7.
- [45] Pecina-Slaus N, Kafka A, Salamon I, Bukovac A. Mismatch Repair Pathway, Genome Stability and Cancer. *Front Mol Biosci* 2020;7(122). <https://doi.org/10.3389/fmolb.2020.00122>.
- [46] Brimmell M, Mendiola R, Mangion J, Packham G. BAX frameshift mutations in cell lines derived from human haemopoietic malignancies are associated with resistance to apoptosis and microsatellite instability. *Oncogene* 1998;16(14): 1803–12.
- [47] Klingler H, Hemmerle C, Bannwart F, Haider R, Cattaruzza MS, Marra G. Expression of the hMSH6 mismatch-repair protein in colon cancer and HeLa cells. *Swiss Med Wkly* 2002;132(5–6):57–63.
- [48] Shaw JA, Walsh T, Chappell SA, Carey N, Johnson K, Walker RA. Microsatellite instability in early sporadic breast cancer. *Br J Cancer* 1996;73(11):1393–2137.
- [49] Basil JB, Goodfellow PJ, Rader JS, Mutch DG, Herzog TJ. Clinical significance of microsatellite instability in endometrial carcinoma. *Cancer* 2000;89(8):1758–64.
- [50] Tang R, Changchien CR, Wu M-C, Fan C-W, Liu K-W, Chen J-S, et al. Colorectal cancer without high microsatellite instability and chromosomal instability—an alternative genetic pathway to human colorectal cancer. *Carcinogenesis* 2004;25 (5):841–86.
- [51] Velasco A, Corvalan A, Wistuba II, Riquelme E, Chuaqui R, Majerson A, et al. Mismatch repair expression in testicular cancer predicts recurrence and survival. *Int J Cancer* 2008;122(8):1774–7.
- [52] Fleisher AS, Esteller M, Harpaz N, Leytin A, Rashid A, Xu Y, et al. 2000. Microsatellite instability in inflammatory bowel disease-associated neoplastic lesions is associated with hypermethylation and diminished expression of the DNA mismatch repair gene, hMLH1. *Cancer Res* 2000; 60 (17): 4864–48.
- [53] Lamberti C, Kruse R, Ruelfs C, Caspari R, Wang Y, Jungck M, et al. Microsatellite instability—a useful diagnostic tool to select patients at high risk for hereditary non-polyposis colorectal cancer: a study in different groups of patients with colorectal cancer. *Gut* 1999;44:839–43.
- [54] Ruszkiewicz A, Bennett G, Moore J, Manavis J, Rudzki B, Shen L, et al. Correlation of mismatch repair genes immunohistochemistry and microsatellite instability status in HNPCC-associated tumours. *Pathology* 2002;34(6):541–7.
- [55] Romanowicz-Makowska H, Smolarz B, Langer E, Kozłowska E, Kulig A, Dziki A. Analysis of microsatellite instability and BRCA1 mutations in patients from hereditary nonpolyposis colorectal cancer (HNPCC) family. *Pol J Pathol* 2005;56 (1):21–6.
- [56] Belvederesi L, Bianchi F, Galizia E, Loretell C, Bracci R, Catalani R, et al. MSH2 missense mutations and HNPCC syndrome: pathogenicity assessment in a human expression system. *Hum Mutat* 2008;29(11):296–309.
- [57] Karamurzin Y, Zeng Z, Stadler ZK, Zhang L, Ouassafi I, Al-Ahmadie HA, et al. Unusual DNA mismatch repair-deficient tumors in Lynch syndrome: a report of new cases and review of the literature. *Hum Pathol* 2012;43(10):1677–87.
- [58] Boissière-Michot F, Frugier H, Ho-Pun-Cheung A, Lopez-Crapez E, Duffour J, Bibeau F. Immunohistochemical staining for p16 and BRAFV600E is useful to distinguish between sporadic and hereditary (Lynch syndrome-related) microsatellite unstable colorectal carcinomas. *Virchows Arch* 2016;469(2): 135–44.
- [59] Latham A, Srinivasan P, Kemel Y, Shia J, Bandlamudi C, Mandelker D, et al. Microsatellite Instability Is Associated With the Presence of Lynch Syndrome Pan-Cancer. *J Clin Oncol* 2019;37(4):286–95.
- [60] Yamamoto G, Takenoya T, Takahashi A, Osanai Y, Kakuta M, Tachikawa T, et al. Quantitative evaluation of MSI testing using NGS detects the imperceptible microsatellite changed caused by MSH6 deficiency. *Fam Cancer* 2021;20(2): 137–43.
- [61] Ellis NA, Roe AM, Kozloski J, Proytcheva M, Falk C, German J. Linkage disequilibrium between the FES, D15S127, and BLM loci in Ashkenazi Jews with Bloom syndrome (3). *Am J Hum Genet* 1994;55:453–60.
- [62] Foucault F, Vaury C, Barakat A, Thibout D, Planchon P, Jaulin C, et al. Characterization of a new BLM mutation associated with a topoisomerase II alpha defect in a patient with Bloom's syndrome. *Hum Mol Genet* 1997;6(9):1427–34.
- [63] Hiort O, Holterhus P-M, Horter T, Schulze W, Kremke B, Bals-Pratsch M, et al. Significance of Mutations in the Androgen Receptor Gene in Males with Idiopathic Infertility. *J Clin Endocrinol Metab* 2000;85(8):2810–3285.
- [64] Laven JSE, Mulders AGMJ, Suryandari DA, Gromoll J, Nieschlag E, Fauser BCJM, et al. Follicle-stimulating hormone receptor polymorphisms in women with normogonadotropic anovulatory infertility. *Fertil Steril* 2003;80(4): 986–92.
- [65] Savino M, Borriello A, d'Apolito M, Criscuolo M, Del Vecchio M, Bianco AM, et al. Spectrum of FANCA mutations in Italian Fanconi anemia patients: Identification of six novel alleles and phenotypic characterization of the S858R variant. *Hum Mutat* 2003;22(4):338–9.
- [66] Ferlin A, Vinanzi C, Garolla A, Selice R, Zuccarello D, Cazzadore C, et al. Male infertility and androgen receptor gene mutations: clinical features and identification of seven novel mutations. *Clin Endocrinol* 2006;65(5):606–10.
- [67] Meyer S, Kingston H, Taylor AMR, Byrd PJ, Last JIK, Brennan BMD, et al. Rhabdomyosarcoma in Nijmegen breakage syndrome: strong association with perianal primary site. *Cancer Genet Cytogenet* 2004;154(2):169–74.
- [68] Baglietto L, Lindor NM, Dowty JG, White DM, Wagner A, Garcia EBG, et al. Risks of Lynch Syndrome Cancers for MSH6 Mutation Carriers. *J Natl Cancer Inst* 2010; 102 (3): 193–201.
- [69] Du J-W, Xu K-Y, Qi X-L, 2011. Association between mutations of the luteinizing hormone β subunit and female infertility. *Mol Med Rep* 2011; 5 (2): 473–76.
- [70] Schmid M, Steinlein C, Tain Q, Newell AEH, Gessler M, Olson SB, et al. Mosaic variegated aneuploidy in mouse B6R1 deficient embryos and pregnancy loss in human. *Chromosome Res* 2014; 22 (3): 375–92.
- [71] Karimian M, Nikzad H, Azami-Tameh A, Taherian A, Darvishi FZ, Haghighatnia MJ. SPO11-C631T Gene Polymorphism: Association With Male Infertility and an in Silico-Analysis. *J Family Reprod Health* 2015;9(4):155–63.
- [72] Aljaser MT, Bahzad S. The use of whole exome sequencing to identify carriers of Mosaic Variegated Aneuploidy (MVA): A case study. *Integr Mol Med* 2018; 5 (3): doi: 10.15761/IMM.1000332.
- [73] Chen B, Wang W, Peng X, Jiang H, Zhang S, Li Da, et al. The comprehensive mutational and phenotypic spectrum of TUBB8 in female infertility. *Eur J Hum Genet* 2019;27(2):300–7.
- [74] Wyrwoll MJ, Temel SG, Nagirnaja L, Oud MS, Lopes AM, van der Heijden GW, et al. Bi-allelic Mutations in MIAP Are a Frequent Cause of Meiotic Arrest and Severely Impaired Spermatogenesis Leading to Male Infertility. *Am J Hum Genet* 2020;107(2):342–51.
- [75] Wang H, Zhou Z, Xu M, Li J, Xiao J, Xu Z-Y, et al. A spermatogenesis-related gene expression profile in human spermatozoa and its potential clinical applications. *J Mol Med* 2004;82(5):317–24.
- [76] Stoop H, Honecker F, Cools M, de Krijger R, Bokemeyer C, Looijenga LHJ. Differentiation and development of human female germ cells during prenatal gonadogenesis: an immunohistochemical study. *Hum Reprod* 2005;20(6): 1466–76.



- [77] Baker SM, Bronner CE, Zhang L, Plug AW, Robatzek M, Warren G, et al. Male Mice Defective in the DNA Mismatch Repair Gene PMS2 Exhibit Abnormal Chromosome Synapsis in Meiosis. *Cell* 1995;82(2):309–19.
- [78] Ferrás C, Zhou X-I, Sousa M, Lindblom A, Barros A. DNA mismatch repair gene hMLH3 variants in meiotic arrest. *Fertil Steril* 2007;88(6):1681–2164.
- [79] Wei K, Clark AB, Wong E, Kane MF, Mazur DJ, Parris T, et al. Inactivation of Exonuclease 1 in mice results in DNA mismatch repair defects, increased cancer susceptibility, and male and female sterility. *Genes Dev* 2003;17(5):603–14.
- [80] Muné S, Weier HUG, Grifo J, Cohen J. Chromosome Mosaicism in Human Embryo. *Biol Reprod* 1994;51(3):373–439.
- [81] Jaruzelska J, Korcz A, Wojda A, Jedrzejczak . Mosaicism for 45,X cell line may accentuate the severity of spermatogenic defects in men with AZFc deletion. *J Med Genet* 2001; 38 (11), 798–802.
- [82] Guillem E, Svrcek M, Bardier-Dupas A, Basset N, Coulet F, Colas C. Molecular tumor testing in patients with Lynch-like syndrome reveals a de novo mosaic variant of a mismatch repair gene transmitted to offspring. *Eur J Hum Genet* 2020;28(11):1624–2168.
- [83] Kanopienė D, Smalytė G, Vidugiriene J, Bacher J. Impact of microsatellite instability on survival of endometrial cancer patients. *Medicina* 2014;50(4): 216–22.
- [84] Fujimoto A, Ichinose M, Harada M, Hirata T, Osuga Y, Fujii T. The outcome of infertility treatment in patients undergoing assisted reproductive technology after conservative therapy for endometrial cancer. *J Assist Reprod Genet* 2014;31(9): 1189–94.
- [85] Black D, Soslow RA, Levine DA, Tornos C, Chen SC, Hummer AJ, et al. Clinicopathologic Significance of Defective DNA Mismatch Repair in Endometrial Carcinoma. *J Clin Oncol* 2006;24(11):1745–53.
- [86] Kanopienė D, Vidugiriene J, Valuckas KP, Smalytė G, Uleckiene S, Bacher J. Endometrial cancer and microsatellite instability status. *Open Med* 2015;10(1): 70–6.
- [87] Huddart RA, Wooster R, Horwich A, Cooper CS. Microsatellite instability in human testicular germ cell tumours. *Br J Cancer* 1995;72(3):642–65.
- [88] Spermon RJ, Kiemeny LALM, Meuleman EJH, Ramos L, Wetzels AMM, Witjes AJ. Fertility in men with testicular germ cell tumors. *Fertil Steril* 2003;79 (3):1543–2159.
- [89] Kato M, Kato T, Hosoba E, Ohashi M, Fujisaki M, Ozaki M, et al. PCS/MVA syndrome caused by an Alu insertion in the BUB1B gene. *Hum Genome Var* 2017; 4(1). <https://doi.org/10.1038/hgv.2017.21>.
- [90] Laberko A, Balashov D, Deripapa E, Soldatkina O, Raikina E, Maschan A, et al. Hematopoietic stem cell transplantation in a patient with type 1 mosaic variegated aneuploidy syndrome. *Orphanet J Rare Dis* 2019;14(1). <https://doi.org/10.1186/s13023-019-1073-x>.
- [91] Leland S, Nagarajan P, Polyzos A, Thomas S, Samaan G, Donnell R, et al. Heterozygosity for a Bub1 mutation causes female-specific germ cell aneuploidy in mice. *Proc Natl Acad Sci U S A* 2009;106(31):12776–81.
- [92] Tilston V, Taylor SS, Perera D. Inactivating the spindle checkpoint kinase Bub1 during embryonic development results in a global shutdown of proliferation. *BMC Res Notes* 2009;2(1):190.
- [93] Singh P, Fragoza R, Blengini CS, Tran TN, Pannafino G, Al-Sweel N, et al. Human MLH1/3 variants causing aneuploidy, pregnancy loss, and premature reproductive aging. *Nature Comms* 2021;12(1). <https://doi.org/10.1038/s41467-021-25028-1>.
- [94] Edelmann W, Yang K, Kuraguchi K, Heyer J, Lia M, Kneitz B, et al. Tumorigenesis in Mlh1 and Mlh1/Apc1638N mutant mice. *Cancer Res* 1999;59(6):1301–7.
- [95] Lujan SA, Clausen AR, Clark AB, MacAlpine HK, MacAlpine DM, Malc EP, et al. Heterogeneous polymerase fidelity and mismatch repair bias genome variation and composition. *Genome Res* 2014;24(11):1751–64.
- [96] Frolova AI, Babb SA, Zantow E, Hagemann AR, Powell MA, Thaker PH, et al. Impact of an immunohistochemistry-based universal screening protocol for Lynch syndrome in endometrial cancer on genetic counseling and testing. *Gynecol Oncol* 2015;137(1):7–13.
- [97] Ouyang KJ, Woo LL, Zhu J, Huo D, Matunis MJ, Ellis, NA. SUMO Modification Regulates BLM and RAD51 Interaction at Damaged Replication Forks. *PLoS Biol* 2011; 7 (12): e1000252.
- [98] Ayed F-BF, Douira-Khoms W, Rhaymen S, Jelassi M, Zribi H, Chaadouni M, et al. Burkitt lymphoma in a child with Bloom syndrome. *Arch Pédiat* 2016; 23 (4): 382–384.
- [99] Guo R-B, Rigolet P, Ren H, Zhang B, Zhang X-D, Dou S-X, et al. Structural and functional analyses of disease-causing missense mutations in Bloom syndrome protein. *Nucleic Acids Res* 2007;35(18):6297–310.
- [100] Rio Frio T, Lavoie J, Hamel N, Geyer FC, Kushner YB, Novak DJ, et al. Homozygous BUB1B Mutation and Susceptibility to Gastrointestinal Neoplasia. *N Engl J Med* 2010;363(27):2628–37.
- [101] Vleugel M, Hoek TA, Tromer E, Slidrecht T, Groenewold V, Omerzu M, et al. Dissecting the roles of human BUB1 in the spindle assembly checkpoint. *J Cell Sci* 2015;128(16):2975–82.
- [102] García-Castillo H, Vázquez-Velásquez AI, Rivera H, Barros-Núñez P. *Am J Med Genet A* 2008;146A(13):1687–95.
- [103] Shi Q, Hu M, Luo M, Liu Q, Jiang F, Zhang, Y, Wang S, et al. Reduced expression of Mad2 and Bub1 proteins is associated with spontaneous miscarriages. *Mol Hum Reprod* 2010; 17 (1): 14–21.
- [104] Micale MA, Schran D, Emch S, Kurczynski TW, Rahman N, Dyke DLV. Mosaic variegated aneuploidy without microcephaly: Implications for cytogenetic diagnosis. *Am J Med Genet A* 2007;143(16):1890–2183.
- [105] Lin SM, Luk HM, Lo IFM, Tam W-K, Chan KYK, Tse H-Y, et al. Prenatal diagnosis and long-term follow-up of a Chinese patient with mosaic variegated aneuploidy and its molecular analysis. *Clin Case Rep* 2020;8(8):1369–75.
- [106] Baker DJ, Jegannathan KB, Cameron JD, Thompson M, Juneja S, Kopecka A, et al. BubR1 insufficiency causes early onset of aging-associated phenotypes and infertility in mice. *Nat Genet* 2004;36(7):744–9.
- [107] He R, Wu Q, Zhou H, Huang N, Chen J, Teng J. Cep57 Protein Is Required for Cytokinesis by Facilitating Central Spindle Microtubule Organization. *J Biol Chem* 2013;288(20):14384–90.
- [108] Watanabe K, Takao D, Ito KK, Takahashi M, Kitagawa D. The Cep57-pericentriole module organizes PCM expansion and centriole engagement. *Nat Commun* 2019; 10(1). <https://doi.org/10.1038/s41467-019-08862-2>.
- [109] LaRocque JR, Stark JM, Oh J, Bojilova E, Yusa K, Horie K, et al. Interhomolog recombination and loss of heterozygosity in wild-type and Bloom syndrome helicase (BLM)-deficient mammalian cells. *Proc Natl Acad Sci U S A* 2011;108 (29):11971–6.
- [110] Calin G, Ranzani GN, Amadori D, Herlea V, Matei I, Barbanti-Brodano G, et al. Somatic frameshift mutations in the Bloom syndrome BLM gene are frequent in sporadic gastric carcinomas with microsatellite mutator phenotype. *BMC Genet* 2001;2(14). <https://doi.org/10.1186/1471-2156-2-14>.
- [111] Kaneko H, Fukao T, Kondo N. The function of RecQ helicase gene family (especially BLM) in DNA recombination and joining. *Adv Biophys* 2004;38:45–64.
- [112] Ham MF, Takakuwa T, Luo W-J, Liu A, Horii A, Aozasa K. Impairment of double-strand breaks repair and aberrant splicing of ATM and MRE11 in leukemia-lymphoma cell lines with microsatellite instability. *Cancer Sci* 2006;97 (3):226–34.
- [113] Bhisitkul RB, Rizen M. Bloom syndrome: multiple retinopathies in a chromosome breakage disorder. *Br J Ophthalmol* 2001;88(3). <https://doi.org/10.1136/bjo.2002.011643>.
- [114] Diaz A, Vogiatzi MG, Sanz MM, German J. Evaluation of Short Stature, Carbohydrate Metabolism and Other Endocrinopathies in Bloom's Syndrome. *Horm Res* 2006;66(3):111–7.
- [115] Tayebi N, Khodaei H. Bloom's syndrome in a 12-year-old Iranian girl. *Indian J Hum Genet* 2008;14(3):103–5.
- [116] Adams M. Acute Myeloid Leukaemia after Treatment for Acute Lymphoblastic Leukaemia in Girl with Bloom Syndrome. *J Genet Syndr Gene Ther* 2013;04(08).
- [117] Mulcahy MT, French M. Pregnancy in Bloom's Syndrome. *Clin Genet* 1981;19(3): 156–218.
- [118] Chisholm CA, Bray MJ, Karns LB. Successful pregnancy in a woman with Bloom syndrome. *Am J Genet* 2001;102(2):136–218.
- [119] German J. Bloom syndrome: a mendelian prototype of somatic mutational disease. *Medicine* 1993;72(6):393–406.
- [120] El Ghamrasni S, Cardoso R, Halaby MJ, Zeegers D, Harding S, Kumareswaran R, et al. Cooperation of BLM and Mus81 in development, fertility, genomic integrity and cancer suppression. *Oncogene* 2015;34(14):1780–9.
- [121] Sonoda E, Sasaki MS, Morrison C, Yamaguchi-Iwai Y, Takata M, Takeda S. Sister Chromatid Exchanges Are Mediated by Homologous Recombination in Vertebrate Cells. *Mol Cell Biol* 1999;19(7):5166–519.
- [122] Howell RT, Davies T. Diagnosis of Bloom's syndrome by sister chromatid exchange evaluation in chorionic villus cultures. *Prenat Diagn* 1994;14(11): 1071–103.
- [123] Londoño-Vallejo AJ, Der-Sarkissian H, Cazes L, Bacchetti S, Reddel RR. Alternative Lengthening of Telomeres Is Characterized by High Rates of Telomeric Exchange. *Cancer Res* 2004;64(7):2324–3237.
- [124] Du X, Shen J, Kugan N, Furth EE, Lombard DB, Cheung C, et al. Telomere Shortening Exposes Functions for the Mouse Werner and Bloom Syndrome Genes. *Mol Cell Biol* 2004;24(19):8437–46.
- [125] Takagi S, Kinouchi Y, Hiwatashi N, Nagashima F, Chida M, Takahashi S, et al. Relationship between microsatellite instability and telomere shortening in colorectal cancer. *Dis Colon Rectum* 2000;43(10):S12–7.
- [126] Fang D-C, Yang S-M, Zhou X-D, Wang D-X, Luo Y-H. Telomere erosion is independent of microsatellite instability but related to loss of heterozygosity in gastric cancer. *World J Gastroenterol* 2001;7(4):522–56.
- [127] Campbell MR, Wang Y, Andrew SE, Liu Y. *Msh2* deficiency leads to chromosomal abnormalities, centrosome amplification, and telomere capping defect. *Oncogene* 2006;2006(25):2531–3256.
- [128] Rampazzo E, Bertorelle R, Serra L, Terrin L, Candiotto C, Pucciarelli S, et al. Relationship between telomere shortening, genetic instability, and site of tumour origin in colorectal cancers. *Br J Cancer* 2010;102(8):1300–5.
- [129] Lee PP, Fitzpatrick DR, Beard C, Jessup HK, Lehar S, Makar KW, et al. A Critical Role for Dnmt1 and DNA Methylation in T Cell Development, Function, and Survival. *Immunity* 2001;15(5):763–74.
- [130] Etoh T, Kanai Y, Ushijima S, Nakagawa T, Nakanishi Y, Sasako M, et al. Increased DNA Methyltransferase 1 (DNMT1) Protein Expression Correlates Significantly with Poorer Tumor Differentiation and Frequent DNA Hypermethylation of Multiple CpG Islands in Gastric Cancers. *Am J Pathol* 2004;164(2):689–99.
- [131] Watt F, Molloy PL. Cytosine methylation prevents binding to DNA of a HeLa cell transcription factor required for optimal expression of the adenovirus major late promoter. *Genes Dev* 1988;2(9):1136–43.
- [132] Boyes J, Bird A. Repression of genes by DNA methylation depends on CpG density and promoter strength: evidence for involvement of a methyl-CpG binding protein. *EMBO J* 1992;11(1):327–33.
- [133] Clark SJ, Harrison J, Molloy PL. Sp1 binding is inhibited by <sup>m</sup>Cp<sup>m</sup>CpG methylation. *Gene* 1997;195(1):67–71.



- [134] Kim M, Trinh BN, Long TI, Oghamian S, Laid PW. Dnmt1 deficiency leads to enhanced microsatellite instability in mouse embryonic stem cells. *Nucleic Acids Res* 2004;32(19):5742–5759.
- [135] Foucault F, Buard J, Praz F, Jaulin C, Stoppa-Lyonnet D, Vergnaud G, et al. Stability of microsatellites and minisatellites in Bloom syndrome, a human syndrome of genetic instability. *Mutat Res* 1996;362(3):227–36.
- [136] Kaneko H, Inoue R, Fukao T, Kasahara K, Tashita H, Teramoto T, et al. Two Japanese siblings with Bloom syndrome gene mutation and B-cell lymphoma. *Leuk Lymphoma* 1997;27(5-6):539–42.
- [137] Harkness EF, Barrow E, Newton K, Green K, Clancy T, Lalloo F, et al. Lynch syndrome caused by MLH1 mutations is associated with an increased risk of breast cancer: a cohort study. *J Med Genet* 2015;52(8):553–6.
- [138] Moreno-Ortiz JM, Ayala-Madrigal MdL, Corona-Rivera JR, Centeno-Flores M, Maciel-Gutiérrez V, Franco-Topete RA, et al. Novel Mutations in MLH1 and MSH2 Genes in Mexican Patients with Lynch Syndrome. *Gastroenterol Res Pract* 2016;2016:1–6.
- [139] Senter L, Clendenning M, Sotamaa K, Hampel H, Green J, Potter JD, et al. The Clinical Phenotype of Lynch Syndrome Due to Germ-Line PMS2 Mutations. *Gastroenterology* 2008;135(2):419–428.e1.
- [140] Walsh MD, Buchanan DD, Cummings MC, Pearson S-A, Arnold ST, Clendenning M, et al. Lynch Syndrome-Associated Breast Cancers: Clinicopathological Characteristics of a Case Series from the Colon CFR. *Clin Cancer Res* 2011;16(7):2214–24.
- [141] Möller P, Seppälä T, Bernstein I, Holinski-Feder E, Sala P, Evans GD, et al. Patients receiving colonoscopic and gynaecological surveillance: first report from the prospective Lynch syndrome database. *Gut* 2015;66(3). <https://doi.org/10.1136/gutjnl-2015-309675>.
- [142] Bauer CM, Ray AM, Halstead-Nussloch BA, Dekker RG, Raymond VM, Gruber SB, et al. Hereditary Prostate Cancer as a Feature of Lynch Syndrome. *Fam Cancer* 2011;10(1):37–42.
- [143] Lobo J, Pinto C, Pinheiro M, Lobo F, Sousa N, Lopes P, et al. Widening the spectrum of Lynch syndrome: first report of testicular seminoma attributable to MSH2 loss. *Histopathology* 2020;76(3):486–9.
- [144] Stupart D, Win AK, Winship IM, Jenkins M. Fertility after young onset colorectal cancer: a study of subjects with Lynch syndrome. *Colorectal Dis* 2016;17(9):787–93.
- [145] Ji G, Long Y, Zhou Y, Huang C, Gu A, Wang X. Common variants in mismatch repair genes associated with increased risk of sperm DNA damage and male infertility. *BMC Med* 2012;10(1). <https://doi.org/10.1186/1741-7015-10-49>.
- [146] Zhang X, Ding M, Ding X, Li T, Chen H. Six polymorphisms in genes involved in DNA double-strand break repair and chromosome synapsis: association with male infertility. *Syst Biol Reprod Med* 2015;61(4):187–93.
- [147] Zhao X, Mu C, Ma J, Dai X, Jiao H. The association of four SNPs in DNA mismatch repair genes with idiopathic male infertility in northwest China. *Int J Immunogenet* 2019;46(6):451–548.
- [148] Kadhim ZA, Abdul-Hassan IA. The Association of Mismatch Repair (MLH1) Gene Polymorphism with The Risk of Oligozoospermia in Iraqi Patients. *Iraqi J Biotechnol* 2020;19(1):56–68.
- [149] Horimoto Y, Thinzar Hlaing M, Saeki H, Kitano S, Nakai K, Sasaki R, et al. Microsatellite instability and mismatch repair protein expressions in lymphocyte-predominant breast cancer. *Cancer Sci* 2020;111(7):2647–54.
- [150] Watatani M, Yoshida T, Kuroda K, Ieda S, Yasutomi M. Allelic Loss of Chromosome 17p, Mutation of the p53 Gene, and Microsatellite Instability in Right- and Left- Sided Colorectal Cancer. *Cancer* 1996;77(8):1688–93.
- [151] Watanabe M, Imai H, Kato H, Shiraishi T, Ushijima T, Nagao M, et al. Microsatellite instability in latent prostate cancers. *Int J Cancer* 1996;69(5):394–7.
- [152] Kiaris H, Koumantakis E, Ergazaki M, Sifakis S, Spandidos D. Instability at Microsatellite Sequences in Spontaneously Aborted Human Embryos Provides Evidence for a Novel Mechanism for Recurrent Miscarriages. *Oncol Rep* 1995;2(5):805–9.
- [153] Tsai AF, Kaufman KA, Walker MA, Karrison TG, Odem RR, Barnes RB, et al. Transmission disequilibrium of maternally-inherited CTLA-4 microsatellite alleles in idiopathic recurrent miscarriage. *J Reprod Immunol* 1998;40(2):147–57.
- [154] Spandidos DA, Koumantakis E, Sifakis S, Sourvinos G. Microsatellite mutations in spontaneously aborted embryos. *Fertil Steril* 1998;70(5):892–985.
- [155] Wang L, Wang ZC, Xie C, Liu XF, Yang MS. Genome-Wide Screening for Risk Loci of Idiopathic Recurrent Miscarriage in a Han Chinese Population: A Pilot Study. *Reprod Sci* 2010;17(6):578–84.
- [156] Shukla V, Høffding MK, Hoffman ER. Genome diversity and instability in human germ cells and preimplantation embryos. *Semin Cell Dev Biol* 2021;113:132–47.
- [157] Cui X, Zhang J, Du R, Wang L, Archacki S, Zhang Y, et al. HSF4 is involved in DNA damage repair through regulation of Rad51. *Biochim Biophys Acta* 2012;1822(8):1308–15.
- [158] Guo XJ, P-h Ni, Deng W-w. Genetic alterations of microsatellite markers at chromosome 17 in non-small cell lung cancer. *Chin. J Cancer Res* 2001;13(3):181–4.
- [159] Samara KD, Tzortzaki EG, Neofytou E, Karatzanis AD, Lambiri I, Tzanakis N, et al. Somatic DNA alterations in lung epithelial barrier cells in COPD patients. *Pul Pharmacol Ther* 2010;23(3):208–14.
- [160] Thomou C, Paraskakis E, Neofytou E, Kalmanti M, Sifakis NM, Tzortzaki EG. Acquired somatic mutations in the microsatellite DNA, in children with bronchial asthma. *Pediatr Pulmonol* 2009;44(10):1017–24.
- [161] Hou TZ, Qureshi OS, Wang CJ, Baker J, Young SP, Walker LSK, et al. A Transendocytosis Model of CTLA-4 Function Predicts Its Suppressive Behavior on Regulatory T Cells. *J Immunol* 2015;194(5):2148–59.
- [162] Ovcinnikovs V, Ross EM, Petersone L, Edner NM, Heuts F, Ntavli E, et al. CTLA-4-mediated transendocytosis of costimulatory molecules primarily targets migratory dendritic cells. *Sci Immunol* 2019;4(35). <https://doi.org/10.1126/sciimmunol.aaw0902>.
- [163] Misra MK, Mishra AM, Phadke SR, Agrawal S. Association of functional genetic variants of CTLA4 with reduced serum CTLA4 protein levels and increased risk of idiopathic recurrent miscarriages. *Fertil Steril* 2016;106(5):1115–23.
- [164] Ding J-L, Diao L-H, Yin T-L, Huang C-Y, Yin B, Chen C et al. Aberrant expressions of endometrial Id3 and CTLA-4 are associated with unexplained repeated implantation failure and recurrent miscarriage. *Am J Reprod Immunol* 2017; 78(2): e12632.
- [165] Vento-Tormo R, Efremova M, Botting RA, Turco MY, Vento-Tormo M, Meyer KB, et al. Single-cell reconstruction of the early maternal-fetal interface in humans. *Nature* 2018;563(7731):347–53.
- [166] Guo C, Cai P, Jin L, Sha Q, Yu Q, Zhang W, et al. Single-cell profiling of the human decidual immune microenvironment in patients with recurrent pregnancy loss. *Cell Discov* 2021;7(1). <https://doi.org/10.1038/s41421-020-00236-z>.
- [167] Thomas DC, Umar A, Kunkel TA. Microsatellite instability and mismatch repair defects in cancer cells. *Mut Res* 1996;350(1):201–5.
- [168] Iino H, Simms L, Young J, Arnold YJ, Winship IM, Webb SI, et al. DNA microsatellite instability and mismatch repair protein loss in adenomas presenting in hereditary non-polyposis colorectal cancer. *Gut* 2000;47(1):37–42.
- [169] Diogo F, Berta C, Ana N, Silva J, Póva AM, Barros A, et al. Preimplantation genetic testing for Huntington disease: the perspective of one Portuguese center. *Porto Biomed J* 2019;4(5):e48.
- [170] Ou Z, Chen Z, Yin M, Deng Yu, Liang Y, Wang W, et al. Re-analysis of whole blastocysts after trophectoderm biopsy indicated chromosome aneuploidy. *Hum Genomics* 2020;14(1). <https://doi.org/10.1186/s40246-019-0253-z>.
- [171] Huang L, Bogale B, Tang Y, Lu S, Xie XS, Racowsky C. Noninvasive preimplantation genetic testing for aneuploidy in spent medium may be more reliable than trophectoderm biopsy. *Proc Natl Acad Sci U S A* 2019;116(28):14105–12.