

Lead Exposure in Adult Males in Urban Transvaal Province, South Africa during the Apartheid Era

Catherine A. Hess¹, Matthew J. Cooper², Martin J. Smith¹, Clive N. Trueman², Holger Schutkowski^{1*}

1 School of Applied Sciences, Bournemouth University, Talbot Campus, Poole, United Kingdom, **2** Ocean and Earth Sciences, National Oceanography Centre, University of Southampton, Southampton, United Kingdom

Abstract

Human exposure to lead is a substantial public health hazard worldwide and is particularly problematic in the Republic of South Africa given the country's late cessation of leaded petrol. Lead exposure is associated with a number of serious health issues and diseases including developmental and cognitive deficiency, hypertension and heart disease. Understanding the distribution of lifetime lead burden within a given population is critical for reducing exposure rates. Femoral bone from 101 deceased adult males living in urban Transvaal Province (now Gauteng Province), South Africa between 1960 and 1998 were analyzed for lead concentration by Inductively Coupled Plasma Mass Spectrometry (ICP-MS). Of the 72 black and 29 white individuals sampled, chronic lead exposure was apparent in nearly all individuals. White males showed significantly higher median bone lead concentration ($ME = 10.04 \mu\text{g}\cdot\text{g}^{-1}$), than black males ($ME = 3.80 \mu\text{g}\cdot\text{g}^{-1}$) despite higher socioeconomic status. Bone lead concentration covaries significantly, though weakly, with individual age. There was no significant temporal trend in bone lead concentration. These results indicate that long-term low to moderate lead exposure is the historical norm among South African males. Unexpectedly, this research indicates that white males in the sample population were more highly exposed to lead.

Citation: Hess CA, Cooper MJ, Smith MJ, Trueman CN, Schutkowski H (2013) Lead Exposure in Adult Males in Urban Transvaal Province, South Africa, during the Apartheid Era. PLoS ONE 8(3): e58146. doi:10.1371/journal.pone.0058146

Editor: Jaymie Meliker, Stony Brook University, Graduate Program in Public Health, United States of America

Received: October 13, 2012; **Accepted:** January 30, 2013; **Published:** March 7, 2013

Copyright: © 2013 Hess et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The research was funded, in part, by the University of Bournemouth Post Graduate Global Exchange Fund, the Bournemouth University School of Applied Sciences Hilary Williams Fund, and the British Council Overseas Student Research Award Scheme. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: hschutkowski@bournemouth.ac.uk

Introduction

Population-wide exposure to lead pollution is a problem that has, for the most part, been addressed in Europe and North America. However inorganic environmental pollution, specifically that of lead and other toxic heavy metals is a major public health concern in sub-Saharan Africa [1–5]. Lead in particular, is of growing concern because of its known toxicity at low levels. The problem of environmental lead pollution was largely overlooked in South Africa in the 20th century, despite its substantial mining and industrial activities, and perhaps more surprisingly, despite the country's persistent use of leaded petrol [1,6]. Few studies of human lead exposure in Transvaal were published prior to the formation of the New South African Republic in 1994, which has left a prominent lack of baseline data with which to compare the growing body of public health research into the issue. Among the aims of this research is to address this gap in data through analysis of the skeletal remains of South African individuals who died before 1998. The Pretoria Bone collection, from which the study population derives, contains fully identified individuals and its use in this research comprises an unparalleled opportunity to study recent historical trends in human lead exposure. In addition, bone tissue is an endogenous repository for lead. Due to its low turnover rate compared to other human tissues – approximately 10 years for compact bone – bone lead concentration is an excellent indicator of chronic lead exposure [7]. Within Africa, the highly industrial Gauteng Province is among the more polluted regions.

The region forms the backbone of the industrial and mining economy of South Africa and is home to the country's most industrial city, Johannesburg and its capital, Pretoria. Urban pollution is a significant public health concern as is human exposure to lead [4,5]. The results of this research may provide valuable background information to more recent studies involving human blood lead concentration in the region.

South Africa began monitoring lead exposure in children in the 1980s [8–11]. Studies conducted by von Schirnding et al. found that as many as 13 percent of children living in Cape Town had blood lead levels greater than or equal to 25 $\mu\text{g}/\text{dL}$ – more than twice the threshold considered dangerous by the US Centers for Disease Control – and noted that proximity to traffic was a significant risk factor for elevated blood lead, as was lower socioeconomic status, overcrowding and homes in disrepair [10–12]. Deveaux et al. also conducted blood lead monitoring in young children in Cape Town and found that children whose blood lead was greater than or equal to 29 $\mu\text{g}/\text{dL}$ were also living in homes with leaded paint [9].

Analysis of teeth from individuals buried in Cape Town before the introduction of leaded petrol show higher than expected lead concentrations which were also significantly higher than those measured in the mid-1980s and before the reduction in the lead concentration of petrol [13]. It was determined that the prevalent use of lead pipes in residential plumbing was responsible. To date only one study of bone lead has been conducted in South Africa. Todd et al. measured tibia lead in employees of a lead-acid battery

factory [14]. They report a mean bone lead concentration of $53.4 \mu\text{g}\cdot\text{g}^{-1}$.

Despite these early studies, as late as 2005 the country had no national lead monitoring program [15]. In addition, we could find no studies of lead exposure conducted in Transvaal during the apartheid era, leaving a gap in the understanding of the historical and demographic patterns associated with lead exposure. In addition, because of the cumulative nature of bone lead, this measure is widely considered to be a valuable indicator of chronic, as opposed to acute, lead exposure, and from an epidemiological standpoint, may be a more reliable indicator of demographic and long-term exposure patterns than blood lead [7,16–18]. In light of these observations, this study aims to quantify lead exposure among urban South African males during apartheid by measuring bone lead concentration in an identified skeletal collection.

The authors wish to note that the racial terms “black” and “white” are used in this study to denote ethnic ancestry. This is wholly due to the fact that the sampled population is classified in this way in associated cadaver records and because the population would have been segregated purely by racial classification during the time period being studied. These terms have social, demographic and political connotations the implications of which appear to have influenced patterns of lead exposure within the study population.

Materials

Skeletal material was sampled from the Pretoria Identified Bone Collection at the University of Pretoria, South Africa and the Dart Student Bone Collection at Witwatersrand University, Johannesburg. The Pretoria Bone Collection is an identified reference collection held at the University of Pretoria, School of Medicine. The skeletal remains are those of individuals who died in the Pretoria area between 1943 and 2012 and whose bodies were either unclaimed or donated. In the former case, unclaimed bodies become the property of the University of Pretoria to be used for teaching and research, subject to the South Africa Human Tissues Act of 1983 [19]. The collection consists of individuals who range in age from neonates to 95 years of age. The predominant demographic within the collection is black males. This is largely to do with both overall demographic patterns within South Africa and to economic conditions during Apartheid, in which circulating migration brought black males to urban areas from Bantustans for work [19–21]. No information regarding the occupation of any of the individuals in either collection was available. The Raymond Dart Collection is housed at the University of Witwatersrand, School of Medicine and is similar in demographic composition to the Pretoria Collection. Skeletal remains in the Dart collection date to 1928 [22]. Only 12 of the femora included in this study are from the Dart collection. For both collections, ancestry was determined by the admitting hospital and based on the racial classification set forth in the 1950 Population Registration Act, which categorized individuals as black, white or colored based on physical appearance, parentage (an individual with one white and one black or colored parent could not be classified as white) and socio-cultural considerations. For the purpose of this paper, these classifications were not re-examined, as this research is primarily concerned with the way this racial division would have contributed to different lead exposure rates. In addition, because of the unique lack of fluidity between racial groups imposed by Apartheid, and because these groups largely defined socio-economic status at the time, the two factors are considered one and the same in this instance.

Table 1. Pb concentration and recovery rate for NIST 1486 Bone Meal.

Sample	Pb $\mu\text{g}\cdot\text{g}^{-1}$	% Recovery
NIST 1486	1.33±.014	–
CRM 1	1.166	87.2
CRM2	1.034	78.0
CRM3	1.167	87.7
CRM4	1.217	92.0
CRM5	1.344	101.0
CRM7	1.229	92.4
CRM 8	1.248	75.1
CRM 10	1.259	95.0
Avg.		88.5

doi:10.1371/journal.pone.0058146.t001

Ethics Statement

This research was approved by the Bournemouth University Ethical Review Committee and the University of Pretoria, Department of Anatomy. In addition, the project met the requirements set by the UK Human Tissues Act (1994) and bone samples were imported into the UK and analyzed in accordance with the Act.

Table 2. Median bone Pb ($\mu\text{g}\cdot\text{g}^{-1}$) in black and white urban South African males in relation to age.

Race	N	Pb	SD	IRQ	Min.	Max.
Black	72	3.92	5.69	4.12	1.22	32.23
White	29	10.04	13.61	9.58	1.55	64.09
Black						
Age	20–29	9	2.22	2.7	1.7	9.25
	30–39	12	4.14	3.47	1.87	13.56
	40–49	18	3.3	2.48	1.22	11.56
	50–59	12	3.67	5.1	1.61	18.1
	60–69	11	4.53	1.72	1.9	6.73
	70–79	10	7.2	11.66	2.02	32.23
	80–89	1	12.95			
Median	49				18	80
White						
	20–29	0				
	30–39	0				
	40–49	5	10.04	7.9	7.53	26.54
	50–59	7	10.85	7.02	6.38	27.6
	60–69	9	12.7	10.7	2.78	37
	70–79	3	3.41	12.56	1.55	24.18
	80–89	3	7.59	0.22	7.45	7.88
	90–99	2	49.07	21.25	34.04	64.1
Median	62				42	95

doi:10.1371/journal.pone.0058146.t002

Table 3. Total bone lead concentration in $\mu\text{g}\cdot\text{g}^{-1}$ dry weight in femora of black males.

Specimen	Age	Death	Pb	Specimen	Age	Death	Pb
	(years)	(year)	$\mu\text{g}\cdot\text{g}^{-1}$		(years)	(year)	$\mu\text{g}\cdot\text{g}^{-1}$
2	30	1987	4.26	134	20	1964	1.80
3	38	1988	6.60	137	50	1965	14.71
4	36	1988	7.89	141	40	1961	2.19
5	67	1991	2.23	143	49	1966	4.49
6	40	1988	3.92	144	70	1969	7.63
7	30	1987	7.93	147	42	1979	2.72*
8	65	1985	4.76	148	48	1975	1.95
9	50	1988	3.45	150	61	1983	4.17
10	59	1987	18.05	151	65	1972	6.19
11	75	1988	32.23	152	50	1979	3.60
12	51	1991	6.14	156	55	1972	1.61
20	35	1988	2.51	158	50	1972	8.79
23	50	1983	2.95*	159	60	1982	4.53
29	30	1984	2.44	168	70	1979	6.74
48	30	1970	13.56	169	69	1983	1.85
51	58	1975	3.65	174	65	1979	6.73
61	56	1983	3.37	192	35	1967	1.87
63	48	1967	7.65	198	27	1964	6.00
64	40	1967	2.98	199	35	1966	2.48
79	45	1979	5.52	300	44	1979	3.53
83	40	1972	2.96	301	65	1965	2.72
86	44	1979	2.05	306	49	1976	3.05
88	24	1967	9.25	312	25	1966	2.13
89	80	1970	12.95	313	66	1967	6.02
90	70	1966	8.56	314	70	1966	15.66
92	56	1979	5.81	315	26	1972	3.08
95	47	1963	1.91	317	60	1983	4.95
99	40	1965	4.37	319	49	1967	11.56
101	48	1969	1.22	320	37	1966	4.36*
104	26	1966	1.69	321	43	1967	5.76
113	20	1979	2.18*	325	72	1979	2.99
115	70	1979	2.02	326	73	1980	32.13
121	40	1965	3.83	329	70	1983	3.93
123	59	1964	3.69	333	28	1965	6.46
125	70	1973	2.25	334	60	1982	2.38*
131	34	1970	4.03	335	18	1982	2.22

*Denotes averaged Pb concentration between right and left femora.

doi:10.1371/journal.pone.0058146.t003

Methods

Analytical Methods

Cortical bone samples of approximately 0.250 g were removed from the right or left femora of 101 individuals who lived in Gauteng Province at the time of their death between 1961 and 1998. Bone samples were removed from femora with a 10 mm diamond-tipped core drill attached to a drill press. Cores were taken from the posterior-distal surface of the right or left femur, just above the intercondylar fossa and placed into sealed plastic bags until analysis. Due to the demographic composition of the

collections which are biased heavily towards black males, the remains sampled were primarily black males. Analysis was conducted at the University of Southampton Geochemistry Class 100 Clean laboratory at the National Oceanography Centre Southampton. All reagents used were Fisher Trace Element grade and further sub-boiled in Teflon® stills to ensure ultra-purity. Water used was MilliQ® Millipore ultra-pure water (18.2 M Ω).

Sample Preparation

Samples were weighed, washed three times with MilliQ® water to remove any surface contaminants and placed into acid-washed

Table 4. Total bone lead concentration in $\mu\text{g}\cdot\text{g}^{-1}$ dry weight in femora of white males.

Specimen	Age	Death	Pb	Specimen	Age	Death	Pb
	(years)	(year)	$\mu\text{g}\cdot\text{g}^{-1}$		(years)	(year)	$\mu\text{g}\cdot\text{g}^{-1}$
17	67	1983	15.18	124	83	1993	7.59
39	62	1980	9.70	126	56	1982	8.30
59	84	1998	7.45	178	47	1977	7.53
60	71	1977	24.18	183	62	1983	12.70*
74	72	1998	3.40	185	68	1975	2.95
78	52	1983	6.38	190	66	1997	3.91
82	95	1982	64.09	191	74	1972	1.55
84	82	1997	7.88*	195	60	1973	2.78*
85	50	1976	13.26	295	44	1977	12.30
93	56	1979	9.34*	298	67	1976	37.00
94	42	1977	10.04	305	68	1964	18.44*
105	43	1964	26.54	322	69	1984	15.69
116	59	1982	10.85	324	57	1976	27.58
119	56	1982	12.15*	332	48	1973	7.82
120	91	1979	34.04				

*Denotes averaged Pb concentration between right and left femora.

doi:10.1371/journal.pone.0058146.t004

13 mL polyethylene tubes. 1 mL of concentrated, sub-boiled HNO_3 (69%) was added to each tube and left at room temperature for 72 hours. After initial digestion, 9 mL MilliQ[®] was added to each tube and samples were left to digest at room temperature (approx. 20°C) for a further 72 hours. To facilitate ICP-MS analysis, all samples were diluted to approximately 100 $\mu\text{g}\cdot\text{g}^{-1}$ calcium concentration with 3% sub-boiled HNO_3 .

Sample Analysis

Samples were analyzed by ICP-MS (Thermo Scientific XSeries 2) calibrated with synthetic mixed element standards made from single element ICP-MS standards (Inorganic Ventures). All samples and standards contained 20 $\text{ng}\cdot\text{g}^{-1}$ Be and 5 $\text{ng}\cdot\text{g}^{-1}$ In and Re as internal standards. The elements were analyzed in one of two instrument modes depending on signal size and susceptibility to interferences. These were standard mode and CCT mode with 2 mL/min. of a mixed He/H₂ gas added to reduce interferences. Ten reagent blanks of 3% HNO_3 were analyzed and Pb concentration in all blanks was below the limit of detection. Detection limit for Pb is 0.0004 $\mu\text{g}/\text{L}^{-1}$. Method validation was established by the inclusion of ten, 0.1 g samples of NIST SRM 1486 Bone Meal and Pb concentration is reported in Table 1. Mean Pb recovery rate in CRM was 90%. Sample duplicate precision was measured at 0.82 ($SD=0.32$).

Statistical Methods

Kolmogorov-Smirnov tests confirmed that bone Pb concentration was not normally distributed for either black males, $D(74)=0.255$, $p<0.001$ or white males $D(29)=0.277$, $p=0.001$. Pb concentration data was log transformed and was found to be normally distributed with $D(74)=0.084$, $p>0.05$ and $D(29)=0.133$, $p>0.05$, in black and white males respectively. Independent t-tests, ANCOVA and multiple regression were performed on log-transformed data. All Pb concentrations reported are back-transformed values.

Results

Median bone Pb by race and age group are presented in Table 2. There was a high degree of variability within the subject population as a whole. Of 72 black males, the median Pb concentration is 3.80 $\mu\text{g}\cdot\text{g}^{-1}$. For the 29 white males median Pb concentration is 10.04 $\mu\text{g}\cdot\text{g}^{-1}$. Results of Pb concentration for black males are presented in Table 3 and white males in Table 4. In some cases, samples from both right and left femora were taken from the same individual, in these cases Pb concentration in both femora within a single individual was averaged and is indicated by an asterisk.

White males show significantly higher bone Pb concentration than black males (Fig. 1). An independent t-test confirmed that the difference in means is significant, $t(100)=5.5$, $p<0.001$. Among all samples, the highest individual concentrations occur in white




Figure 1. Boxplot of median bone Pb concentrations (in $\mu\text{g}\cdot\text{g}^{-1}$) in black and white males. White males show significantly higher bone Pb concentration than black males. Horizontal line = median, boxes = 2nd and 3rd quartiles, error bars = range.

doi:10.1371/journal.pone.0058146.g001

males – samples 82 ($64.09 \mu\text{g}\cdot\text{g}^{-1}$) and 60 ($24.8 \mu\text{g}\cdot\text{g}^{-1}$). Among black males, the highest concentrations occur in samples 11 ($32.23 \mu\text{g}\cdot\text{g}^{-1}$) and 10 ($18.05 \mu\text{g}\cdot\text{g}^{-1}$).

There were significant difference in bone Pb concentrations between the 12 black males from the Dart Collection, and black males in the Pretoria Collection, $t(71)=2.23$, $p<0.05$. Median bone Pb from males in the Dart collection is $6.14 \mu\text{g}\cdot\text{g}^{-1}$ and $3.36 \mu\text{g}\cdot\text{g}^{-1}$ in males from the Pretoria Collection. However this result must be accepted with caution, as there are only 12 samples from the Dart collection and there is considerable overlap in the admitting hospitals between the two collections, indicating that the individual remains in the two collections did not come from two distinct populations.

It is possible that some of the apparent differences in bone lead concentrations between black and white origin males could be attributed to age, as the white individuals are on average older than the black individuals (Table 2). Analysis of Covariance (ANCOVA) demonstrates that the covariate, age affects bone Pb concentration, $F(1, 99)=6.66$, $p<0.05$, though the effect size is small, $r=0.06$. After controlling for the effect of age, the effect of ancestry on bone Pb concentration remains significant, $F(1, 99)=19.20$, $p<0.001$, though the effect size is relatively small, $r=0.145$.

When multiple regression was used to test the relationship between age and ancestry, a significant linear trend between age, ancestry and bone Pb concentration is apparent. Both age and ancestry explain 28.4% of the variance in bone Pb concentration ($R^2=.284$, $F(1, 102)=19.82$, $p<.001$). Both age and ancestry significantly predicts bone Pb ($\beta=.235$, $p<.001$ and $\beta=.395$, $p>.001$).

Discussion

Demographic Trends and Lead Exposure

The results of this study are particularly informative, in that they do not correspond to world-wide trends in human lead exposure. In general, and especially in developing countries, the poorest and most disadvantaged sectors within the population tend to encounter greater exposure to lead [23–25]. These groups also tend to yield the highest body concentrations of lead (bone or blood) [26,27]. This is the case in present day South Africa. von Schirnding et al. [10] have reported blood lead levels among children residing in Cape Town, with children from lower income households having the highest levels. Mathee et al. [28] reported that high blood lead levels were associated with lower socioeconomic status in a study of children in Johannesburg. Other researchers have reported similar findings [2,3].

The higher bone lead levels of white individuals reported in this study are interesting and may be the result of a variety of factors. The finding is counter to results of lead studies conducted in the United States. Research from the National Health and Nutrition Examination Surveys (NHANES) have consistently reported higher blood and bone lead concentrations in African American individuals regardless of age or sex [29,30]. Data from the US Veterans' Association Normative Aging Survey have also shown that white males tend to have lower bone and blood lead than African American males [31]. Similar results were found in the Baltimore Memory Study, in which authors reported significantly higher cortical bone lead in African American versus white males [32]. These patterns have persisted in the United States, even as overall lead exposure rates have fallen [33]. Most notably, Hu et al. [31] report a median bone lead concentration of $20 \mu\text{g}\cdot\text{g}^{-1}$ in community exposed males living in Boston. This is higher than the median concentrations values found in South African males in

this study. Hu et al. also report higher bone lead concentration in black males, and a significant increase in bone lead in males who did not complete high school versus those who completed graduate or professional school. The latter indicates a strong socioeconomic relationship with lead exposure.

With regards to our study, apartheid-mandated urban residential patterns, with white residents living closer to urban core and major roadways (with subsequent exposure to lead from petrol), may be the significant factors. Recent studies have reported greater atmospheric lead concentration in central business areas in Pretoria and Johannesburg, which during Apartheid were primarily white areas [34–37]. von Schirnding et al. found that atmospheric lead levels in the Cape Town city center were 2.5 times greater than in suburban areas [38]. The presence of lead paint in residential buildings and the possibility that homes built in the early 20th century and before may be plumbed with lead pipes is another, though the latter appears to be rare [39–42]. Investigation into the source of lead in bone by analysis of lead isotopic ratios is currently underway, which may shed light on this phenomenon.

Bone Lead and Age

This research confirms, though weakly, the previously reported association between age and bone lead concentration [43–45]. It has been estimated that 90% of the lead that is stored in the human body is stored in bone tissue [7,46,47]. This has the effect of sequestering lead from other tissues and organs where it may cause toxicity. However as individuals age and bone is resorbed, lead is released from bone tissue. The correlation between age and bone lead is well established and given the likelihood that bone acts as an endogenous source of lead within the body, releasing lead into the bloodstream as bone is resorbed and remodeled, it is clear that high lead levels in old age may have a significant impact on individual health [17].

Bone Lead and Public Health

With regards to toxicity, it has been previously reported that bone lead levels as low as $5 \mu\text{g}\cdot\text{g}^{-1}$ have been associated with clinical symptoms of toxicity such as hypertension [48–52]. In this study, 38% of black individuals and 86% of white individuals had bone lead levels above this threshold. Overall, however, the bone lead concentration in males in this study population is relatively moderate. Baranowska et al. [53] reported bone lead levels between 100 and $200 \mu\text{g}\cdot\text{g}^{-1}$ in an industrial district in Poland. Nevertheless, in the past decade it has become increasingly clear that chronic low-level exposure to lead is a substantial threat to individual and public health [54].

Reported health effects of chronic lead exposure include renal disease, diminished IQ and developmental delay (in children), and impaired cognitive function in adults [55–64]. Most recently, the drop in violent crime rates in urban areas in the United States has been attributed to the fall in lead pollution following the banning of lead in petrol [56,65,66]. Many of these pathologies are evident even at the subclinical level and at relatively low levels of exposure. Norman et al. [51] report that in South Africa in 2000, nearly 1,500 deaths could be attributed directly to lead exposure. Other studies have found that low-level lead exposure in men leads to diminished cognitive function on the order of five years accelerated mental aging [67]. From the results of this study, it is likely the negative effects of lead on public health have been acting on the population for some time. In addition, the data above suggest that persistent lower-level exposure to lead may be the norm in South Africa (even after the cessation of the use of lead in

petrol). This low but chronic level of exposure may be particularly pernicious, as subclinical or sub-acute symptoms are often overlooked in marginalised populations due, in part, to differential access to medical care and lifestyle [68–70]. Potentially then, despite lower lead exposure overall, black males may be more susceptible to unfavourable health effects.

It is critical to acknowledge that, though black individuals may show lower bone lead concentrations, the burden of disease resulting from lead may be higher in this demographic group. Numerous studies have demonstrated that individuals who may be physically or nutritionally stressed are also likely to suffer from the effects of lead toxicity at lower exposure levels than healthier individuals [71–73]. Within these populations lead exposure may also be associated with other illnesses such as asthma and iron deficiency anemia, both of which are prevalent in low income households in South Africa [74–78].

In summary, bone lead analysis of apartheid-era skeletal remains has yielded unexpected results. White males show

significantly higher bone lead concentration than black males. This difference could be attributed to use of exposure to leaded petrol and exacerbated by residential patterns in urban areas in which white individuals resided closer to the congested urban core.

Acknowledgments

The authors would like to thank Marius Loots and Ericka L'Abbe at the University of Pretoria for their generous assistance and access to the Pretoria Bone Collection. In addition, the authors would like to thank the curators of the Dart Bone Collection at Witwatersrand University for access to the Student Bone Collection.

Author Contributions

Conceived and designed the experiments: CAH MC CT HS MJS. Performed the experiments: CAH. Analyzed the data: CAH. Contributed reagents/materials/analysis tools: MC CT. Wrote the paper: CAH.

References

- Diab RD (1999) A note on changes in atmospheric lead content in seven cities in South Africa. *South African Journal of Science* 95: 117–121.
- Harper CC, Mathee A, von Schirnding Y, De Rosa CT, Falk H (2003) The health impact of environmental pollutants: a special focus on lead exposure in South Africa. *International Journal of Hygiene and Environmental Health* 206: 315–322.
- Nriagu JO (1992) Toxic metal pollution in Africa. *The Science of The Total Environment* 121: 1–37.
- Nriagu JO, Blankson ML, Ocran K (1996) Childhood lead poisoning in Africa: a growing public health problem. *Science of The Total Environment* 181: 93–100.
- Nweke OC, Sanders WH (2009) Modern Environmental Health Hazards: A Public Health Issue of Increasing Significance in Africa. *Environmental Health Perspectives* 117: 863–870.
- Mathee A, Rollin H, von Schirnding Y, Levin J, Naik I (2006) Reductions in blood lead levels among school children following the introduction of unleaded petrol in South Africa. *Environmental Research* 100: 319–322.
- Rabinowitz MB (1991) Toxicokinetics of bone lead. *Environmental Health Perspectives* 91: 33–37.
- White NW, Dempster WS, Pocock F, Kibel MA (1982) Lead absorption in Cape children - a preliminary report. *South African Medical Journal* 62: 799–802.
- Deveaux P, Kibel MA, Dempster WS, Pocock F, Formenti K (1986) Blood lead levels in pre-school children in Cape Town. *SAMJ-South African Medical Journal* 69: 421–424.
- von Schirnding Y (1991) Blood lead levels in South African inner-city children. *Environmental Health Perspectives* 94: 125–130.
- von Schirnding Y, Fuggle RF, Bradshaw D (1991) Factors associated with elevated blood lead levels in inner-city Cape Town children. *South African Medical Journal* 79: 454–456.
- Centers for Disease Control (2005) Preventing Lead Poisoning in Young Children. Atlanta: CDC.
- Grobler SR, Theunissen FS, Maresky LS (1996) Evidence of undue lead exposure in Cape Town before the advent of leaded petrol. *South African Journal of Medicine* 86: 169–171.
- Todd AC, Ehrlich RI, Selby P, Jordaan E (2000) Repeatability of Tibia Lead Measurement by X-Ray Fluorescence in a Battery-Making Workforce. *Environmental Research* 84: 282–289.
- Mathee A, von Schirnding Y, Montgomery M, Rollin H (2004) Lead poisoning in South African children: the hazard is at home. *Reviews on Environmental Health* 19: 347–361.
- Bergdahl IA, Skerfving S (2008) Biomonitoring of lead exposure-alternatives to blood. *Journal of Toxicology and Environmental Health-Part A-Current Issues* 71: 1235–1243.
- Hu H, Rabinowitz M, Smith D (1998) Bone lead as a biological marker in epidemiologic studies of chronic toxicity: Conceptual paradigms. *Environmental Health Perspectives* 106: 1–8.
- Hu H (1998) Bone lead as a new biologic marker of lead dose: Recent findings and implications for public health. *Environmental Health Perspectives* 106: 961–967.
- L'Abbe E, Loots M, Meiring JH (2005) The Pretoria Bone Collection: A modern South African Skeletal Sample. *Homo* 56: 197–205.
- Byerlee D (1974) Rural - urban immigration in Africa - Theory, policy and research implications. *International Migration Review* 8: 543–566.
- Smit R (2001) The impact of labor migration on African families in South Africa: Yesterday and today. *Journal of Comparative Family Studies* 32: 533.
- Dayal MR, Kegley ADT, Strkalj G, Bidmos MA, Kuykendall KL (2009) The history and composition of the Raymond A. Dart Collection of Human Skeletons at the University of the Witwatersrand, Johannesburg, South Africa. *American Journal of Physical Anthropology* 140: 324–335.
- Brown RW, Longoria T (2010) Multiple Risk Factors for Lead Poisoning in Hispanic Sub-Populations: A Review. *Journal of Immigrant and Minority Health* 12: 715–725.
- Grosse SD, Matte TD, Schwartz J, Jackson RJ (2002) Economic gains resulting from the reduction in children's exposure to lead in the United States. *Environmental Health Perspectives* 110: 563–569.
- Pirkle JL, Brody DJ, Gunter EW, Kramer RA, Paschal DC, et al. (1994) The Decline in Blood Lead Levels in the United States: The National Health and Nutrition Examination Surveys (NHANES). *JAMA: The Journal of the American Medical Association* 272: 284–291.
- Ezzati M, Lopez AD, Rodgers A, Vander Hoorn S, Murray CJL, et al. (2002) Selected major risk factors and global and regional burden of disease. *Lancet* 360: 1347–1360.
- Thomas KJA (2007) Child mortality and socioeconomic status: An examination of differentials by migration status in South Africa. *International Migration Review* 41: 40–74.
- Mathee A, von Schirnding YER, Levin J, Ismail A, Huntley R, et al. (2002) A survey of blood lead levels among young Johannesburg school children. *Environmental Research* 90: 181–184.
- Harlan WR (1988) The relationship of blood lead levels to blood pressure in the US population. *Environmental health perspectives* 78: 9.
- Lauphear BP, Weitzman M, Eberly S (1996) Racial differences in Urban children's environmental exposures to lead. *American journal of public health* 86: 1460–1463.
- Hu H, Payton M, Kornic S, Aro A, Sparrow D, et al. (1996) Determinants of Bone and Blood Lead Levels among Community-exposed Middle-aged to Elderly Men: The Normative Aging Study. *American Journal of Epidemiology* 144: 749–759.
- Martin D, Glass TA, Bandeen-Roche K, Todd AC, Shi W, et al. (2006) Association of Blood Lead and Tibia Lead with Blood Pressure and Hypertension in a Community Sample of Older Adults. *American Journal of Epidemiology* 163: 467–478.
- Muntner P, Madkbrfab V (2005) Continued decline in blood lead levels among adults in the United States: The National Health and Nutrition Examination Surveys. *Archives of Internal Medicine* 165: 2155–2161.
- Monna F, Poujol M, Losno R, Dominik J, Annegarn H, et al. (2006) Origin of atmospheric lead in Johannesburg, South Africa. *Atmospheric Environment* 40: 6554–6566.
- Olowoyo JO, van Heerden E, Fischer JL, Baker C (2010) Trace metals in soil and leaves of Jacaranda mimosifolia in Tshwane area, South Africa. *Atmospheric Environment* 44: 1826–1830.
- Olowoyo JO, van Heerden E, Fischer JL (2011) Trace element concentrations from lichen transplants in Pretoria, South Africa. *Environmental Science and Pollution Research* 18: 663–668.
- Lemon A (1991) *Homes apart : South Africa's segregated cities*. London: Chapman.
- Von Schirnding YER, Fuggle RF (1996) A study of the distribution of urban environmental lead levels in Cape Town, South Africa. *Science of The Total Environment* 188: 1–8.
- Von Schirnding YER, Fuggle RF, Bradshaw D (1991) Factors associated with elevated blood lead levels in inner-city Cape Town children. *SAMJ-South African Medical Journal* 79: 454–456.
- Mathee A, Rollin H, Levin J, Naik I (2006) Lead in Paint: Three Decades Later and Still a Hazard for African Children? *Environ Health Perspect* 115: 321–322.

41. Mathee A, Singh E, Mogotsi M, Timothy G, Maduka B, et al. (2009) Lead-based paint on playground equipment in public children's parks in Johannesburg, Tshwane and Ekurhuleni. *SAMJ- South African Medical Journal* 99: 819–821.
42. Montgomery M, Mathee A (2005) A preliminary study of residential paint lead concentrations in Johannesburg. *Environmental Research* 98: 279–283.
43. Morrow L, Needleman HL, McFarland C, Aethenv K, Tobin M (2007) Past occupational exposure to lead: Association between current blood lead and bone lead. *Archives of Environmental & Occupational Health* 62: 183–186.
44. Barry PSI, Mossman DB (1970) Lead concentrations in human tissues. *British Journal of Industrial Medicine* 27: 339–351.
45. Barry PS (1975) A comparison of concentrations of lead in human tissues. *British Journal of Industrial Medicine* 32: 119–139.
46. Wittmers LE, Wallgren J, Alich A, Aufderheide AC, Rapp G (1988) Lead in Bone IV. Distribution of lead in the human skeleton. *Archives of Environmental Health* 43: 381–391.
47. Rabinowitz MB, Wetherill GW, Kopple JD (1976) Kinetic analysis of lead metabolism in healthy humans. *The Journal of Clinical Investigation* 58: 260–270.
48. Cheng YW, Schwartz J, Sparrow D, Aro A, Weiss ST, et al. (2001) Bone lead and blood lead levels in relation to baseline blood pressure and the prospective development of hypertension - The Normative Aging Study. *American Journal of Epidemiology* 153: 164–171.
49. Glenn BS, Bandeen-Roche K, Lee BK, Weaver VM, Todd AC, et al. (2006) Changes in systolic blood pressure associated with lead in blood and bone. *Epidemiology* 17: 538–544.
50. Navas-Acien A, Schwartz BS, Rothenberg SJ, Hu H, Silbergeld EK, et al. (2008) Bone lead levels and blood pressure endpoints. *Epidemiology* 19: 496–504.
51. Norman R, Mathee A, Barnes B, van der Merwe L, Bradshaw D, et al. (2007) Estimating the burden of disease attributable to lead exposure in South Africa in 2000. *SAMJ -South African Medical Journal* 97: 773–780.
52. Wedeen RP (1988) Bone lead, hypertension, and lead nephropathy. *Environmental Health Perspectives* 78: 57–60.
53. Baranowska I, Czernicki K, Aleksandrowicz R (1995) The analysis of lead, cadmium, zinc, copper and nickel content in human bones from the Upper Silesian Industrial District. *Science of the Total Environment* 159: 155–162.
54. Needleman H (2004) Lead poisoning. *Annual Review of Medicine* 55: 209–222.
55. Baghurst PA, McMichael AJ, Wigg NR, Vimpani GV, Robertson EF, et al. (1992) Environmental Exposure to Lead and Children's Intelligence at the Age of Seven Years. *New England Journal of Medicine* 327: 1279–1284.
56. Carpenter DO, Nevin R (2010) Environmental causes of violence. *Physiology & Behavior* 99: 260–268.
57. Koller Ka (2004) Recent developments in low-level lead exposure and intellectual impairment in children. *Environmental Health Perspectives* 112: 987–994.
58. Needleman HL, Riess JA, Tobin MJ, Biesecker GE, Greenhouse JB (1996) Bone lead levels and delinquent behavior. *JAMA-Journal of the American Medical Association* 275: 363–369.
59. Needleman HL, McFarland C, Ness RB, Fienberg SE, Tobin MJ (2002) Bone lead levels in adjudicated delinquents - A case control study. *Neurotoxicology and Teratology* 24: 711–717.
60. Nevin R (2000) How lead exposure relates to temporal changes in IQ, violent crime, and unwed pregnancy. *Environmental Research* 83: 1–22.
61. Nevin R (2009) Trends in preschool lead exposure, mental retardation, and scholastic achievement: Association or causation? *Environmental Research* 109: 301–310.
62. Bellinger DC, Stiles KM, Needleman HL (1992) Low level lead exposure, intelligence and academic achievement - a long term follow up study. *Pediatrics* 90: 855–861.
63. Bandeen-Roche K, Glass TA, Bolla KI, Todd AC, Schwartz BS (2009) Cumulative Lead Dose and Cognitive Function in Older Adults. *Epidemiology* 20: 831–839.
64. van Wijngaarden E, Campbell JR, Cory-Slechta DA (2009) Bone lead levels are associated with measures of memory impairment in older adults. *Neurotoxicology* 30: 572–580.
65. Wright JP, Boisvert D, Vaske J (2009) Blood Lead Levels in Early Childhood Predict Adulthood Psychopathy. *Youth Violence and Juvenile Justice* 7: 208–222.
66. Mielke HW, Zahran S (2012) The urban rise and fall of air lead (Pb) and the latent surge and retreat of societal violence. *Environment International* 43: 48–55.
67. Weiskopf MG, Wright RO, Schwartz J, Spiro A, Sparrow D, et al. (2004) Cumulative lead exposure and prospective change in cognition among elderly men. *American Journal of Epidemiology* 160: 1184–1193.
68. Kon ZR, Lackan N (2008) Ethnic Disparities in Access to Care in Post-Apartheid South Africa. *American Journal of Public Health* 98: 2272–2277.
69. Peltzer K (2001) Psychosocial correlates of healthy lifestyles in black and white South Africans. *Social Behavior and Personality* 29: 249–255.
70. Peltzer K (2002) Health behaviour among Black and White South Africans. *Journal of the Royal Society for the Promotion of Health* 122: 187–193.
71. Herman DS, Geraldine M, Venkatesh T (2009) Influence of minerals on lead-induced alterations in liver function in rats exposed to long-term lead exposure. *Journal of Hazardous Materials* 166: 1410–1414.
72. Mahaffey KR (1983) Biototoxicity of lead - Influence of various factors. *Federation Proceedings* 42: 1730–1734.
73. Goyer RA (1995) Nutrition and metal toxicity. *American Journal of Clinical Nutrition* 61: S646–S650.
74. Motosue AM, Petronella S, Sullivan J, Castillo S, Garcia T, et al. (2009) Lead Exposure Risk is Associated with Asthma in a Low-income Urban Hispanic Population: Results of the Communities Organized against Asthma and Lead (COAL) Project. *Journal of Allergy and Clinical Immunology* 123: S20.
75. van Gemert F, van der Molen T, Jones R, Chavannes N (2011) The impact of asthma and COPD in sub-Saharan Africa. *Primary Care Respiratory Journal* 20: 240–248.
76. Heckman J, Samie A, Bessong P, Ntsieni M, Hamandi H, et al. (2010) Anaemia among clinically well under-fives attending a community health centre in Venda, Limpopo Province. *SAMJ -South African Medical Journal* 100: 445–448.
77. Faber M, Wenhold F (2007) Nutrition in contemporary South Africa. *Water Sa* 33: 393–400.
78. Nojilana B, Norman R, Dhansay MA, Labadarios D, van Stuijvenberg ME, et al. (2007) Estimating the burden of disease attributable to iron deficiency anaemia in South Africa in 2000. *SAMJ-South African Medical Journal* 97: 741–746.