Rapid Response: COVID-19 and Black Africans in the UK: Disparities linked to underlying inequalities in health by F. Alloh, P. Regmi & E. van Teijlingen

https://www.bmj.com/content/369/bmj.m1548/rr-18

Rapid response to: Is ethnicity linked to incidence or outcomes of covid-19? *BMJ* 2020; 369 doi: <u>https://doi.org/10.1136/bmj.m1548</u> (Published 20 April 2020)

Article	Related content	Article metrics	Rapid responses	Response
Rapid Re:	sponse:			
		n the UK: Disparitie	es linked to underlyi	ng inequalities
health				-0 q
loann				
Dear Editor,				
The COVID-1	19 nandemic has drawn	attention to the impa	ct of health inequality o	on the virus outco
			ancet around Black and	
and the disp	roportionate severity of	f COVID-19 among thi	s group in the UK [1,2]. E	BAMEs account fo
than 34% of	critically ill patients from	m COVID-19, keeping	in mind that this group (only make up 14
total nonula	tion in the LIK [3]. In con	oparison to the White	population have only 73	8.6% of deaths w

Dear Editor,

The COVID-19 pandemic has drawn attention to the impact of health inequality on the virus outcome. In response to the important conversations in the BMJ and Lancet around Black and Ethnic Minorities (BAMEs) and the disproportionate severity of COVID-19 among this group in the UK [1,2]. BAMEs account for more than 34% of critically ill patients from COVID-19, keeping in mind that this group only make up 14% of the total population in the UK [3]. In comparison to the White population have only 73.6% of deaths while representing more than 85% of the total population, this brings -14% deaths in this group when compared to the total population.

However, we would like to focus this conversation on disproportionate infection and mortality of the black-African population due to COVID-19 in the UK. This is important because Black-African has been reported to be more disadvantaged than other BAMEs [4]. It has been reported that black-Africans make up 5.8% of the cases of COVID deaths while representing only 3.5% of the total population. They have highest hospital deaths with three times that of the White population [5]. In comparison, Asian group made up 7.6% of deaths due to COVID-19 while representing 7.7% of the total population. Similarly in London, black-African accounted for more than 16% of the deaths while only representing 13% of the total population. This is higher in comparison to Whites 55.9% deaths with 59.8% representation of the population and Asian 16.4% deaths while representing 18.5% of the population [6]. This prompted us to review publications in relation to COVID-19 outcome among BAMEs in the UK and found that no study has focused on Black-Africans and the severity of COVID-19 outcome among individuals from this group.

Black- African population in the UK have continuously faced health inequality which has disadvantaged this group in relation to COVID-19 outcome. African population have a higher burden of non-communicable diseases such as diabetes, hypertension, mental health in the UK [7]. Furthermore, cultural beliefs in regard to conditions such as HIV and AIDs and

tuberculosis have contributed to the higher burden of these diseases among African populations in the UK [8]. These burdens of chronic disease among black-African populations have implications for the higher burden of COVID-19 reported within these groups.

Other factors such as genetic composition, vitamin D deficiency, vaccine from country of birth, immune system performance might be important in COVID-19 outcome among Africans in the UK [9]. In addition, limited access to health-care services, working in high-risk jobs such as care worker, health worker has made African immigrants more susceptible to COVID-19 infection and mortality. This is in addition to the overly-representation of this group in low socio-economic status which has important implications on health and wellbeing.

Finally, COVID-19 has highlighted the disparity of African population health in the UK as compared to the general population. We feel it is time that policy address this long-overdue inequality in health among BAMEs and African population in particular. It should not be expected as normal that these groups are worse hit by COVID-19 or any other diseases. Targeted research is needed to understanding the implications of inequality on health of Black-Africans that are disproportionately affected by disease conditions.

References

1. Khunti Kamlesh, Singh Awadhesh Kumar, Pareek Manish, Hanif Wasim. Is ethnicity linked to incidence or outcomes of covid-19? BMJ 2020; 369 :m1548

2. Pareek, M., Bangash, M.N., Pareek, N., Pan, D., Sze, S., Minhas, J.S., Hanif, W. and Khunti, K. Ethnicity and COVID-19: an urgent public health research priority. The Lancet. 2020.

3. Intensive Care National Audit and Research Centre, London 2020 Available from https://www.icnarc.org/Our-Audit/Audits/Cmp/Reports Accessed April, 2020

4. Alloh, F., Hemingway, A. and Turner-Wilson, A.L. Systematic review of diabetes management among black African immigrants, white and South Asian populations. Journal of Global Health Reports. 2019;3.

5. Platt and Warwick 2020 Are some ethnic groups more vulnerable to COVID-19 than others? Available from: https://www.ifs.org.uk/inequality/wp-content/uploads/2020/04/Are-some-et... Accessed May, 2020

6. NHS statistics. COVID-19 Daily Deaths https://www.england.nhs.uk/statistics/statisticalwork-areas/covid-19-da... Accessed May, 2020

7. Regmi PR, Alloh F, Pant PR, Simkhada P, Van Teijlingen E. Mental health in BME groups with diabetes: an overlooked issue?. The Lancet. 2017;4;389(10072):904-5.

8. Gerrish, K., Naisby, A. and Ismail, M. The meaning and consequences of tuberculosis among Somali people in the United Kingdom. Journal of Advanced Nursing. 2012;68(12), pp.2654-2663.

9. Agyemang, C., de-Graft Aikins, A. and Bhopal, R. Ethnicity and cardiovascular health research: pushing the boundaries by including comparison populations in the countries of origin. Ethnicity & Health. 2012;17(6), pp.579-596.