

Table 2. Pearson correlations between baseline (B) and follow-up (FU) variables

	1. B-carer burden ^Ψ		2. FU-carer burden ^Ψ		3. B-carer quality of life [†]		4. FU-carer quality of life [†]		5. B-person with dementia quality of life [†]		6. FU-person with dementia quality of life [†]		7. B-person with dementia cognitive function [†]		8. FU-person with dementia cognitive function [†]		9. Trial arm		
	R	p	r	p	r	p	r	p	r	p	r	p	R	p	r	p	r	p	
1 ¹																			
2 ²	.79	<.001																	
3 ¹	-.66	<.001	-.49	<.001															
4 ³	-.62	<.001	-.66	<.001	.66	<.001													
5 ¹	-.05	.67	-.11	.44	.03	.83	-.09	.53											
6 ²	.06	.66	.07	.65	-.26	.06	-.01	.94	.40	<.001									
7 ¹	-.07	.59	-.06	.65	-.04	.77	.16	.26	-.27	.03	-.14	.30							
8 ³	-.28	.04	-.24	.09	.13	.37	.29	.04	-.14	.32	-.06	.68	.83	<.001					
9 ⁴	-.00	.97	.03	.85	.14	.27	-.05	.73	.21	.10	-.08	.58	-.26	.04	-.14	.32			

Notes: B = baseline, FU = follow-up. 1: N = 64, 2: N = 52, 3: N = 53, 4: N = 65. Values in bold indicate significance at $p < 0.10$.

^ΨLower score = better outcome; [†]Higher score = better outcome;