
Effect of Exposure to Asynchronous Virtual Clinical Environments on Actual/ Perceived Competence in Drug Dosage Calculation: A Pilot Study

Sandra Goldsworthy¹, Keith Weeks², Naim Abdulmohdi³, Sue Baron⁴,
Karey McCullough⁵, Nita Muir⁶, Kim Sears⁷, Alex Weeks⁸,
Grace Perez⁹, Laurence Moseley¹⁰, Matt Brown¹¹, David Pontin¹²

¹PhD, MSc, RN, CNCC(C), CMSN(C), CCSNE, Professor and Chair, School of Nursing and Midwifery, Mount Royal University, Calgary, Alberta, ²PhD, RN Authentic World, UK, ³PhD, RGN, Anglia Ruskin University, Chelmsford, UK, ⁴PhD, RN, RNT Bournemouth University, Poole, UK, ⁵PhD, RN Nipissing University, North Bay, ON, Canada, ⁶PhD, RN Queen's University, Kingston, ON, Canada, University of Chichester, Chichester, UK, ⁷PhD, RN Queen's University, Kingston, ON, Canada, University of Chichester, Chichester, UK, ⁸BSc Mathematics Authentic World, UK, ⁹MSc University of Calgary, Calgary, AB, Canada, ¹⁰PhD Authentic World, UK, ¹¹BA (Hons) Authentic World, UK, ¹²PhD, RN Numeric Generics, UK.

How to cite this article: Sandra Goldsworthy, Keith Weeks, Naim Abdulmohdi et al. Effect of Exposure to Asynchronous Virtual Clinical Environments on Actual/Perceived Competence in Drug Dosage Calculation: A Pilot Study. *International Journal of Nursing Education* / Vol. 18 No. 1, January-March 2026.

Abstract

Introduction: Nursing students are expected to be 'practice ready' on qualifying. This includes safe medication administration. This pilot study investigates the relationship between exposure duration to asynchronous virtual drug dosage calculation scenarios and nursing student actual and perceived competence. Methodology design planned for larger scale main study was tested and piloted.

Methods: A randomised quasi-experimental research design (pre- and post-test) was used. Purposive sampling was used to recruit six groups of second/third-year pre-registration undergraduate nursing students from six sites (UK and Canada). Students were randomly assigned to four groups of different exposure to the safeMedicate® COVID-19 education module.

Results: Student actual competence increased across all four groups, and their perceived competence mirrored this. There was no clear dose-response relationship demonstrated.

Conclusion: Valuable insights into the effects of asynchronous virtual learning on drug dosage calculation competence among nursing students were generated. Improvement in actual and perceived competence was found, but no clear dose-response relationship. Further research on a larger scale is needed to explore the impact of instructional design, feedback, and interaction on learning outcomes.

Keywords: medication errors; medication calculation; simulation; competence; nursing student; patient safety.

Corresponding Author: Sandra Goldsworthy, Professor and Chair, School of Nursing and Midwifery, Mount Royal University, Calgary, Alberta.

E-mail: sgoldsworthy@mtroyal.ca

Submission date: Aug 20, 2025

Revision date: Sept 30, 2025

Published date:

This is an Open Access journal, and articles are distributed under a Creative Commons license- CC BY-NC 4.0 DEED. This license permits the use, distribution, and reproduction of the work in any medium, provided that proper citation is given to the original work and its source. It allows for attribution, non-commercial use, and the creation of derivative work.

Introduction

Administering medications is one of the most high-risk tasks in health care. Few studies have examined the barriers nurses experience, and the strategies used to prevent medication related harm. The World Health Organisation (WHO) highlight medication errors as a serious global issue. It is the third leading cause of death in the USA⁽¹⁾; 1:20 patients globally experience preventable medication-related harm⁽²⁾. In 2017 WHO launched 'Medication Without Harm' to address medications as a top priority⁽²⁾ and stressed the topic's importance in healthcare curriculum⁽³⁾⁽⁴⁾.

Medication competence is complex and requires strong theoretical pathophysiology, pharmacology knowledge, critical analysis, numeracy and calculation skills⁽⁵⁾⁽⁶⁾. Nursing students report increased anxiety about medication calculation and administration, which can result in medication errors, near misses or actual patient harm⁽⁷⁾⁽⁸⁾. These frequently occur at administration due to inaccurate dosage calculation⁽⁹⁾⁽¹⁰⁾. Nursing student under-reporting of medication errors may be due to shame, guilt, fear of repercussion, decreased confidence, and anxiety⁽¹¹⁾⁽¹²⁾. It is prudent to understand the complexity of how nurses are educated in preparing to administer medication.

Since the COVID-19 pandemic, medication safety concerns have continued due to the pandemic's lasting impact on the ongoing global nursing shortage. COVID-19 triggered a call for nursing students to be practice ready and competent in calculating and administering medication on qualifying⁽⁴⁾⁽¹³⁾⁽¹⁴⁾⁽¹⁵⁾⁽¹⁶⁾⁽¹⁷⁾⁽¹⁸⁾. The pandemic highlighted that all nurses must be ready to safely practice where rapid mobilization/healthcare resource adaptation was needed to minimize COVID-19's impact on populations and health systems⁽¹⁹⁾⁽²⁰⁾⁽²¹⁾⁽²²⁾⁽²³⁾.

Medication administration is a practice-based skill. Nurses need the opportunity and right environment to acquire this skill before

administering medications to real patients. Consistency and practice opportunity availability throughout each nursing curricula year is important for improving medication safety competence and confidence⁽⁵⁾⁽²⁴⁾⁽²⁵⁾. These may not be routinely available in practice settings⁽²⁸⁾⁽⁸⁾. Medication errors account for 38% of adverse practice events reported by undergraduate nursing students⁽³⁾⁽¹⁰⁾; difficulty in dosage calculations and poor mathematics skills are key contributory factors⁽¹²⁾⁽²⁶⁾⁽²⁷⁾. Traditional clinical learning environments are not the best, safest or only locations for students to develop competency and confidence in medication calculation and administration skills⁽⁷⁾⁽²⁹⁾⁽³⁰⁾.

safeMedicate® developed an international COVID-19 version of its Authentic Virtual Drug Dosage Calculation Clinical Learning Environment (VLE) in 2020 to support RN competence adaptation. It included typical COVID-19 therapies for hospital patients with pneumonia and pyrexia and for patients requiring critical care therapies⁽³⁹⁾⁽⁴⁰⁾

Given the knowledge and experience of the nursing student sample in this study, actual competence assessments excluded critical care therapies, focussing on minimal requirements for safe pandemic practice⁽⁴¹⁾. The 12 test items included varied drug formulations/administration routes and complexity levels (Table 1). Perceived competence is a pertinent corollary for investigation given the commonly reported decreased confidence and anxiety associated with drug dosage calculation, i.e. how confident students perceive they correctly solve each test-item.

Simulation-based education has a positive effect on developing knowledge and confidence for medication administration⁽³¹⁾⁽³²⁾⁽³³⁾⁽⁶⁾. There is opportunity to understand how simulation-based education and technology may improve medication administration due to increased use of technology-driven strategies to support quality and safety teaching in nursing education⁽³⁴⁾⁽³⁵⁾⁽³⁶⁾⁽³⁷⁾.

Table 1: 12-item-safeMedicate® COVID-19-specific drug therapy test of actual competence

Number of Pre- & Post-Test items (questions)	Drug Formulation	Drug Administration Route (PO=Oral; IV= Intra-venous)	Item Rubric (Complexity) Level	Item- Score weighted by total problem-solving phases (% per phase)	Total Score
2	Tablets & Capsules	PO	Unit Dose	3 x (%)	6
2	Oral Suspension	PO	Multiple Unit-Dose	3 x (%)	6
1	Intravenous Injection	IV	Slow IV-Injection (Timed)	4 x (25%)	4
2	Intravenous Infusion	IV	Intermittent Infusion (Standard Dose)	3 x (%)	6
5	Intravenous Infusion	IV	Intermittent Infusion (Bodyweight-Based Dose)	6 x (%)	30
					52

Method

Aim

To investigate the effect of exposure to asynchronous online virtual clinical environments of differing duration on actual and perceived competence in COVID-19-related drug dosage calculation.

Research Questions

1. What is the effect of exposure duration to an online virtual clinical environment on nursing students' actual competence in COVID-19 support drug therapy?
2. What is the effect of exposure duration to an online virtual clinical environment on nursing students' perceived competence in COVID-19 support drug therapy?

Design

A pilot pre/post-test randomised quasi-experimental research design was used.

Sample/setting

Second- and third-year pre-registration Bachelor of Science nursing students were recruited via purposive sampling from four UK sites: University

of Chichester, Anglia Ruskin University and Bournemouth University (England), and Robert Gordon University (Aberdeen, Scotland), and two Canadian sites: Nipissing University and Queen's University. We recruited 128 students (n=38 completed the study following high attrition rates); the International Standard Randomised Controlled Trial Number registry identifies a sample size of 30-40 for feasibility studies⁽³⁸⁾.

Data collection

Ethical approval was obtained from each site in late 2022. The target population received an invitation-to-participate email from the study administrator. Controls were put in place for numeracy skill variation and orientation to the virtual learning environment (VLE) (safeMedicate® VLE/competence-perceived competence metric). Data was collected January-October 2023 at each site using:

1. 12-item-safeMedicate® COVID-19-specific drug therapy test (0%-100%) to measure actual competence.
2. safeMedicate® VLE/competence-perceived competence metric (0%-100%) to measure perceived competence.

Concept and Operational Definitions

Actual Competence is an objective measure of individual knowledge, skills and abilities in a specific context. The safeMedicate competence model was used as the operational definition.

Supplementary Material: Video 1⁽⁴²⁾: Models drug dosage calculation problem-solving and error diagnosis for:

- Conceptual competence: Dose/Rate of infusion equation set-up ability.
- Calculation competence: Dose/Rate of infusion computation ability.
- Measurement/Technical Measurement competence: Dose/Rate of infusion measurement ability.

Perceived competence is a psychological construct based on individual self-evaluation of capability in a specific context. We defined it as how confident students were that a test-item was correctly solved. A slide-bar scale metric (range 0%–100%) was used to record perceived competence. We looked for any discrepancy between actual and perceived competence, i.e. were students under/over-confident, or accurate in their perception of their actual competence.

The intervention used the safeMedicate® Authentic Virtual Drug Dosage Calculation Clinical Learning Environment (VLE) to engage students. The VLE is based on an adapted cognitive apprenticeship model⁽⁴²⁾. This is a method for externalizing expert problem-solving processes that are often obscured from students. It uses constructivist-based modelling, coaching, scaffolding and abstracted replay feedback that facilitate learners' understanding, development and application of competence, and reflection on their own and expert problem-solving practice⁽⁴²⁾.

Students undertook authentic diagnostic assessments compared with expert problem-solving models (**Supplementary Material: Videos 2, 3 & 4**

⁽⁴²⁾. They had detailed diagnostic reports and feedback on their actual and perceived competence assessment activities and outcomes compared to correct answers throughout the study. Abstracted replay feedback loops and error diagnosis help students identify their strengths and areas needing remediation. They had variable 0–3-week exposure to the safeMedicate COVID-19 Education Support Module (Intervention).

Supplementary Material: Video 2: Illustrates a comparison of student conceptual, calculation and technical-measurement competence assessment processes and outcomes, compared with an expert problem-solving model.

Supplementary Material: Video 3: Illustrates an example of abstracted-replay feedback, diagnosis of an actual competence problem-solving error, and self-rated over-estimation of competence. Over-estimation of competence was a limited finding in the study (Figure 2). It has been illustrated here to highlight early diagnosis of patient-safety critical errors, and how this can act as a powerful learning event in the education process.

Supplementary Material: Video 4: Shows diagnosis of actual-competence development, and self-rated under-estimation of competence. Under-estimation of competence was a common finding (Figure 2).

Healthcare Numeracy Assessment

The safeMedicate® 25-item Healthcare Numeracy Assessment (HNA) assessed student baseline competence in 14/20 essential healthcare numeracy skills⁽⁴¹⁾. Test items focused on fundamental mathematics stripped of specific nursing contexts. Test item-score reliability was used as a criterion for item selection. HNA test results were used to assign students to four groups via stratified competence profiling. This controlled for student numeracy skill variability.

Procedure

Phase	Activity	Group	Weeks							
			1	2	3	4	5	6	7	8
1	Orientation	All								
	HNA Assessment	All								

Continue.....

2	Group assignment	All									
3	safeMedicate orientation module	All									
4	12 item COVID-19 Pre-Test	All									
5	Exposure to safeMedicate COVID-19 Education Support module intervention	Control									
		1									
		2									
		3									
6	12 item COVID-19 Post-Test	Control									
		1									
		2									
		3									

Figure 1: Study design

Figure 1 documents the study design. It sets out project activities and timelines.

Ethics considerations

Ethics approval was obtained at all sites.

Data analysis

Data analysis methodology consisted of:

- Descriptive and inferential statistical analyses conducted using SPSS V29. Data were quantified using means and standard deviations (SDs). Non-parametric statistical tests were carried out as participant attrition resulted in a dataset (n=38) that did not meet assumptions for parametric analysis. Statistical significance was set at an alpha level of $p < 0.05$.
- Piloting analysis of correlation between actual and perceived competence [Spearman's Correlation]; and analysis of

treatment effect for pre-test/post-test actual and perceived competence [ANCOVA] as part of methodology design testing. This will be used for larger sample size data analysis in the main study.

Results

Study Assessments

Students' actual competence and perceived competence results were compared before/after variable exposure to the intervention. The four groups were homogeneous at baseline, i.e. no differences in the mean pre-test assessments. Actual and perceived competence mean scores increased in the post-test assessments, including the control group which did not have any intervention exposure (Table 2; Figure 2). However, abstracted replay pre-test competence/error diagnostic feedback (see Video 3) may have acted as a learning event; this will be considered during the design of the main study.

Table 2: Analysis of difference between mean HNA and pre-test to post-test actual and perceived competence percentage scores (*Based on Kruskal-Wallis test)⁽⁴³⁾

Intervention Group	Statistic	HNA	Pre-Test		Post-Test	
			Actual Competence	Perceived Competence	Actual Competence	Perceived Competence
Group 1	N	15	15	15	15	15
	Mean	83.47	77.18	64.53	95	82.78
	SD	14.09	22.6	27.05	6.77	25.45
	Min	52	38.46	15.83	80.77	0
	Max	100	100	99.17	100	100
	Median	88	86.54	77.33	98.08	91.25

Continue.....

Group 2	N	11	11	11	11	11
	Mean	88.73	79.72	73.48	87.94	83.9
	SD	11.98	24.16	22.89	21.71	20.98
	Min	60	32.69	39.42	34.62	38.25
	Max	100	100	100	100	100
	Median	92	86.54	77	100	92.92
Group 3	N	8	8	8	8	8
	Mean	81.5	78.85	77.93	95.67	97.68
	SD	17.75	24.59	20.92	5.51	3.78
	Min	48	40.38	39.83	86.54	89
	Max	100	100	100	100	100
	Median	86	87.5	79.79	98.08	99.17
Control Group	N	4	4	4	4	4
	Mean	84	67.31	66.98	84.62	82.63
	SD	13.86	19.8	24.75	20.05	9.31
	Min	64	44.23	40.83	55.77	69.25
	Max	96	86.54	99.58	100	89.5
	Median	88	69.23	63.75	91.35	85.88
Group Comparison*	p-value	0.712	0.682	0.627	0.738	0.085




Figure 2: Mean pre-test/post-test actual/perceived competence scores

Students performed better on the 12 item COVID-19 post-test of actual competence after the intervention. Figure 3 shows the change between

individual student pre-test score and post-test score. Table 3 shows the change in average pre-test score (mean 77%; median 87%), to post-test score (mean

92%; median 98%). A Wilcoxon Test indicated this was statistically significant ($p < 0.0004$). Although a small pilot study sample, students achieved these

outcomes over 0-3 weeks rather than the typical 3-4 years associated with pre-registration/licensure nurse education programmes.




Figure 3: Change in Pre-Test/Post-Test Actual Competence scores

Table 3: Wilcoxon Signed Rank Test Analysis

Descriptive Statistics

	N	Mean	SD	Minimum	Maximum	Percentiles		
						25 th	50 th (Median)	75 th
Pre-Test Actual Competence	38	77	22.59	32.69	100	59.61	86.54	96.15
Post-Test Actual Competence	38	92	14.16	34.62	100	88.94	98.08	100

Ranks

	Mean Rank	Sum of Ranks
Pre-Test Actual Competence	20.5	532.5
Post-Test Actual Competence	10.8	97.5

Rank Differences

Negative Ranks	26
Positive Ranks	9
Ties	3
Total	38

Test Statistics

p-values	
1-tail	0.0002
2-tail	0.0004

Discussion

Findings suggest general improvement across all groups in actual and perceived competence including the control. Significant outcome differences between groups with varied exposure duration were not evident. Increasing exposure time did not translate into substantial competence improvement.

Improvement in actual and perceived competence suggests initial VLE and baseline assessment exposure might have enhanced competency. Assessment outcome feedback can foster incremental learning, boosting competence and confidence⁽⁴⁴⁾⁽⁴²⁾. Increased post-test perceived competence aligns with research on VLE and self-assessment influence on learners' perceived proficiency^{(45) (46)}. Lack of significant

differences between groups with varying exposure durations contradicts earlier studies advocating for prolonged engagement with VLEs to achieve competence⁽⁴⁷⁾.

Our findings indicate a shift in students' self-awareness and competence perception. This counters existing evidence that suggests students overestimate their competence⁽⁴⁸⁾. Our work suggests students are cautious in overestimating their competence, which is an understandable safety conscious response.

Our study included VLE feedback on student performance for review, and indicates that simulation-based intervention effectiveness is based on instructional design and not dependent on exposure time. Findings may be related to small sample size and need repeating with a larger sample. The differences in perceived competence before/after intervention highlight the psychological and confidence-building value of VLEs, consistent with previous research on self-efficacy enhancement effects of digital learning^{(36) (49)}.

The results suggest revisiting VLE design. Prolonged exposure alone did not result in significantly better outcomes. Our findings indicate the potential of short-term exposure to VLEs to positively influence actual and perceived competence. Brief, focused interventions may rapidly bolster nursing students' confidence and awareness. This may be valuable in pandemic preparedness or acute care settings where speedy upskilling to mobilize healthcare staff adaptation is critical. However, we would strongly advocate for repeated periodic assessment to evaluate competence retention and address any skill decay.

Implications

This pilot study provides valuable insights into the use of asynchronous virtual learning environments (VLEs) for developing drug dosage calculation competence i.e. brief exposure, actual competence development, and feedback design.

Repeated simulation exposure is associated with improvements in nursing students' self-efficacy, perceived competence, and learning satisfaction⁽⁵⁰⁾. However, our findings suggest that even brief exposure to accessible, simulation-based learning

with structured feedback and opportunities for repetition, can enhance actual as well as perceived competence. This has practical implications for nursing education and healthcare provider strategies where rapid upskilling is required, such as public health emergency response or workforce shortages.

Our study adds weight to Koukourikos et al's findings⁽⁵¹⁾ of increased perceived clinical competence through VLE use during COVID-19, by showing positive changes in actual competence. Although we faced challenges as a multisite, international collaboration over differing time zones, this allows us to share diverse perspectives on undergraduate nursing learning strategies in a global context. Our findings echo Bae et al's systematic review⁽⁵²⁾. This supports using VLEs to enhance clinical reasoning, problem-solving, and communication competencies when traditional clinical placements are limited and may not provide consistent or safe opportunities for skill acquisition. Our work helps educators incorporate targeted, technology-driven interventions to complement traditional clinical placements.

The results show the importance of feedback design, where instructional strategies may be more influential than exposure duration alone. Future research should expand on these findings by using larger samples, and longitudinal designs to assess skill retention and explores stematic integration of VLEs into curricula to build confidence and competence in drug dosage calculation.

Limitations

This study has several limitations. First, the small sample size (n=38) as a result from the high attrition rate, limit the statistical power of the findings and restrict generalizability beyond the study population. As a pilot study, results should be interpreted with caution until replicated with larger, more diverse cohorts. Second, participants were recruited from selected universities in the UK and Canada; therefore, findings may not reflect the experiences of students in other regions or curricula. Third, the reliance on self-reported measures of perceived competence introduces the possibility of response bias, as students may have under- or over-estimated their abilities despite efforts to mitigate this through diagnostic

feedback. Fourth, the short duration of exposure (0–3 weeks) provides only a snapshot of competence development; longer-term retention of knowledge and skill decay were not assessed. Finally, while the intervention targeted drug dosage calculations related to COVID-19 therapies, exclusion of critical care scenarios limits insight into competence in more complex, high-acuity contexts.

Conclusion

This pilot offers valuable insights into the effects of asynchronous VLEs on drug dosage calculation competence among nursing students. Improvements were observed in actual and perceived competence, but lack of a clear dose-response relationship points to future research to explore the impact of instructional design, feedback, and interaction on learning outcomes. Simulation-based education remains a promising avenue, but its implementation should be refined to maximize actual and perceived competency in high-risk tasks like medication administration.

Acknowledgements: The authors acknowledge the Canadian Association of Schools for Nursing (CASN) for the Pat Griffin grant. Special thanks to Fiona Budden, Ruan Muldoon and Jan Hutt from Bournemouth University.

Acknowledgements

The research team would like to acknowledge funding from the Canadian Association of

Schools of Nursing (CASN) through the Pat Griffin Grant and Ms. Lisa Fasken,

Research Coordinator for all of her contributions to this project.

We would also like to recognize one of our retired contributors Dr. Kate Goodhand formerly of Robert Gordon, Aberdeen, UK.

Funding Source

This research was made possible through the Canadian Association of Schools of Nursing (CASN) Pat Griffin Grant.

Ethical Clearance : This research received Ethical approval from the Research Ethics Board at Nipissing University, North Bay, Ontario, Canada.

Conflicts of interest statement

Dr Keith W. Weeks, Alex Weeks and Matt Brown are directors of Authentic World Ltd, a nursing spin-out company of the University of South Wales and Cardiff University, UK.

Authentic World Ltd was the first joint university nursing spin-out company in the UK (founded 2004), and its Thought Leadership team is responsible for the international translational research, knowledge transfer, design, development, health professional education technology evaluation and international distribution of the safeMedicate® suite of authentic virtual medicines calculations and nursing mathematics clinical education environments. Dr David Pontin is a director of Numeric Generics Ltd, an associate company of Authentic World Ltd.

A university-engaged statistician independent of the research team carried out the data analysis reported here.

The authors declare no conflict of interest in the completion of this research.

Supplementary Material

1. Supplementary Material: Video 1: safeMedicate competence model.https://safemedicate.com/publications/comp_pilot_study/sm_model.html
2. Supplementary Material: Video 2: Example of safeMedicate authentic actual/perceived competence pre-test assessment process.https://safemedicate.com/publications/comp_pilot_study/sm_pre_assessment.html
3. Supplementary Material: Video 3: Example of safeMedicate pre-test abstracted-replay feedback on student and expert problem-solving processes. https://safemedicate.com/publications/comp_pilot_study/sm_pre_feedback.html
4. Supplementary Material: Video 4: Example of safeMedicate post-test abstracted-replay feedback on student and expert problem-solving processes. https://safemedicate.com/publications/comp_pilot_study/sm_post_feedback.html.

References

1. Makary MA, Daniel M. Medical error-the third leading cause of death in the US. *BMJ - British Medical Journal*. 2016;353:i2139.

2. World Health Organization. Global burden of preventable medication-related harm in health care: A systematic review. Geneva: World Health Organization; 2023. Available from: <https://iris.who.int/bitstream/handle/10665/376203/9789240088887-eng.pdf?sequence=1>.
3. World Health Organization. Medication without harm: WHO global patient safety challenge on medication safety Geneva: World Health Organization; 2017 [Available from: <https://iris.who.int/bitstream/handle/10665/255263/WHO-HIS-SDS-2017.6-eng.pdf?sequence=1>].
4. World Health Organization. Patient safety curriculum guide: multi-professional edition 2011 [Available from: <https://www.who.int/publications/i/item/9789241501958>].
5. Jarvill M. Nursing student medication administration performance: a longitudinal assessment. *Nurse Educ.* 2021;46(1):59-62.
6. Brauneis L, Badowski D, Maturin L, Simonovich SD. Impact of low-fidelity simulation-based experiences in a pharmacology classroom setting in prelicensure graduate nursing education. *Clinical Simulation in Nursing.* 2021;50(January):43-7.
7. Fusco LA, Alfes CM, Weaver A, Zimmermann E. Medication safety competence of undergraduate nursing students. *Clinical Simulation in Nursing.* 2021;52(March):1-7.
8. Schneidereith TA. Using simulations to identify nursing student behaviors: a longitudinal study of medication administration. *Journal of Nurse Education.* 2014;53(2):89-92.
9. Williams B, Davis S. Maths anxiety and medication dosage calculation errors: a scoping review. *Nurse Educ Pract.* 2016;20(September):139-46.
10. Asensi-Vicente J, Jiménez-Ruiz I, Vizcaya-Moreno MF. Medication errors involving nursing students: a systematic review. *Nurse Educ.* 2018;43(5):e1-e5.
11. Treiber LA, Jones JH. After the medication error: recent nursing graduates' reflections on adequacy of education. *J Nurs Educ.* 2018;57(5):275-80.
12. García-Gámez M, Morales-Asencio JM, García-Mayor S, Kaknani-Uttumchandani S, Martí-García C, Lopez-Leiva I, et al. Adverse events encountered during clinical placements by undergraduate nursing students in Spain. *Nurse Educ Today.* 2020;91(August).
13. Curtis K, Brysiewicz P, Shaban RZ, Fry M, Considine J, Gamboa FEA, et al. Nurses responding to the World Health Organization (WHO) priority for emergency care systems for universal health coverage. *Int Emerg Nurs.* 2020;50(May).
14. Low TY, Hartman M, Chee CYJ, Mohankumar B, Ang SBL, San MT, et al. Restructuring the surgical service during the COVID-19 pandemic: experience from a tertiary institution in Singapore. *Am J Surg.* 2020;220(3):553-5.
15. Torlinski T, Rakasz L, Wysota B, Czyz M, Snelson C. An interdisciplinary approach to the management of critically ill patients during COVID-19 pandemic; an experience of a university hospital in England. *Waidomosci Lekarskie.* 2020;LXXIII(7):1576-9.
16. Health Education England. Back to clinical practice COVID-19: Health Education England; 2020 [Available from: <https://www.hee.nhs.uk/our-work-capitalnurse/back-clinical-practice-covid-19>].
17. Sutton-Smith L. Planning for a COVID-19 crisis. *Kai Tiaki Nursing New Zealand.* 2020;26(4):26-7.
18. Wanless S, Winterman E, Chapman J. Skills teaching in COVID lockdown in the UK: lessons learnt. *Pielegniarstwo XXI Wieku-- Nursing in the 21 Century.* 2020;19(3):171-3.
19. NHS England. Joint statement on developing immediate critical care nursing capacity 2020 [Available from: <https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/52/2020/03/critical-care-joint-statement-25-march-2020.pdf>].
20. NHS England. Coronavirus: principles for increasing the nursing workforce in response to exceptional increased demand in adult critical care. Specialty guides for management during the coronavirus pandemic 2020. Available from: https://www.cc3n.org.uk/uploads/9/8/4/2/98425184/specialty_guide_critical_care_workforce_v1_25_march.pdf.
21. NHS Health Education England. Aftercare needs of inpatients recovering from COVID-19. Version 2 2020. Available from: https://madeinheene.hee.nhs.uk/Portals/0/C0705_Aftercare%20needs%20of%20inpatients%20recovering%20from%20COVID-19_3Aug%20-%20Copy.pdf.
22. Public Health Agency of Canada. COVID-19 pandemic guidance for health care sector Ottawa, ON.: Government of Canada; 2020 [Available from: <https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirus-infection/health-professionals/covid-19-pandemic-guidance-health-care-sector.html>].
23. Welsh Government. All Wales critical care escalation guidance for the management of large unplanned

- increases in demand - December 2016 Cardiff: Welsh Government; 2016 [Available from: <https://www.gov.wales/sites/default/files/publications/2019-11/all-wales-critical-care-escalation-guidance-for-the-management-of-large-unplanned-increases-in-demand.pdf>].
24. Pettigrew J, Stunden A, McGlynn S. Contextualising numeracy skill development and assessment in a first year undergraduate nursing subject: A mixed methods research study. *Nurse Educ Today*. 2020;92:8.
 25. Moloney M, Kingston L, Doody O. Fourth year nursing students' perceptions of their educational preparation in medication management: an interpretative phenomenological study. *Nurse Educ Today*. 2020;92.
 26. Mariani B, Ross JG, Paparella S, Allen LR. Medication safety simulation to assess student knowledge and competence. *Clinical Simulation in Nursing*. 2017;13(5):210-6.
 27. Weeks K, Coben D, Lum G, Pontin D. Developing nursing competence: future proofing nurses for the changing practice requirements of 21st century healthcare. *Nurse Educ Pract*. 2017;27(November):A3-A4.
 28. Jarvill M, Jenkins S, Akman O, Astroth KS, Pohl C, Jacobs PJ. Effect of simulation on nursing students' medication administration competence. *Clinical Simulation in Nursing*. 2018;14:3-7.
 29. Lee SE, Quinn BL. Incorporating medication administration safety in undergraduate nursing education: a literature review. *Nurse Educ Today*. 2019;72(January):77-83.
 30. Sears K, Goldsworthy S, Goodman WM. The relationship between simulation in nursing education and medication safety. *J Nurs Educ*. 2010;49(1):52-5.
 31. Zahara-Such RM. Improving Medication Calculations of Nursing Students through Simulation: An Integrative Review. *Clinical Simulation in Nursing*. 2013;9(9):e379-e83.
 32. Hewitt J, Tower M, Latimer S. An education intervention to improve nursing students' understanding of medication safety. *Nurse Educ Pract*. 2015;15(1):17-21.
 33. Aggar C, Bloomfield JG, Frotjold A, Thomas THT, Koo F. A time management intervention using simulation to improve nursing students' preparedness for medication administration in the clinical setting: a quasi-experimental study. *Collegian: Journal of the Royal College of Nursing, Australia*. 2018;25(1):105-11.
 34. Wyres M, Taylor N. Covid-19: using simulation and technology-enhanced learning to negotiate and adapt to the ongoing challenges in UK healthcare education. *BMJ Simulation & technology Enhanced Learning*. 2020;6(6):317-9.
 35. Altmiller G, Pepe LH. Influence of technology in supporting quality and safety in nursing education. *Nurs Clin North Am*. 2022;57(4):551-62.
 36. Goldsworthy S, Muir N, Baron S, Button D, Goodhand K, Hunter S, et al. The impact of virtual simulation on the recognition and response to the rapidly deteriorating patient among undergraduate nursing students. *Nurse Educ Today*. 2022;110.
 37. Sanko JS, McKay M. Impact of simulation-enhanced pharmacology education in prelicensure nursing education. *Nurse Educ*. 2014;42:s32-s7.
 38. Totton N, Lin JF, Julious S, Chowdhury M, Brand A. A review of sample sizes for UK pilot and feasibility studies on the ISRCTN registry from 2013 to 2020. *Pilot and Feasibility Studies*. 2023;9(1).
 39. World Health Organization. Strengthening the health system response to COVID-10: maintaining the delivery of essential health care services while mobilizing the health workforce for the COVID-19 response (18 April 2020) Copenhagen: World Health Organization Regional Office for Europe; 2020 [Available from: <https://iris.who.int/bitstream/handle/10665/332559/WHO-EURO-2020-669-40404-54161-eng.pdf>].
 40. World Health Organization Regional Office for the Western P. Vital roles of nurses and midwives in the Western Pacific Region Manila: World Health Organization; 2020 [Available from: <https://apps.who.int/iris/bitstream/handle/10665/333042/9789290619185-eng.pdf?sequence=1&isAllowed=y>].
 41. Weeks KW, Pontin D, Coben D, Weeks A, Clochesy JM, Rowe D. Decoding the DNA of healthcare numeracy: establishing a mathematics and healthcare numeracy benchmark for nursing. Cardiff: Numeric Generics Authentic World; 2022. Available from: https://safemedicate.com/sm_site/promo/hna.php.
 42. Weeks KW, Coben D, O'Neill D, Jones A, Weeks A, Brown M, et al. Developing and integrating nursing competence through authentic technology-enhanced clinical simulation education: pedagogies for reconceptualising the theory-practice gap. *Nurse Educ Pract*. 2019;37(May):29-38. [Available from: <https://doi.org/10.1016/j.nepr.2019.04.010>]
 43. Kruskal WH, Wallis WA. Use of ranks in one-criterion variance analysis. *Journal of the American Statistical Association*. 1952;47(260):583-621.

44. Mahou FZ, Decormeille G, Changuiti O, Mouhaoui M, Khattabi A. The effects of screen-based simulation on nursing students' acquisition of medication administration and dosage calculation skills: a randomized controlled trial. *BMC Nurs.* 2024;23(1).
45. Mettiäinen S, Luojus K, Salminen S, Koivula M. Web course on medication administration strengthens nursing students' competence prior to graduation. *Nurse Educ Pract.* 2014;14(4):368-73.
46. Sowan AK, Abu Idhail J. Evaluation of an interactive web-based nursing course with streaming videos for medication administration skills. *Int J Med Inform.* 2014;83(8):592-600.
47. Sato SN, Moreno EC, Rubio-Zarapuz A, Dalamitros AA, Yanez-Sepulveda R, Tornero-Aguilera JF, et al. Navigating the new normal: adapting online and distance learning in the post-pandemic era. *Education Sciences.* 2024;14(1):1-25.
48. Bradley CS, Dreifuerst KT, Johnson BK, Loomis A. More than a meme: the Dunning-Kruger effect as an opportunity for positive change in nursing education. *Clinical Simulation in Nursing.* 2022;66:58-65.
49. Heyn LG, Brembo EA, Byermoen KR, Cruaud C, Eide H, Flo J, et al. Exploring facilitation in virtual simulation in nursing education: a scoping review. *PEC Innovation.* 2023;3.
50. Hung CC, Kao HS, Liu HC, Liang HF, Chu TP, Lee BO. Effects of simulation-based learning on nursing students' perceived competence, self-efficacy, and learning satisfaction: A repeat measurement method. *Nurse Educ Today.* 2021 Feb;97:104725.
51. Koukourikos K, Tsaloglidou A, Kourkouta L, Papathanasiou IV, Iliadis C, Fratzana A, Panagiotou A. Simulation in clinical nursing education. *Acta Informatica Medica.* 2021 ;29(1):15.
52. Bae J, Choi M. Learning Outcomes of Education on Visual Thinking Strategies for Healthcare Professionals: A Systematic Review. *Nurse Education in Practice.* 2025;13:104555.